



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07D 249/08, 249/10, 403/10 A61K 31/41	A2	(11) International Publication Number: WO 92/04335 (43) International Publication Date: 19 March 1992 (19.03.92)
<p>(21) International Application Number: PCT/US91/05897</p> <p>(22) International Filing Date: 26 August 1991 (26.08.91)</p> <p>(30) Priority data: 574,314 28 August 1990 (28.08.90) US</p> <p>(60) Parent Application or Grant (63) Related by Continuation US 574,314 (CIP) Filed on 28 August 1990 (28.08.90)</p> <p>(71) Applicant (for all designated States except US): G.D. SEARLE & CO. [US/US]; Corporate Patent Department, P.O. Box 5110, Chicago, IL 60680-5110 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): REITZ, David, B. [US/US]; 14814 Pleasant Ridge Court, Chesterfield, MO 63017 (US). MANNING, Robert, E. [US/US]; 1298 South Mason Road, St. Louis, MO 63131 (US).</p>	<p>(74) Agents: KEANE, J., Timothy et al.; G.D. Searle & Co., Corporate Patent Department, P.O. Box 5110, Chicago, IL 60680-5110 (US).</p> <p>(81) Designated States: AT, AT (European patent), AU, BB, BE (European patent), BF (OAPI patent), BG, BJ (OAPI patent), BR, CA, CF (OAPI patent), CG (OAPI patent), CH, CH (European patent), CI (OAPI patent), CM (OAPI patent), CS, DE, DE (European patent), DK, DK (European patent), ES, ES (European patent), FI, FR (European patent), GA (OAPI patent), GB, GB (European patent), GN (OAPI patent), GR (European patent), HU, IT (European patent), JP, KP, KR, LK, LU, LU (European patent), MC, MG, ML (OAPI patent), MN, MR (OAPI patent), MW, NL, NL (European patent), NO, PL, RO, SD, SE, SE (European patent), SN (OAPI patent), SU*, TD (OAPI patent), TG (OAPI patent), US.</p> <p>Published Without international search report and to be republished upon receipt of that report.</p>	
<p>(54) Title: RENAL-SELECTIVE BIPHENYLALKYL 1H-SUBSTITUTED-1,2,4-TRIAZOLE ANGIOTENSIN II ANTAGONISTS FOR TREATMENT OF HYPERTENSION</p>		
<div style="text-align: center;"> <p>(I)</p> </div>		
<p>(57) Abstract</p> <p>Renal-selective compounds are described which, in one embodiment, are prodrugs preferentially converted in the kidney to compounds capable of blocking angiotensin II (AII) receptors. These prodrugs are conjugates formed from two components, namely, a first component provided by an AII antagonist compound and a second component which is capable of being cleaved from the first component when both components are chemically linked within the conjugate. The two components are chemically linked by a bond which is cleaved selectively in the kidney, for example, by an enzyme. The liberated AII antagonist compound is then available to block AII receptors within the kidney. Conjugates of particular interest are glutamyl derivatives of biphenylmethyl 1H-substituted-1,2,4-triazole compounds, of which N-acetylglutamic acid, 5-[[4'-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]]carbonylhydrazide is an example.</p>		

+ DESIGNATIONS OF "SU"

Any designation of "SU" has effect in the Russian Federation. It is not yet known whether any such designation has effect in other States of the former Soviet Union.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU ⁺	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TC	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark				

**RENAL-SELECTIVE BIPHENYLALKYL 1H-SUBSTITUTED-1,2,4-
TRIAZOLE ANGIOTENSIN II ANTAGONISTS FOR
TREATMENT OF HYPERTENSION**

5 **Related Application**

This a continuation-in-part of U.S. Application
Ser. No. 07/574,314 filed August 28, 1990.

10 **Field of the Invention**

This invention is in the field of cardiovascular
therapeutics and relates to a class of compounds useful in
control of hypertension. Of particular interest is a class
15 of prodrugs of angiotensin II antagonists which, when
selectively hydrolyzed in the kidney, provide hypertension
control.

20 **Background of the Invention**

The renin-angiotensin system is one of the
hormonal mechanisms involved in regulation of
pressure/volume homeostasis and in expression of
hypertension. Activation of the renin-angiotensin cascade
25 begins with renin secretion from the juxtaglomerular
apparatus of the kidney and culminates in the formation of
angiotensin II, an octapeptide which is the primary active
species of this system. Angiotensin II is a potent
vasoconstrictor agent and also produces other physiological
30 effects such as promoting aldosterone secretion, promoting
sodium and fluid retention, inhibiting renin secretion,
increasing sympathetic nervous system activity, increasing
vasopressin secretion, causing positive cardiac inotropic
effect and modulating other hormonal systems.

35 Previous studies have shown that antagonizing
angiotensin II at its receptors is a viable approach to
inhibit the renin-angiotensin system, given the pivotal

role of this octapeptide which mediates the actions of the renin-angiotensin system through interaction with various tissue receptors. There are several known angiotensin II antagonists, most of which are peptidic in nature. Such
5 peptidic compounds are of limited use due to their lack of oral bioavailability or their short duration of action. Also, commercially-available peptidic angiotensin II antagonists (e.g., Saralasin) have a significant residual agonist activity which further limit their therapeutic
10 application.

Non-peptidic compounds with angiotensin II antagonist properties are known. For example, the sodium salt of 2-n-butyl-4-chloro-1-(2-chlorobenzyl)imidazole-5-
15 acetic acid has specific competitive angiotensin II antagonist activity as shown in a series of binding experiments, functional assays and in vivo tests [P. C. Wong et al, J. Pharmacol. Exp. Ther., 247 (1), 1-7 (1988)]. Also, the
20 sodium salt of 2-butyl-4-chloro-1-(2-nitrobenzyl)imidazole-5-acetic acid has specific competitive angiotensin II antagonist activity as shown in a series of binding experiments, functional assays and in vivo tests [A. T. Chiu et al, European J. Pharmacol., 157, 3121 (1988)]. A family of 1-benzylimidazole-5-acetate
25 derivatives has been shown to have competitive angiotensin II antagonist properties [A. T. Chiu et al, J. Pharmacol. Exp. Ther., 250 (3), 867-874 (1989)]. U.S. Patent No. 4,816,463 to Blankey et al describes a family of 4,5,6,7-tetrahydro-1H-imidazo(4,5-c)-tetrahydro-pyridine
30 derivatives useful as antihypertensives, some of which are reported to antagonize the binding of labelled angiotensin II to rat adrenal receptor preparation and thus cause a significant decrease in mean arterial blood pressure in
35 conscious hypertensive rats. EP No. 253,310, published 20 January 1988, describes a series of aralkyl imidazole compounds, including in particular a family of biphenylmethyl substituted imidazoles, as antagonists to

the angiotensin II receptor. EP No. 323,841, published 12 July 1989, describes four classes of angiotensin II antagonists, namely, biphenylmethylpyrroles, biphenylmethylpyrazoles, biphenylmethyl-1,2,3-triazoles and
5 biphenylmethyl 4-substituted-4H-1,2,4-triazoles, including the compound 3,5-dibutyl-4-[(2'-carboxybiphenyl-4-yl)methyl]-4H-1,2,4-triazole. U.S. Patent No. 4,880,804 to Carini et al describes a family of
10 biphenylmethylbenzimidazole compounds as angiotensin II receptor blockers for use in treatment of hypertension and congestive heart failure.

One disadvantage of these angiotensin II antagonist compounds is that the desired hypertension-
15 reducing effect may be offset by hypotension-induced compensatory stimulation of the renin-angiotensin system or stimulation of the sympathetic nervous system, either of which may result in promotion of sodium and water retention. Also, some angiotensin II antagonists may have
20 toxicological effects systemically which preclude their use at doses necessary to be effective in reducing blood pressure.

To avoid such systemic side effects, drugs may
25 be targetted to the kidney by creating a conjugate compound that would be a renal-specific prodrug containing the targetted drug modified with a chemical carrier moiety. Cleavage of the drug from the carrier moiety by enzymes predominantly localized in the kidney releases the drug in
30 the kidney. Gamma glutamyl transpeptidase and acylase are examples of such cleaving enzymes found in the kidney which have been used to cleave a targetted drug from its prodrug carrier within the kidney.

35 Renal targetted prodrugs are known for delivery of a drug selectively to the kidney. For example, the compound L- γ -glutamyl amide of dopamine when administered to dogs was reported to generate dopamine in vivo by

specific enzymatic cleavage by γ -glutamyl transpeptidase [J. J. Kyncl et al, Adv. Biosc., 20, 369-380 (1979)]. In another study, γ -glutamyl and N-acyl- γ -glutamyl derivatives of the anti-bacterial compound sulfamethoxazole were shown to deliver relatively high concentrations of sulfamethoxazole to the kidney which involved enzymatic cleavage of the prodrug by acylamino acid deacylase and γ -glutamyl transpeptidase [M. Orlowski et al, J. Pharmacol. Exp. Ther., 212, 167-172 (1980)]. The N- γ -glutamyl derivatives of 2-, 3-, or 4-aminophenol and p-fluoro-L-phenylalanine have been found to be readily solvolyzed in vitro by γ -glutamyl transpeptidase [S.D.J. Magnan et al, J. Med. Chem., 25, 1018-1021 (1982)]. The hydralazine-like vasodilator 2-hydrazino-5- γ -butylpyridine (which stimulates guanylate cyclase activity) when substituted with the N-acetyl- γ -glutamyl residue resulted in a prodrug which provided selective renal vasodilation [K. G. Hofbauer et al, J. Pharmacol. Exp. Ther., 212, 838-844 (1985)]. The dopamine prodrug γ -L-glutamyl-L-dopa ("gludopa") has been shown to be relatively specific for the kidney and to increase renal blood flow, glomerular filtration and urinary sodium excretion in normal subjects [D. P. Worth et al, Clin. Sci., 69, 207-214 (1985)]. In another study, gludopa was reported to be an effective renal dopamine prodrug whose activity can be blocked by the dopa-decarboxylase inhibitor carbidopa [R. F. Jeffrey et al, Br. J. Clin. Pharmacol., 25, 195-201 (1988)]. A class of 4-ureido derivatives of isoquinolin-3-ol has been investigated for renal specific effects, such as increases in renal vasodilation and renal blood flow [R. M. Kanojia et al, J. Med. Chem., 32, 990-997 (1989)].

BRIEF DESCRIPTION OF THE DRAWING FIGURES

Fig. 1 is a graph showing reduction in mean arterial pressure by intravenous administration of a

conjugate of the invention to rats over a period of four days.

Fig. 2 is a graph showing angiotensin II pressor response by intravenous administration of a conjugate of the invention to rats over a period of four days.

Fig. 3 is a graph showing change in heart rate upon intravenous administration of a conjugate of the invention to rats over a period of four days.

Description of the Invention

Treatment of circulatory disorders, which include cardiovascular disorders, such as chronic hypertension, sodium-retaining disorders, congestive heart failure, cirrhosis and nephrosis, may be accomplished by administering to a susceptible or afflicted subject a therapeutically-effective amount of a renal-selective prodrug capable of causing blood-pressure reducing effects by selective action in the kidney. An advantage of such renal-selective prodrug therapy resides in reduction or avoidance of adverse side effects associated with systemically-acting drugs.

Advantages of a renal-selective antihypertensive compound are several. First, the renal-selective compound is targetted at those pathophysiological mechanisms which occur primarily in the kidney. Second, the regulation of other organ systems is unaffected; thus, normal physiological regulation of other organ systems is maintained. Third, fewer side-effects may be anticipated, since the compound remains inactive until cleaved in the kidneys. Similarly, fewer negative drug-drug interactions may be anticipated. Finally, since a renal-selective accumulation of active compound may occur, which is not dependent on plasma levels of the parent compound, lower

doses of the renal-selective compound compared to active parent compound may be used.

A renal-selective prodrug is provided by a conjugate comprising a residue of an angiotensin II antagonist compound, which conjugate is renal selective. The conjugate will typically comprise a first component and a second component connected together by a cleavable or hydrolyzable bond. The term "renal-selective", as used to characterize a conjugate of the invention, embraces any of the following four pharmacological events: (1) the conjugate is selectively taken up by the kidney and is selectively cleaved in the kidney; (2) the conjugate is not taken up selectively by the kidney, but is selectively cleaved in the kidney; (3) the conjugate is selectively taken up by the kidney and then cleaved in the kidney; or (4) where the conjugate itself is active as an angiotensin II antagonist, the conjugate is selectively taken up by the kidney without cleavage of the hydrolyzable bond.

The first component of a conjugate of the invention is a residue derived from an antagonist compound capable of inhibiting angiotensin II (AII) receptors, especially those AII receptors located in the kidney. The second residue is capable of being cleaved from the first residue preferential-ly. Cleaving of the first and second residues may be accomplished by a variety of mechanisms. For example, the bond may be cleaved by an enzyme in the kidney.

The residue providing the first component may be characterized as the "AII antagonist active" residue. Such "active" residue may be provided by a compound having AII antagonist activity or by a metabolite of such compound having AII antagonist activity. The residue providing the second component may be characterized in being capable of forming a cleavable bond connecting the "active" first residue and the second residue. Such bond is cleavable by

an enzyme located in the kidney. In a preferred embodiment, this cleavable bond is typically a hydrolyzable amide bond, that is, a bond between a carbonyl-terminated moiety and an reactive nitrogen-terminated moiety, such as an amino-terminated moiety, which may be cleaved by enzyme found in the kidney, but which is not cleaved substantially by enzymes located in other organs or tissues of the body. Preferred bond-cleaving enzymes would be found predominantly in the kidney.


10

The conjugate containing the residue of an AII antagonist compound and containing the cleavable fragment or residue may possess AII antagonist activity comparable to, or more than, or less than, the AII antagonist compound which forms the conjugate. In one embodiment of the invention, the conjugate will have AII receptor blocking activity comparable to the AII antagonist component forming the conjugate. In another embodiment of the invention, the conjugate will have AII receptor blocking activity less than the AII receptor blocking activity forming the conjugate. One advantage of such differential activity between the conjugate and the AII antagonist component is that certain side effects associated with non-renal, systemic AII receptor blocking may be avoided or reduced. For example, at least one conjugate of the invention has been found to have very large differential in AII receptor blocking activities between the conjugate and the AII antagonist component forming the conjugate. Such differential activity is advantageous in that therapeutically-effective antihypertensive doses of the conjugate may be administered to give renal-selective AII receptor blocking action resulting from kidney-specific enzyme hydrolysis or metabolism of the conjugate to free the active AII receptor blocker within the kidney. Inasmuch as this renal-selective conjugate has relatively low AII receptor blocking activity, compared to the AII receptor compound forming the conjugate, this conjugate will have fewer adverse side effects associated with

unwanted systemic interaction with non-renal AII receptors
such as found in the vascular bed.

DETAILED DESCRIPTION OF THE INVENTION


The first residue of the conjugate may be selected from any class of compounds, or metabolites thereof, having angiotensin II antagonist activity. An example of one such class of angiotensin II antagonist compounds is provided by a class of biphenylalkyl 1H-substituted-1,2,4-triazole compounds defined by Formula I:



wherein m is a number selected from one to four, inclusive;

wherein each of R¹ through R¹¹ is independently selected from hydrido, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, formyl, alkoxy, aralkyl, aryl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkylcarbonylalkyl, alkoxy carbonyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxy carbonylalkyl, aralkoxy carbonylalkyl, aralkylcarbonyloxyalkyl, mercaptocarbonyl, mercaptothiocarbonyl, mercaptoalkyl, alkoxy carbonyloxy, alkylthio, cycloalkylthio, alkylthiocarbonyl, alkylcarbonylthio, alkylthiocarbonyloxy, alkylthiocarbonylthio, alkylthiothiocarbonyl, alkylthiothiocarbonylthio, arylthio, arylthiocarbonyl, arylcarbonylthio, arylthiocarbonyloxy, arylthiocarbonylthio, arylthiothiocarbonyl, arylthiothiocarbonylthio, aralkylthio, aralkylthiocarbonyl, aralkylcarbonylthio, aralkylthiocarbonyloxy,

aralkylthiocarbonylthio, alkylthiocarbonyl,
 aralkylthiocarbonylthio, mercapto, alkylsulfinyl,
 alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl,
 arylsulfinyl, arylsulfonyl, phthalimido, phthalimidoalkyl,
 5 heteroaryl, heteroarylalkyl, cycloheteroalkyl,
 cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl
 wherein each of said heteroaryl- and cyclohetero-containing
 groups has one or more ring atoms selected from oxygen,
 sulfur and nitrogen atoms, and wherein each of R¹ through
 10 R¹¹ may be further independently selected from amino and
 amido radicals of the formula



15

wherein X is oxygen atom or sulfur atom;

wherein each n is a number independently selected from zero
 to six, inclusive;


20

wherein each of R¹² through R²⁴ is independently selected
 from hydrido, alkyl, cycloalkyl, cyano, amino,
 monoalkylamino, dialkylamino, hydroxyalkyl,
 cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl, and wherein
 25 R¹² and R¹³ taken together, R¹⁴ and R¹⁵ taken together, R¹⁶
 and R¹⁷ taken together, R¹⁹ and R²⁰ taken together and R²¹
 and R²² taken together may each form a heterocyclic group
 having five to seven ring members including the nitrogen
 atom of said amino or amido radical and which heterocyclic
 30 group may further contain one or more hetero atoms as ring
 members selected from oxygen, nitrogen and sulfur atoms and
 which heterocyclic group may be saturated or partially


- unsaturated; wherein R¹² and R¹³ taken together, R¹⁴ and R¹⁵ taken together, R¹⁹ and R²⁰ taken together and R²¹ and R²² taken together may each form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino or amido radical and which aromatic heterocyclic group may further contain one or more hetero atoms as ring atoms selected from oxygen, nitrogen and sulfur atoms;
- and wherein each of R³ through R¹¹ may be further independently selected from hydroxy and from acidic moieties of the formula




- wherein n is a number selected from zero through three, inclusive, and wherein A is an acidic group selected to contain at least one acidic hydrogen atom, and the amide, ester and salt derivatives of said acidic moieties;
- wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, aryl, aralkyl and heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms;
- and wherein any of the foregoing R¹ through R²⁴, Y and A groups having a substitutable position may be substituted with one or more groups selected from hydroxy, alkyl, alkenyl, alkynyl, aralkyl, hydroxyalkyl, trifluoromethyl, difluoroalkyl, oxo, alkoxy, aryloxy, aralkoxy, aralkylthio, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aroyl, cycloalkenyl, cyano, cyanoamino, nitro, alkylcarbonyloxy, alkoxycarbonyloxy, alkylcarbonyl, alkoxycarbonyl, carboxyl, mercapto, mercaptocarbonyl, alkylthio, arylthio, alkylthiocarbonyl, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms, and amino and amido radicals of the formula



- wherein X is selected from oxygen atom and sulfur atom;
 5 wherein each of R²⁵, R²⁶, R²⁷, R²⁸ and R²⁹ is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl, DR³⁰ and



- 10 wherein D is selected from oxygen atom and sulfur atom and R³⁰ is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl and aryl; wherein each of R²⁵, R²⁶, R²⁷, R²⁸, R²⁹, R³¹ and R³² is independently selected
 15 from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, alkylcarbonyl, alkoxycarbonyl, carboxyl, alkylsulfinyl, alkylsulfonyl, arylsulfinyl, arylsulfonyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl, and wherein each of
 20 R²⁵, R²⁶, R²⁷, R²⁸, R²⁹, R³¹ and R³² is further independently selected from amino and amido radicals of the formula



- 25 wherein X is oxygen atom or sulfur atom;

- wherein each of R³³, R³⁴, R³⁵, R³⁶, R³⁷ and R³⁸ is
 30 independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, haloalkylsulfinyl,

haloalkylsulfonyl, aralkyl and aryl, and wherein R²⁶ and R²⁷ taken together and R²⁸ and R²⁹ taken together may each form a heterocyclic group having five to seven ring members including the nitrogen atom of said amino or amido radical, which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms and which heterocyclic group may be saturated or partially unsaturated; wherein R²⁶ and R²⁷ taken together and R³¹ and R³² taken together may each form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino or amido radical and which aromatic heterocyclic group may further contain one or more hetero atoms as ring atoms selected from oxygen, nitrogen and sulfur atoms;

with the proviso that at least one of said R¹ through R²⁴, Y and A substituents contains a terminal primary or secondary amino moiety or a moiety convertible to a primary or secondary amino moiety.

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

Conjugates of the invention are therapeutically effective in treatment of cardiovascular disorders by acting directly on, or by providing cleavable components selected from Formula I compounds which act directly on, or as antagonists to, or as blockers of, the angiotensin II (AII) receptor. Thus, conjugates of Formula I would be therapeutically effective in treatment of cardiovascular disorders or would be precursors to, or prodrugs of, therapeutically-effective compounds.

Preferred compounds of Formula I, from which a cleavable component may be selected, are all characterized in having a substituent, other than hydrido, at each of the three- and five-positions of the triazole ring. Such substituents are selected from the aforementioned R¹ and R²

groups. Also especially useful are compounds having one of the R^1 and R^2 substituents selected from alkylcarbonyl, monoalkoxyalkyl, dialkoxyalkyl and difluoroalkyl groups. When the selected substituent for R^1 and R^2 is

5 difluoroalkyl, then it is particularly useful for both of the fluoro atoms of the difluoroalkyl group to be substituted on the difluoroalkyl group carbon atom attached at the R^1 or R^2 positions of the triazole ring. Such difluoroalkyl group may be characterized as an "alpha-

10 carbon difluoro-substituted difluoroalkyl group", or as an "alpha,alpha-difluoro-substituted alkyl group". When the selected substituent for R^1 or R^2 is monoalkoxyalkyl or dialkoxyalkyl, then it is particularly useful for the single alkoxy group or the two alkoxy groups, respectively,

15 to be substituted on the carbon atom of the selected substituent which is attached at the R^1 or R^2 positions of the triazole ring. Such alkoxyalkyl groups may be characterized as "alpha-carbon monoalkoxy- or dialkoxy-

20 substituted alkoxyalkyl groups", respectively, or "alpha-monoalkoxy-substituted or alpha,alpha-dialkoxy-substituted alkyl groups", respectively. When the selected substituent is alkylcarbonyl, then it is particularly useful for the carbonyl group to be attached at the R^1 or R^2 positions of the triazole ring. Such alkylcarbonyl group may be

25 characterized as an "alpha-oxo-substituted alkyl group", and may be exemplified by the substituents 1-oxoethyl, 1-oxopropyl and 1-oxobutyl. Where compounds of Formula I contain any of these above-mentioned particularly-useful alpha-carbon substituted R^1 or R^2 groups at the triazole


30 ring three- or five-position, it is preferred that such particularly-useful group be attached at the three-position, that is, as an R^1 substituent.

35 The phrase "acidic group selected to contain at least one acidic hydrogen atom", as used to define the $-Y_nA$ moiety, is intended to embrace chemical groups which, when attached to any of the R^3 through R^{11} positions of Formula I, confers acidic character to the compound of Formula I.

"Acidic character" means proton-donor capability, that is, the capacity of the compound of Formula I to be a proton donor in the presence of a proton-receiving substance such as water. Typically, the acidic group should be selected to have proton-donor capability such that the product compound of Formula I has a pK_a in a range from about one to about twelve. More typically, the Formula I compound would have a pK_a in a range from about two to about seven. An example of an acidic group containing at least one acidic hydrogen atom is carboxyl group ($-COOH$). Where n is zero and A is $-COOH$, in the $-Y_nA$ moiety, such carboxyl group would be attached directly to one of the R^3 through R^{11} positions. The Formula I compound may have one $-Y_nA$ moiety attached at one of the R^3 through R^{11} positions, or may have a plurality of such $-Y_nA$ moieties attached at more than one of the R^3 through R^{11} positions, up to a maximum of nine such $-Y_nA$ moieties. There are many examples of acidic groups other than carboxyl group, selectable to contain at least one acidic hydrogen atom. Such other acidic groups may be collectively referred to as "bioisosteres of carboxylic acid" or referred to as "acidic bioisosteres". Specific examples of such acidic bioisosteres are described hereinafter. Compounds of Formula I having the $-Y_nA$ moiety attached at one of positions R^5 , R^6 , R^8 and R^9 would be expected to have preferred properties, while attachment at R^5 or R^9 would be more preferred.

A preferred class of compounds within the subclass defined by Formula I consists of those compounds wherein m is one; wherein each of R^1 through R^{11} is independently selected from alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, aralkyl, aryl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxycarbonylalkyl, aralkoxycarbonylalkyl,

aralkylcarbonyloxyalkyl, mercaptocarbonyl,
 mercaptothiocarbonyl, mercaptoalkyl, alkoxycarbonyloxy,
 alkylthio, cycloalkylthio, alkylthiocarbonyl,
 alkylcarbonylthio, alkylthiocarbonyloxy, alkylthio-
 5 carbonylthio, alkylthiothiocarbonyl, alkylthiothio-
 carbonylthio, arylthio, arylthiocarbonyl, arylcarbonylthio,
 arylthiocarbonyloxy, arylthiocarbonylthio,
 arylthiothiocarbonyl, arylthiothiocarbonylthio,
 aralkylthio, aralkylthiocarbonyl, aralkylcarbonylthio,
 10 aralkylthiocarbonyloxy, aralkylthiocarbonylthio,
 aralkylthiocarbonyl, aralkylthiocarbonylthio, mercapto,
 alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl,
 aralkylsulfonyl, arylsulfinyl, arylsulfonyl, phthalimido,
 phthalimidoalkyl, heteroaryl, heteroarylalkyl,
 15 cycloheteroalkyl, cycloheteroalkylalkyl and
 cycloheteroalkylcarbonylalkyl wherein each of said
 heteroaryl- and cycloheteroalkyl-containing groups has
 one or more hetero ring atoms selected from oxygen, sulfur
 and nitrogen atoms, and wherein each of R¹ through R¹¹ may
 20 be further independently selected from amino and amido
 radicals of the formula



25

wherein X is selected from oxygen atom or sulfur atom;

wherein each n is a number independently selected from zero
 to six, inclusive;

30

wherein each of R¹² through R²⁴ is independently selected
 from hydrido, alkyl, cycloalkyl, cyano, amino,

monoalkylamino, dialkylamino, hydroxyalkyl,
cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;


- and wherein each of R³ through R¹¹ may be further
5 independently selected from hydroxy and from acidic
moieties of the formula



- 10 wherein n is a number selected from zero through three,
inclusive; wherein A is an acidic group selected from acids
containing one or more atoms selected from oxygen, sulfur,
phosphorus and nitrogen atoms, and wherein said acidic
group is selected to contain at least one acidic hydrogen
15 atom, and the amide, ester and salt derivatives of said
acidic moieties; wherein Y is a spacer group independently
selected from one or more of alkyl, cycloalkyl,
cycloalkylalkyl, alkenyl, alkynyl, aryl, aralkyl and
heteroaryl having one or more ring atoms selected from
20 oxygen, sulfur and nitrogen atoms;


- and wherein any of the foregoing R¹ through R²⁴, Y and A
groups having a substitutable position may be substituted
with one or more groups selected from alkyl, alkenyl,
25 aralkyl, hydroxyalkyl, trifluoromethyl, difluoroalkyl,
alkoxy, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl,
alkoxycarbonyl, carboxyl, mercaptocarbonyl, alkylthio,
alkylthiocarbonyl, and amino and amido radicals of the
formula

30




wherein X is selected from oxygen atom and sulfur atom;
wherein each of R²⁵, R²⁶, R²⁷, R²⁸ and R²⁹ is selected from

hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl,
and DR³⁰ and



5

wherein D is selected from oxygen atom and sulfur atom,
and R³⁰ is selected from hydrido, alkyl, cycloalkyl,
cycloalkylalkyl, aralkyl and aryl; wherein each of R²⁵,
R²⁶, R²⁷, R²⁸, R²⁹, R³¹ and R³² is independently selected
10 from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl,
haloalkyl, cycloalkylalkyl, alkoxyalkyl, alkanoyl,
alkoxycarbonyl, carboxyl, haloalkylsulfinyl,
haloalkylsulfonyl, aralkyl and aryl, and wherein each of
R²⁵, R²⁶, R²⁷, R²⁸, R²⁹, R³¹ and R³² is further
15 independently selected from amino and amido radicals of the
formula



20 wherein X is selected from oxygen atom or sulfur atom;

wherein each of R³³ through R³⁸ is independently selected
from hydrido, alkyl, cycloalkyl, cyano, amino,
monoalkylamino, dialkylamino, hydroxyalkyl,
25 cycloalkylalkyl, alkoxyalkyl, haloalkylsulfinyl,
haloalkylsulfonyl, aralkyl and aryl;

with the proviso that at least one of said R¹ through R²⁴,
Y and A substituents contains a terminal primary or
30 secondary amino moiety or a moiety convertible to a primary
or secondary amino moiety;

or a tautomer thereof or a pharmaceutically-acceptable salt
thereof.

A more preferred class of compounds within the sub-class defined by Formula I consists of those compounds wherein m is one; wherein each of R¹ through R¹¹ is


5 independently selected from alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, aralkyl, aryl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl,

10 carboxyalkyl, alkylcarbonyloxy, mercaptocarbonyl, alkoxy carbonyloxy, alkylcarbonyloxyalkyl, alkoxy carbonylalkyl, aralkoxy carbonylalkyl, aralkylcarbonyloxyalkyl, alkylthio, cycloalkylthio, arylthio, aralkylthio, aralkylthiocarbonylthio, mercapto,

15 alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl wherein each of said

20 heteroaryl- and cycloheteroalkyl-containing groups has one or more hetero ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R¹ through R¹¹ may be further independently selected from amino and amido radicals of the formula

25



wherein X is selected from oxygen atom or sulfur atom;

30

wherein each n is a number independently selected from zero to six, inclusive;

wherein each of R¹² through R²⁴ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

and wherein each of R³ through R¹¹ may be an further independently selected from hydroxy and from acidic moieties of the formula


10



wherein n is a number selected from zero through three, inclusive;

15


wherein A is selected from carboxylic acid and bioisosteres of carboxylic acid selected from



20

wherein each W is independently selected from oxygen atom, sulfur atom and NR⁴³; wherein each of R³⁹, R⁴⁰, R⁴¹, R⁴² and R⁴³ is independently selected from hydrido, alkyl, haloalkyl, haloalkylsulfonyl, haloalkylcarbonyl, cycloalkyl, cycloalkylalkyl, aryl and aralkyl; wherein each of R³⁹, R⁴⁰, R⁴¹ and R⁴² may be further independently selected from amino radical of the formula

25



30

wherein each of R⁴⁴ and R⁴⁵ is independently selected from hydrido, alkyl, cycloalkyl, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl, and wherein R⁴⁴ and R⁴⁵ taken together may form a heterocyclic group

having five to seven ring members including the nitrogen atom of said amino radical, which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms and which

5 heterocyclic group may be saturated or partially unsaturated; wherein R^{44} and R^{45} taken together may form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino radical and which aromatic heterocyclic group may further contain one or more

10 hetero atoms as ring atoms selected from oxygen, nitrogen and sulfur atoms; wherein each of R^{44} and R^{45} may be further independently selected from hydroxy, alkoxy, alkylthio, aryloxy, arylthio, aralkylthio and aralkoxy; and the amide, ester and salt derivatives of said acidic

15 groups;

wherein said bioisostere of carboxylic acid may be further selected from heterocyclic acidic groups consisting of heterocyclic rings of four to about nine ring members,

20 which heterocyclic ring contains at least one hetero atom selected from oxygen, sulfur and nitrogen atoms, which heterocyclic ring may be saturated, fully unsaturated or partially unsaturated, and which heterocyclic ring may be attached at a single position selected from R^3 through R^{11}


25 or may be attached at any two adjacent positions selected from R^3 through R^{11} so as to form a fused-ring system with one of the phenyl rings of the biphenyl moiety of Formula I; and the amide, ester and salt derivatives of said heterocyclic acidic groups;

30 wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, aryl and aralkyl;


35 and wherein any of the foregoing R^1 through R^{24} , Y and A groups having a substitutable position may be substituted by one or more groups selected from alkyl, difluoroalkyl, alkenyl, aralkyl, hydroxyalkyl, trifluoromethyl, alkoxy,

aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, carboxyl, mercaptocarbonyl, alkylthio, alkylthiocarbonyl, and amino and amido radicals of the formula

5



wherein X is selected from oxygen atom and sulfur atom;
 wherein each of R²⁵, R²⁶, R²⁷, R²⁸ and R²⁹ is selected from
 10 hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl
 and DR³⁰ and



15 wherein D is selected from oxygen atom and sulfur atom,
 wherein R³⁰ is selected from hydrido, alkyl, cycloalkyl,
 cycloalkylalkyl, aralkyl and aryl;


wherein each of R²⁵, R²⁶, R²⁷, R²⁸, R²⁹, R³¹ and R³² is
 20 independently selected from hydrido, alkyl, cycloalkyl,
 cyano, hydroxyalkyl, haloalkyl, cycloalkylalkyl,
 alkoxyalkyl, alkanoyl, alkoxy carbonyl, carboxyl,
 haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl;

25 with the proviso that at least one of said R¹ through R²⁴,
 Y and A substituents contains a terminal primary or
 secondary amino moiety or a moiety convertible to a primary
 or secondary amino moiety;

30 or a tautomer thereof or a pharmaceutically-acceptable salt
 thereof.

An even more preferred class of compounds within
 the sub-class defined by Formula I consists of those

compounds wherein m is one; wherein each of R¹ and R² is independently selected from alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, aralkyl, aryl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxycarbonylalkyl, aralkoxycarbonylalkyl, aralkylcarbonyloxyalkyl, mercaptocarbonyl, mercaptoalkyl, alkoxy carbonyloxy, alkylthio, cycloalkylthio, arylthio, aralkylthio, mercapto, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl wherein each of said heteroaryl- and cycloheteroalkyl-containing groups has one or more hetero ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R¹ through R¹¹ may be further independently selected from amino and amido radicals of the formula



wherein X is selected from oxygen atom and sulfur atom;

wherein each n is a number independently selected from zero to six, inclusive;

wherein each of R¹² through R²⁴ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino,


monoalkylamino, dialkylamino, hydroxyalkyl,
cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

- wherein each of R^3 through R^{11} is independently selected
 5 from hydrido, hydroxy, alkyl, hydroxyalkyl, halo,
 haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, aralkyl,
 aryl, aroyl, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl,
 alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl,
 cycloalkynyl, cyano, nitro, carboxyl, alkylthio,
 10 aralkylthio, mercapto, alkylsulfinyl, alkylsulfonyl,
 aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl,
 arylsulfonyl and heteroaryl having one or more ring atoms
 selected from oxygen, sulfur and nitrogen atoms;


- 15 and wherein each of R^3 through R^{11} may be an acidic moiety
 further independently selected from acidic moieties of the
 formula



- 20 wherein n is a number selected from zero through three,
 inclusive; wherein A is selected from carboxylic acid and
 bioisosteres of carboxylic acid selected from



- 25 wherein each W is independently selected from oxygen atom,
 sulfur atom and NR^{43} ; wherein each of R^{39} , R^{42} and R^{43} is
 independently selected from hydrido, alkyl, haloalkyl,
 haloalkylsulfonyl, haloalkylcarbonyl, cycloalkyl,
 30 cycloalkylalkyl, aryl and aralkyl; wherein each of R^{39} and
 R^{42} may be further independently selected from amino
 radical of the formula



wherein each of R⁴⁴ and R⁴⁵ is independently selected from hydrido, alkyl, cycloalkyl, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl, and wherein R⁴⁴ and R⁴⁵ taken together may form a heterocyclic group having five to seven ring members including the nitrogen atom of said amino radical, which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms, and which heterocyclic group may be saturated or partially unsaturated; wherein R⁴⁴ and R⁴⁵ taken together may form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino radical and which aromatic heterocyclic group may further contain one or more hetero atoms as ring atoms selected from oxygen, nitrogen and sulfur atoms; and the amide, ester and salt derivatives of said acidic groups; wherein said bioisostere of carboxylic acid may be further selected from heterocyclic acidic groups consisting of heterocyclic rings of four to about nine ring members, which ring contains at least one hetero atom, selected from oxygen, sulfur and nitrogen atoms, which heterocyclic ring may be saturated, fully unsaturated or partially unsaturated, and which heterocyclic ring may be attached at a single position selected from R³ through R¹¹ or may be attached at any two adjacent positions selected from R³ through R¹¹ so as to form a fused-ring system with one of the phenyl rings of Formula I; and the amide, ester and salt derivatives of said heterocyclic acidic groups;

wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, aryl and aralkyl;


wherein each of R¹ through R¹¹, Y and A independently may be substituted at any substitutable position with one or more groups selected from alkyl, cycloalkyl, cycloalkylalkyl, hydroxy, oxo, trifluoromethyl,

difluoroalkyl, alkoxycarbonyl, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, aryl, aralkyl, alkoxy, aryloxy and aralkoxy;

- 5 with the proviso that at least one of said R^1 through R^{24} ,
Y and A substituents contains a terminal primary or
secondary amino moiety or a moiety convertible to a primary
or secondary amino moiety;
- 10 or a tautomer thereof or a pharmaceutically-acceptable salt
thereof.

- A highly preferred class of compounds within the
sub-class defined by Formula I consists of those compounds
15 wherein m is one; wherein each of R^1 and R^2 is
independently selected from alkyl, hydroxyalkyl, halo,
haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, aralkyl,
aryl, benzoyl, phenoxy, phenoxyalkyl, phenalkyloxy,
phenylthio, phenalkylthio, aralkoxy, alkoxyalkyl,
20 alkylcarbonyl, alkoxycarbonyl, alkenyl, cycloalkenyl,
alkynyl, cyano, nitro, carboxyl, carboxyalkyl,
alkylcarbonyloxy, alkylcarbonyloxyalkyl,
alkoxycarbonylalkyl, aralkoxycarbonylalkyl,
aralkylcarbonyloxyalkyl, mercaptocarbonyl, mercaptoalkyl,
25 alkoxycarbonyloxy, alkylthio, cycloalkylthio, phthalimido,
phthalimidoalkyl, heteroaryl, heteroarylalkyl,
cycloheteroalkyl, cycloheteroalkylalkyl and
cycloheteroalkylcarbonylalkyl wherein each of said
heteroaryl- and cycloheteroalkyl-containing groups has one
30 or more hetero ring atoms selected from oxygen, sulfur and
nitrogen atoms, and wherein each of R^1 through R^{11} may be
further independently selected from amino and amido
radicals of the formula

27



wherein X is selected from oxygen atom and sulfur atom;

- 5 wherein each n is a number independently selected from zero to six, inclusive;

wherein each of R¹² through R²⁴ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino,
 10 hydroxyalkyl, alkoxyalkyl, phenalkyl and phenyl;


wherein each of R³ through R¹¹ is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, phenalkyl,
 15 phenyl, benzoyl, phenoxy, phenalkyloxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, alkenyl, cyano, nitro, carboxyl, alkylthio, mercapto and heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms;

20 and wherein each of R³ through R¹¹ may be an acidic moiety further independently selected from acidic moieties of the formula




25

wherein n is a number selected from zero through two, inclusive; wherein A is selected from carboxylic acid and bioisosteres of carboxylic acid selected from



30

wherein each W is independently selected from oxygen atom, sulfur atom and NR^{43} ; wherein each of R^{39} , R^{42} and R^{43} is independently selected from hydrido, alkyl, haloalkyl, haloalkylsulfonyl, haloalkylcarbonyl, cycloalkyl, phenyl and benzyl; wherein each of R^{39} and R^{42} may be further independently selected from amino radical of the formula



10

wherein each of R^{44} and R^{45} is independently selected from hydrido, alkyl, cycloalkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, benzyl and phenyl; and the amide, ester and salt derivatives of said acidic groups;

15

wherein said bioisostere of carboxylic acid may be further selected from heterocyclic acidic groups consisting of heterocyclic rings of four to about nine ring members, which ring contains at least one hetero atom, selected from oxygen, sulfur and nitrogen atoms, which heterocyclic ring may be saturated, fully unsaturated or partially unsaturated, and which heterocyclic ring may be attached at a single position selected from R^3 through R^{11} or may be attached at any two adjacent positions selected from R^3 through R^{11} so as to form a fused-ring system with one of the phenyl rings of the biphenyl moiety of Formula I; and the amide, ester and salt derivatives of said heterocyclic acidic groups;

30

wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, phenyl, phenalkyl and aralkyl;

35

wherein each of R^1 through R^{11} , Y and A and independently may be substituted at any substitutable position with one or more groups selected from alkyl, cycloalkyl,

cycloalkylalkyl, hydroxy, oxo, trifluoromethyl, difluoroalkyl, alkoxycarbonyl, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, aryl, aralkyl, alkoxy, aryloxy and aralkoxy;


5

with the proviso that at least one of said R^1 through R^{24} , Y and A substituents contains a terminal primary or secondary amino moiety or a moiety convertible to a primary or secondary amino moiety;

10

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

An even more highly preferred class of compounds within Formula I consists of those compounds wherein m is one; wherein each of R^1 and R^2 is independently selected from alkyl, aminoalkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, aralkyl, aryl, benzoyl, phenoxy, phenoxyalkyl, phenalkyloxy, phenylthio, phenalkylthio, aralkoxy, alkoxyalkyl, acetyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, mercaptoalkyl, mercaptocarbonyl, alkoxycarbonyloxy, alkylcarbonyloxyalkyl, alkoxycarbonylalkyl, aralkoxycarbonylalkyl, aralkylcarbonyloxyalkyl, phthalimido, phthalimidoalkyl, imidazoalkyl, tetrazole, tetrazolealkyl, alkylthio, cycloalkylthio, and amino and amido radicals of the formula



wherein X is selected from oxygen atom and sulfur atom;


wherein each n is a number independently selected from zero to six, inclusive;

5


wherein each of R¹² through R²⁴ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, hydroxyalkyl, alkoxyalkyl, phenalkyl and phenyl;

10 wherein each of R³ through R¹¹ is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, phenalkyl, phenyl, benzoyl, phenoxy, phenalkyloxy, alkoxyalkyl, acetyl, alkoxycarbonyl, alkenyl, cyano, nitro, carboxyl, 15 alkylthio and mercapto;

and wherein each of R³ through R¹¹ may be an acidic moiety further independently selected from acidic moieties consisting of CO₂H, CO₂CH₃, SH, CH₂SH, C₂H₄SH, PO₃H₂, 20 NHSO₂CF₃, NHSO₂C₆F₅, SO₃H, CONHNH₂, CONHNHSO₂CF₃, CONHOCH₃, CONHOC₂H₅, CONHCF₃, OH, CH₂OH, C₂H₄OH, OPO₃H₂, OSO₃H ,



wherein each of R⁴⁶, R⁴⁷ and R⁴⁸ is independently selected
 5 from H, Cl, CN, NO₂, CF₃, C₂F₅, C₃F₇, CHF₂, CH₂F, CO₂CH₃,
 CO₂C₂H₅, SO₂CH₃, SO₂CF₃ and SO₂C₆F₅; wherein Z is selected
 from O, S, NR⁴⁹ and CH₂; wherein R⁴⁹ is selected from
 hydrido, CH₃ and CH₂C₆H₅; and wherein said acidic moiety
 may be a heterocyclic acidic group attached at any two
 10 adjacent positions of R³ through R¹¹ so as to form a fused
 ring system so as to include one of the phenyl rings of the
 biphenyl moiety of Formula I, said biphenyl fused ring
 system selected from




and the esters, amides and salts of said acidic moieties;






- 5 with the proviso that at least one of said R^1 through R^{24} substituents contains a terminal primary or secondary amino moiety or a moiety convertible to a primary or secondary amino moiety;
- 10 or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

- A class of compounds of particular interest within the sub-class defined by Formula I consists of those
- 15 compounds wherein m is one; wherein each of R^1 and R^2 is independently selected from amino, aminomethyl, aminoethyl, aminopropyl, CH_2OH , CH_2OCOCH_3 , CH_2Cl , Cl , CH_2OCH_3 , $CH_2OCH(CH_3)_2$, I , CHO ,


CH_2CO_2H , $CH(CH_3)CO_2H$, NO_2 , Cl ,



- 20 $-CH_2OCOCH_2CH_2-$ (with a benzene ring attached to the CH_2 group), $-CO_2CH_3$, $-CONH_2$, $-CONHCH_3$, $CON(CH_3)_2$,

- $-\text{CH}_2-\text{NHCO}_2\text{C}_2\text{H}_5$, $-\text{CH}_2\text{NHCO}_2-$ , $-\text{CH}_2\text{NHCO}_2\text{CH}_3$, $-\text{CH}_2\text{NHCO}_2\text{C}_3\text{H}_7$,
 $-\text{CH}_2\text{NHCO}_2\text{CH}_2(\text{CH}_3)_2$, $-\text{CH}_2\text{NHCO}_2\text{C}_4\text{H}_9$, CH_2NHCO_2 -adamantyl,
 $-\text{CH}_2\text{NHCO}_2$ -(1-naphthyl), $-\text{CH}_2\text{NHCONHCH}_3$, $-\text{CH}_2\text{NHCONHC}_2\text{H}_5$,
 $-\text{CH}_2\text{NHCONHC}_3\text{H}_7$, $-\text{CH}_2\text{NHCONHC}_4\text{H}_9$, $-\text{CH}_2\text{NHCONHCH}(\text{CH}_3)_2$,
 5 $-\text{CH}_2\text{NHCONH}(1\text{-naphthyl})$, $-\text{CH}_2\text{NHCONH}(1\text{-adamantyl})$, CO_2H ,
 $-\text{CH}_2\text{CH}_2\text{CO}-\text{N}$ , $-\text{CH}_2\text{CH}_2\text{CO}-\text{N}$ , $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CO}_2\text{H}$,
 $-\text{CH}_2\text{CH}_2\text{F}$, $-\text{CH}_2\text{OCONHCH}_3$, $-\text{CH}_2\text{OCSNHCH}_3$, $-\text{CH}_2\text{NHC}_3\text{H}_7$,
 $-\text{CH}_2\text{CH}_2\text{CH}_2\text{F}$, $-\text{CH}_2\text{ONO}_2$, $-\text{CH}_2-\text{N}$ , $-\text{CH}_2\text{SH}$, $-\text{CH}_2\text{O}-$ ,
 H, Cl, NO_2 , CF_3 , CH_2OH , Br, F, I, methyl, ethyl, n-propyl,
 10 isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-
 pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl,
 cyclohexyl, cyclohexylmethyl, 1-oxoethyl, 1-oxopropyl, 1-
 oxobutyl, 1-oxopentyl, 1,1-dimethoxypropyl, 1,1-
 dimethoxybutyl, 1,1-dimethoxypentyl, hydroxyalkyl, halo,
 15 difluoromethyl, 1,1-difluoroethyl, 1,1-difluoropropyl, 1,1-
 difluorobutyl and 1,1-difluoropentyl; wherein each of R^3
 through R^{11} is hydrido, with the proviso that at least one
 of R^5 , R^6 , R^8 and R^9 is an acidic group selected from CO_2H ,
 SH, PO_3H_2 , SO_3H , CONHNH_2 , $\text{CONHNHSO}_2\text{CF}_3$, OH,

20



wherein each of R^{46} and R^{47} is independently selected from
 Cl, CN, NO_2 , CF_3 , CO_2CH_3 and SO_2CF_3 ;


25

with the proviso that at least one of said R^1 through R^{11}
 substituents contains a terminal primary or secondary amino
 moiety or a moiety convertible to a primary or secondary
 amino moiety;

30

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

- A class of compounds of more particular interest within the sub-class defined by Formula I consists of those compounds wherein m is one; wherein R^1 is selected from amino, aminomethyl, aminoethyl, aminopropyl, methyl, ethyl, n -propyl, isopropyl, n -butyl, sec-butyl, isobutyl, tert-butyl, n -pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclohexyl, cyclohexylmethyl, 1-oxoethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl, 1,1-dimethoxypropyl, 1,1-dimethoxybutyl, 1,1-dimethoxypentyl, hydroxyalkyl, halo, difluoromethyl, 1,1-difluoroethyl, 1,1-difluoropropyl, 1,1-difluorobutyl and 1,1-difluoropentyl; wherein R^2 is selected from ethyl, n -propyl, isopropyl, n -butyl, sec-butyl, isobutyl, 4-methylbutyl, tert-butyl, n -pentyl and neopentyl; wherein each of R^3 , R^4 , R^6 , R^7 , R^8 , R^{10} , and R^{11} is hydrido; wherein one of R^5 and R^9 is hydrido and the other of R^5 and R^9 is an acidic group selected from CO_2H , SH , PO_3H_2 , SO_3H , $CONHNH_2$, $CONHNHSO_2CF_3$, OH ,




- wherein each of R^{46} and R^{47} is independently selected from Cl , CN , NO_2 , CF_3 , CO_2CH_3 and SO_2CF_3 ;

- with the proviso that at least one of said R^1 through R^{11} substituents contains a terminal primary or secondary amino moiety or a moiety convertible to a primary or secondary amino moiety;

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

The second component of a conjugate of the invention is provided by a residue which forms a kidney-enzyme-cleavable amide bond with the residue of the first-component AII antagonist compound. Such residue is preferably selected from a class of compounds of Formula II:



- 10 wherein each of R^{50} and R^{51} may be independently selected from hydrido, alkylcarbonyl, alkoxy carbonyl, alkoxyalkyl, hydroxyalkyl and haloalkyl; and wherein G is selected from hydroxyl, halo, mercapto, $-\text{OR}^{52}$, $-\text{SR}^{53}$ and $-\text{NR}^{54}$ with each of R^{52} , R^{53} and R^{54} independently
- 15 selected from alkyl; and wherein R^{54} may be further selected from hydrido; with the proviso that said cleavable bond is within an amide group formed between said first and second residues, wherein said first residue has a terminal primary or second amino moiety
- 20 provided by one of said R^1 through R^{11} substituents of said Formula I compound or provided by a linker group attached to one of said R^1 - R^{11} substituents of said Formula I, and wherein said second residue has a carbonyl moiety attached at the gamma-position carbon of
- 25 said Formula II compound, whereby said amide bond is formed from said first residue amino moiety and said second residue carbonyl moiety.

More preferred are compounds of Formula II wherein each G is hydroxy.


A more highly preferred class of compounds within Formula II consists of those compounds wherein each

G is hydroxy; wherein R⁵⁰ is hydrido; and wherein R⁵¹ is

selected from $\begin{array}{c} \text{O} \\ \parallel \\ -\text{CR}^{55} \end{array}$

wherein R⁵⁵ is selected from methyl, ethyl, n-propyl,
 5 isopropyl, n-butyl, sec-butyl, iso-butyl, tert-butyl, n-pentyl, neopentyl, n-hexyl and chloromethyl.

A most highly preferred compound of Formula II
 is N-acetyl- γ -glutamic acid which provides a residue for
 10 the second component of a conjugate of the invention as shown below:




15 The phrase "terminal primary or secondary amino moiety or a moiety convertible to a primary or secondary amino terminal moiety" characterizes a structural requirement for selection of a suitable angiotensin II antagonist compound as the "active" first residue of a
 20 conjugate of the invention. Such terminal amino moiety must be available to react with a terminal carboxylic moiety of the cleavable second residue to form a kidney-enzyme-specific hydrolyzable bond.

25 In one embodiment of the invention, the first component used to form a conjugate of the invention provides a first residue derived from an AII antagonist compound containing a terminal primary or secondary amino moiety. Examples of such terminal amino moiety are amino
 30 and linear or branched aminoalkyl moieties containing linear or branched alkyl groups such as aminomethyl, aminoethyl, aminopropyl, aminoisopropyl, aminobutyl, aminosecbutyl, aminoisobutyl, aminotertbutyl, aminopentyl, aminoisopentyl and aminoneopentyl.


In another embodiment of the invention, the first component used to form the conjugate of the invention provides a first residue derived from an AII antagonist compound containing a moiety convertible to a primary or secondary amino terminal moiety. An example of a moiety convertible to an amino terminal moiety is a carboxylic acid group reacted with hydrazine so as to convert the acid moiety to carboxylic acid hydrazide. The hydrazide moiety thus contains the terminal amino moiety which may then be further reacted with the carboxylic acid containing residue of the second component to form a hydrolyzable amide bond. Such hydrazide moiety thus constitutes a "linker" group between the first and second components of a conjugate of the invention.

Suitable linker groups may be provided by a class of diamino-terminated linker groups based on hydrazine as defined by Formula III:

20



wherein each of R²⁰⁰ and R²⁰¹ may be independently selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, alkoxyalkyl, hydroxyalkyl, aralkyl, aryl, haloalkyl, amino, 25 monoalkylamino, dialkylamino, cyanoamino, carboxyalkyl, alkylsulfinyl, alkylsulfonyl, arylsulfinyl and arylsulfonyl; and wherein n is zero or a number selected from three through seven, inclusive. In Table I there is shown a class of specific examples of diamino-terminated linker groups within Formula III, identified as Linker Nos. 1-73. These linker groups would be suitable to form a conjugate between a carbonyl moiety of an AII antagonist (designated as "I") and a carbonyl moiety of a carbonyl terminated 30 second residue such as the carbonyl moiety attached to the gamma carbon of a glutamyl residue (designated as "T").

TABLE I

I = inhibitor

T = acetyl- γ -glutamyl

LINKER NO.	n	R ²⁰⁰	R ²⁰¹
1	0	H	H
2	0	CH ₃	H
3	0	C ₂ H ₅	H
4	0	C ₃ H ₇	H
5	0	CH(CH ₃) ₂	H
6	0	C ₄ H ₉	H
7	0	CH(CH ₃)CH ₂ CH ₃	H
8	0	C(CH ₃) ₃	H
9	0	C ₅ H ₉	H
10	0	C ₆ H ₁₁ (cyclo)	H
11	0	C ₆ H ₅	H
12	0	CH ₂ C ₆ H ₅	H
13	0	H	CH ₃


LINKER NO.	n	R ²⁰⁰	R ²⁰¹
14	0	H	C ₂ H ₅
15	0	H	C ₃ H ₇
16	0	H	CH(CH ₃) ₂
17	0	H	C ₄ H ₉
18	0	H	CH(CH ₃)CH ₂ CH ₃
19	0	H	C(CH ₃) ₃
20	0	H	C ₅ H ₉
21	0	H	C ₆ H ₁₃
22	0	H	C ₆ H ₅
23	0	H	CH ₂ C ₆ H ₅
24	0	H	C ₆ H ₁₁ (cyclo)
25	0	C ₆ H ₁₃	H
26	0	CH ₃	CH ₃
27	0	C ₂ H ₅	C ₂ H ₅
28	0	C ₃ H ₇	C ₃ H ₇
29	0	CH(CH ₃) ₂	CH(CH ₃) ₂
30	0	C ₄ H ₉	C ₄ H ₉
31	0	CH(CH ₃)CH ₂ CH ₃	CH(CH ₃)CH ₂ CH ₃

LINKER NO.	n	R ²⁰⁰	R ²⁰¹
32	0	C(CH ₃) ₃	C(CH ₃) ₃
33	0	C ₅ H ₉	C ₅ H ₉
34	0	C ₆ H ₁₃	C ₆ H ₁₃
35	0	C ₆ H ₁₁ (cyclo)	C ₆ H ₁₁ (cyclo)
36	0	C ₆ H ₅	C ₆ H ₅
37	0	CH ₂ C ₆ H ₅	CH ₂ C ₆ H ₅
38	3	H	H
39	3	CH ₃	H
40	3	H	CH ₃
41	3	C ₆ H ₅	H
42	3	H	C ₆ H ₅
43	3	CH ₃	C ₆ H ₅
44	3	C ₆ H ₅	CH ₃
45	3	CH ₂ C ₆ H ₅	H
46	3	H	CH ₂ C ₆ H ₅
47	4	H	H
48	4	CH ₃	H

LINKER NO.	n	R ²⁰⁰	R ²⁰¹
49	4	H	CH ₃
50	4	C ₆ H ₅	H
51	4	H	C ₆ H ₅
52	4	CH ₃	C ₆ H ₅
53	4	C ₆ H ₅	CH ₃
54	4	CH ₂ C ₆ H ₅	H
55	4	H	CH ₂ C ₆ H ₅
56	5	H	H
57	5	CH ₃	H
58	5	H	CH ₃
59	5	C ₆ H ₅	H
60	5	H	C ₆ H ₅
61	5	CH ₃	C ₆ H ₅
62	5	C ₆ H ₅	CH ₃
63	5	CH ₂ C ₆ H ₅	H
64	5	H	CH ₂ C ₆ H ₅
65	6	H	H


LINKER NO.	n	R ²⁰⁰	R ²⁰¹
66	6	CH ₃	H
67	6	H	CH ₃
68	6	C ₆ H ₅	H
69	6	H	C ₆ H ₅
70	6	CH ₃	C ₆ H ₅
71	6	C ₆ H ₅	CH ₃
72	6	CH ₂ C ₆ H ₅	H
73	6	H	CH ₂ C ₆ H ₅

Another class of suitable diamino terminal linker groups is defined by Formula IV:



5

wherein each of Q and T is one or more groups independently selected from




10

wherein each of R²⁰² through R²⁰⁵ is independently selected from hydrido, hydroxy, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl, alkoxy, aralkoxy, aryloxy, alkoxyalkyl, haloalkyl, hydroxyalkyl, halo, cyano, amino, monoalkylamino, dialkylamino, carboxy, carboxyalkyl, alkanoyl, alkenyl, cycloalkenyl and alkynyl.

15


A preferred class of linker groups within Formula IV is defined by Formula V:

20




wherein each of R²⁰² and R²⁰³ is independently selected from hydrido, hydroxy, alkyl, phenalkyl, phenyl, alkoxy, benzyloxy, phenoxy, alkoxyalkyl, hydroxyalkyl, halo, amino, monoalkylamino, dialkylamino, carboxy, carboxyalkyl and alkanoyl; and wherein each of p and q is a number independently selected from one through six, inclusive; with the proviso that when each of R²⁰² and R²⁰³ is selected from halo, hydroxy, amino, monoalkylamino and dialkylamino, then the carbon to which R²⁰² or R²⁰³ is attached in Formula V is not adjacent to a nitrogen atom of Formula V.

A more preferred class of linker groups of Formula V consists of divalent radicals wherein each of R²⁰² and R²⁰³ is independently selected from hydrido, hydroxy, alkyl, alkoxy, amino, monoalkylamino, carboxy, carboxyalkyl and alkanoyl; and wherein each of p and q is a number independently selected from two through four, inclusive. Even more preferred are linker groups wherein each of R²⁰² and R²⁰³ is independently selected from hydrido, amino, monoalkylamino and carboxyl; and wherein each of p and q is independently selected from the numbers two and three. Most preferred is a linker group wherein each of R²⁰² and R²⁰³ is hydrido; and wherein each of p and q is two; such most preferred linker group is derived from a piperazinyl group and has the structure



In Table II there is shown a class of specific examples of cyclized, diamino-terminated linker groups within Formula V. These linker groups, identified as Linker Nos. 74-95, would be suitable to form a conjugate between a carbonyl moiety of an AII antagonist (designated as "I") and a carbonyl moiety of carbonyl terminated second residue such as the carbonyl moiety attached to the gamma carbon of a glutamyl residue (designated as "T").


TABLE II

I = inhibitor
T = acetyl-γ-glutamyl

LINKER NO.	R ²⁰⁶	R ²⁰⁷	R ²⁰⁸	R ²⁰⁹	R ²¹⁰	R ²¹¹	R ²¹²	R ²¹³
74	H	H	H	H	H	H	H	H
75	CH ₃	H	H	H	H	H	H	H
76	H	H	H	H	CH ₃	H	H	H
77	CH ₃	H	H	H	CH ₃	H	H	H
78	CH ₃	H	CH ₃	H	H	H	H	H
79	CH ₃	H	H	H	H	H	CH ₃	H
80	CH ₃	CH ₃	H	H	H	H	H	H
81	H	H	H	H	CH ₃	CH ₃	H	H
82	CH ₃	CH ₃	H	H	CH ₃	CH ₃	H	H
83	CH ₃	CH ₃	CH ₃	CH ₃	H	H	H	H
84	CH ₃	CH ₃	H	H	H	H	CH ₃	CH ₃
85	H	H	H	H	CH ₃	CH ₃	CH ₃	CH ₃


LINKER NO.	R206	R207	R208	R209	R210	R211	R212	R213
86	C ₆ H ₅	H	H	H	H	H	H	H
87	H	H	H	H	C ₆ H ₅	H	H	H
88	C ₆ H ₅	H	H	H	C ₆ H ₅	H	H	H
89	C ₆ H ₅	H	H	H	H	H	C ₆ H ₅	H
90	C ₆ H ₅	H	C ₆ H ₅	H	H	H	H	H
91	CH ₂ C ₆ H ₅	H	H	H	H	H	H	H
92	H	H	H	H	CH ₂ C ₆ H ₅	H	H	H
93	CH ₂ C ₆ H ₅	H	H	H	CH ₂ C ₆ H ₅	H	H	H
94	CH ₂ C ₆ H ₅	H	H	H	H	H	CH ₂ C ₆ H ₅	H
95	CH ₂ C ₆ H ₅	H	CH ₂ C ₆ H ₅	H	H	H	H	H

Another class of suitable diamino terminal linker groups is defined by Formula VI:



5 wherein each of R²¹⁴ through R²¹⁷ is independently selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, hydroxyalkyl, alkoxyalkyl, aralkyl, aryl, haloalkyl, amino, monoalkylamino, dialkylamino, cyanoamino, carboxyalkyl, alkylsulfinio, alkylsulfonyl, arylsulfinyl and arylsulfonyl;
10 and wherein p is a number selected from one through six inclusive.

A preferred class of linker groups within Formula VI consists of divalent radicals wherein each of R²¹⁴ and R²¹⁵ is hydrido; wherein each of R⁶² and R⁶³ is independently selected from hydrido, alkyl, phenalkyl, phenyl, alkoxyalkyl, hydroxyalkyl, haloalkyl and carboxyalkyl; and wherein p is two or three. A more preferred class of linker groups within Formula VI consists of divalent radicals wherein each of R²¹⁴ and R²¹⁵ is hydrido; wherein each of R²¹⁶ and R²¹⁷ is independently selected from hydrido and alkyl; and wherein p is two. A specific example of a more preferred linker within Formula VI is the divalent radical ethylenediamino. In Table III there is shown a class of specific examples of diamino-terminated linker groups within Formula VI. These linker groups, identified as Linker Nos. 96-134, would be suitable to form a conjugate between a carbonyl moiety of an AII antagonist (designated as "I") and a carbonyl moiety of carbonyl terminated second residue such as the carbonyl moiety attached to the gamma carbon of a glutamyl residue (designated as "T").

TABLE III

I = inhibitor
G = acetyl-γ-glutamyl

LINKER NO.	R ²¹⁸	R ²¹⁹	R ²²⁰	R ²²¹	R ²²²	R ²²³
96	H	H	H	H	H	H
97	H	H	H	H	H	CH ₃
98	H	H	H	CH ₃	H	H
99	H	H	H	CH ₃	H	CH ₃
100	CH ₃	H	H	H	H	H
101	H	CH ₃	H	H	H	H
102	H	H	H	H	CH ₃	CH ₃
103	H	H	CH ₃	CH ₃	H	H
104	CH ₃	CH ₃	H	H	H	H
105	H	H	H	H	H	C ₆ H ₅
106	H	H	H	C ₆ H ₅	H	H
107	H	H	H	C ₆ H ₅	H	C ₆ H ₅
108	C ₆ H ₅	H	H	H	H	H

LINKER NO.	R ²¹⁸	R ²¹⁹	R ²²⁰	R ²²¹	R ²²²	R ²²³
---------------	------------------	------------------	------------------	------------------	------------------	------------------

109	H	C ₆ H ₅	H	H	H	H
110	H	H	H	H	C ₆ H ₅	C ₆ H ₅
111	H	H	C ₆ H ₅	C ₆ H ₅	H	H
112	C ₆ H ₅	C ₆ H ₅	H	H	H	H
113	H	H	H	H	H	C ₂ H ₅
114	H	H	H	C ₂ H ₅	H	H
115	H	H	H	C ₂ H ₅	H	C ₂ H ₅
116	C ₂ H ₅	H	H	H	H	H
117	H	C ₂ H ₅	H	H	H	H
118	H	H	H	H	C ₂ H ₅	C ₂ H ₅
119	H	H	C ₂ H ₅	C ₂ H ₅	H	H
120	C ₂ H ₅	C ₂ H ₅	H	H	H	H
121	CH ₃	H	C ₆ H ₅	H	H	H
122	CH ₃	H	H	H	C ₆ H ₅	H
123	H	CH ₃	C ₆ H ₅	H	H	H
124	H	CH ₃	H	H	C ₆ H ₅	H
125	CH ₃	CH ₃	H	C ₆ H ₅	H	H

LINKER NO.	R ²¹⁸	R ²¹⁹	R ²²⁰	R ²²¹	R ²²²	R ²²³
---------------	------------------	------------------	------------------	------------------	------------------	------------------

126	CH ₃	CH ₃	H	H	H	C ₆ H ₅
127	H	H	H	H	H	CH ₂ C ₆ H ₅
128	H	H	H	CH ₂ C ₆ H ₅	H	H
129	CH ₂ C ₆ H ₅	H	H	H	H	H
130	H	CH ₂ C ₆ H ₅	H	H	H	H
131	CH ₃	H	CH ₂ C ₆ H ₅	H	H	H
132	CH ₃	H	H	H	CH ₂ C ₆ H ₅	H
133	H	CH ₃	CH ₂ C ₆ H ₅	H	H	H
134	H	CH ₃	H	H	CH ₂ C ₆ H ₅	H

The term "hydrido" denotes a single hydrogen atom (H). This hydrido group may be attached, for example, to a carbon atom to form a $\text{CH} \lt$ group or attached to an oxygen atom to form a hydroxyl group; or as another example, two hydrido groups may be attached to a carbon atom to form a $-\text{CH}_2-$ group. Where the term "alkyl" is used, either alone or within other terms such as "haloalkyl" and "hydroxyalkyl", the term "alkyl" embraces linear or branched radicals having one to about twenty carbon atoms or, preferably, one to about twelve carbon atoms. More preferred alkyl radicals are "lower alkyl" radicals having one to about ten carbon atoms. Most preferred are lower alkyl radicals having one to about five carbon atoms. The term "cycloalkyl" embraces cyclic radicals having three to about ten ring carbon atoms, preferably three to about six carbon atoms, such as cyclopropyl and cyclobutyl. The term "haloalkyl" embraces radicals wherein any one or more of the alkyl carbon atoms is substituted with one or more halo groups, preferably selected from bromo, chloro and fluoro. Specifically embraced by the term "haloalkyl" are monohaloalkyl, dihaloalkyl and polyhaloalkyl groups. A monohaloalkyl group, for example, may have either a bromo, a chloro, or a fluoro atom within the group. Dihalalkyl and polyhaloalkyl groups may be substituted with two or more of the same halo groups, or may have a combination of different halo groups. A dihaloalkyl group, for example, may have two fluoro atoms, such as difluoromethyl and difluorobutyl groups, or two chloro atoms, such as a dichloromethyl group, or one fluoro atom and one chloro atom, such as a fluoro-chloromethyl group. Examples of a polyhaloalkyl are trifluoromethyl, 1,1-difluoroethyl, 2,2,2-trifluoroethyl, perfluoroethyl and 2,2,3,3-tetrafluoropropyl groups. The term "difluoroalkyl" embraces alkyl groups having two fluoro atoms substituted on any one or two of the alkyl group carbon atoms. Preferably, when the difluoroalkyl group is attached at the triazole ring R^1 and R^2 positions of Formula I, the two

fluoro atoms are substituted on the carbon atom which is attached directly to the triazole ring. Such preferred difluoroalkyl group may be characterized as an "alpha-carbon difluoro-substituted difluoroalkyl group". The terms

5 "alkylol" and "hydroxyalkyl" embrace linear or branched alkyl groups having one to about ten carbon atoms any one of which may be substituted with one or more hydroxyl groups. The term "alkenyl" embraces linear or branched radicals having two to about twenty carbon atoms,

10 preferably three to about ten carbon atoms, and containing at least one carbon-carbon double bond, which carbon-carbon double bond may have either cis or trans geometry within the alkenyl moiety. The term "alkynyl" embraces linear or branched radicals having two to about twenty carbon atoms,

15 preferably two to about ten carbon atoms, and containing at least one carbon-carbon triple bond. The term "cycloalkenyl" embraces cyclic radicals having three to about ten ring carbon atoms including one or more double bonds involving adjacent ring carbons. The terms "alkoxy"

20 and "alkoxyalkyl" embrace linear or branched oxy-containing radicals each having alkyl portions of one to about ten carbon atoms, such as methoxy group. The term "alkoxyalkyl" also embraces alkyl radicals having two or more alkoxy groups attached to the alkyl radical, that is,

25 to form monoalkoxyalkyl and dialkoxyalkyl groups. The "alkoxy" or "alkoxyalkyl" radicals may be further substituted with one or more halo atoms, such as fluoro, chloro or bromo, to provide haloalkoxy or haloalkoxyalkyl groups. The term "alkylthio" embraces radicals containing a

30 linear or branched alkyl group, of one to about ten carbon atoms attached to a divalent sulfur atom, such as a methylthio group. The term "aryl" embraces aromatic radicals such as phenyl, naphthyl and biphenyl. Preferred aryl groups are those consisting of one, two, or three

35 benzene rings. The term "aralkyl" embraces aryl-substituted alkyl radicals such as benzyl, diphenylmethyl, triphenylmethyl, phenylethyl, phenylbutyl and diphenylethyl. The terms "benzyl" and "phenylmethyl" are

interchangeable. The terms "aryloxy" and "arylthio" denote radical respectively, aryl groups having an oxygen or sulfur atom through which the radical is attached to a nucleus, examples of which are phenoxy and phenylthio. The

5 terms "sulfinyl" and "sulfonyl", whether used alone or linked to other terms, denotes respectively divalent radicals SO and SO₂. The term "aralkoxy", alone or within another term, embraces an aryl group attached to an alkoxy group to form, for example, benzyloxy. The term "acyl"

10 whether used alone, or within a term such as acyloxy, denotes a radical provided by the residue after removal of hydroxyl from an organic acid, examples of such radical being acetyl and benzoyl. The term "heteroaryl" embraces aromatic ring systems containing one or two hetero atoms

15 selected from oxygen, nitrogen and sulfur in a ring system having five or six ring members, examples of which are thienyl, furanyl, pyridinyl, thiazolyl, pyrimidyl and isoxazolyl. Such heteroaryl may be attached as a substituent through a carbon atom of the heteroaryl ring

20 system, or may be attached through a carbon atom of a moiety substituted on a heteroaryl ring-member carbon atom, for example, through the methylene substituent of imidazolemethyl moiety. Also, such heteroaryl may be attached through a ring nitrogen atom as long as

25 aromaticity of the heteroaryl moiety is preserved after attachment. The term "amido" denotes a radical consisting of a nitrogen atom attached to a carbonyl group, which radical may be further substituted in the manner described herein. The amido radical can be attached to the nucleus

30 of a compound of the invention through the carbonyl moiety or the nitrogen atom of the amido radical. The term "alkenylalkyl" denotes a radical having a double-bond unsaturation site between two carbons, and which radical may consist of only two carbons or may be further

35 substituted with alkyl groups which may optionally contain additional double-bond unsaturation. For any of the foregoing defined radicals, preferred radicals are those containing between one and about ten carbon atoms.

Specific examples of alkyl groups are methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, methylbutyl, dimethylbutyl, and neopentyl. Typical alkenyl and alkynyl groups may have one unsaturated bond, such as an allyl group, or may have a plurality or unsaturated bonds, with such plurality of bonds either adjacent, such as allene-type structures, or in conjugation, or separated by several saturated carbons.

10

Conjugates of the invention formed from compounds of Formula I have been found to inhibit the action of angiotensin II in mammals. Thus, conjugates of Formula I are therapeutically useful in methods for treating hypertension by administering to a hypertensive patient a therapeutically-effective amount of a conjugate containing a compound of Formula I, such that the conjugate is hydrolyzed by an enzyme found predominantly in the kidney so as to release an active angiotensin II antagonist species. The phrase "hypertensive patient" means, in this context, a mammalian subject suffering from the effects of hypertension or susceptible to a hypertensive condition if not treated to prevent or control such hypertension.

20

Included within the invention are conjugates of compounds of Formula I which are tautomeric forms of the described compounds, isomeric forms including diastereoisomers, and the pharmaceutically-acceptable salts thereof. The term "pharmaceutically-acceptable salts" embraces salts commonly used to form alkali metal salts and to form addition salts of free acids or free bases. The nature of the salt is not critical, provided that it is pharmaceutically-acceptable. Suitable pharmaceutically-acceptable acid addition salts of compounds of Formula I may be prepared from an inorganic acid or from an organic acid. Examples of such inorganic acids are hydrochloric, hydrobromic, hydroiodic, nitric, carbonic, sulfuric and phosphoric acid. Appropriate organic acids may be selected

25

30

35

from aliphatic, cycloaliphatic, aromatic, araliphatic, heterocyclic, carboxylic and sulfonic classes of organic acids, example of which are formic, acetic, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, ascorbic, glucuronic, maleic, fumaric, pyruvic, aspartic, glutamic, benzoic, anthranilic, p-hydroxybenzoic, salicylic, phenylacetic, mandelic, embonic (pamoic), methanesulfonic, ethanesulfonic, 2-hydroxyethanesulfonic, pantothenic, benzenesulfonic, toluenesulfonic, sulfanilic, mesylic, cyclohexylaminosulfonic, stearic, algenic, β -hydroxybutyric, malonic, galactaric and galacturonic acid. Suitable pharmaceutically-acceptable base addition salts of compounds of Formula I include metallic salts made from aluminium, calcium, lithium, magnesium, potassium, sodium and zinc or organic salts made from N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine) and procaine. All of these salts may be prepared by conventional means from the corresponding compound of Formula I by reacting, for example, the appropriate acid or base with the compound of Formula I. Also, such pharmaceutical salts may be formed with either a compound of Formula I which is contained in the conjugate, or such salts may be formed with the conjugate itself.

25

Conjugates of the invention can possess one or more asymmetric carbon atoms and are thus capable of existing in the form of optical isomers as well as in the form of racemic or non-racemic mixtures thereof. The optical isomers can be obtained by resolution of the racemic mixtures according to conventional processes, for example by formation of diastereoisomeric salts by treatment with an optically active acid or base. Examples of appropriate acids are tartaric, diacetyltartaric, dibenzoyltartaric, ditoluoyltartaric and camphorsulfonic acid and then separation of the mixture of diastereoisomers by crystallization followed by liberation of the optically active bases from these salts. A different process for

separation of optical isomers involves the use of a chiral chromatography column optimally chosen to maximize the separation of the enantiomers. Still another available method involves synthesis of covalent diastereoisomeric molecules by reacting conjugates with an optically pure acid in an activated form or an optically pure isocyanate. The synthesized diastereoisomers can be separated by conventional means such as chromatography, distillation, crystallization or sublimation, and then hydrolyzed to deliver the enantiomerically pure compound. The optically active conjugates can likewise be obtained by utilizing optically active starting materials. These isomers may be in the form of a free acid, a free base, an ester or a salt.

15

The first and second residues are provided by precursor compounds having suitable chemical moieties which react together to form a cleavable bond between the first and second residues. For example, the precursor compound of one of the residues will have a reactable carboxylic acid moiety and the precursor of the other residue will have a reactable amino moiety or a moiety convertible to a reactable amino moiety, so that a cleavable amide bond may be formed between the carboxylic acid moiety and the amino moiety.

25

Conjugates of the invention may be prepared using precursors of highly active angiotensin II antagonists of Formula I. Examples of lesser active, suitable precursors are acid chloride, esters and amides of angiotensin II antagonists of Formula I. For example, ester precursors of angiotensin II antagonists may be reacted with hydrazine to provide an amino terminal moiety which then can be reacted with a glutamic acid derivative to form a conjugate of the invention. Such precursors or intermediates themselves may be relatively strong, relatively weak, or inactive as AII antagonists. Also, conjugates of the invention may be prepared using

35

angiotensin II antagonists lacking a reactive terminal amino moiety. Such angiotensin II antagonists may be modified to contain a terminal acid moiety which then may be connected to a glutamyl residue through a diamino-terminated linker group, such as shown in Tables I-III.

A family of specific angiotensin II antagonist compounds of Formula I, from which a suitable first component of the conjugate may be selected, consists of biphenylmethyl 1H-substituted-1,2,4-triazole compounds listed below having a carboxylic acid terminal moiety or carboxylic acid moiety modified to be a hydrazide terminal moiety. All such compounds are characterized in having such carboxylic acid at one of the R⁵ through R⁹ positions of Formula I. Those compounds having a terminal carboxylic moiety may be reacted with one of the aforementioned linker groups, such as a hydrazine or a piperazine linker, to provide an amino terminal moiety which can then be reacted with the carboxylic acid moiety of a second component of the conjugate, such as a glutamic acid residue to form an enzyme-cleavable bond. Specific examples of these compounds are listed below:

4'-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid, hydrazide;
4'-[(5-butyl-3-chloro-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(3-butyl-5-chloro-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-ethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-propyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-isopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
methyl 4'-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylate;

- 4'-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-butyl-3-secbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 5 4'-[(5-butyl-3-isobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-butyl-3-tertbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-butyl-3-pentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 10 4'-[(5-butyl-3-isopentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-butyl-3-mercapto-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 15 4'-[(5-butyl-3-thiomethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-butyl-3-thioethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-butyl-3-thiopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 20 4'-[(5-butyl-3-hydroxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-butyl-3-methoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 25 4'-[(5-butyl-3-ethoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-butyl-3-propoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-butyl-3-butoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 30 4'-[(5-butyl-3-cyclohexyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-butyl-3-cyclohexylmethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 35 4'-[[5-butyl-3-(2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-butyl-3-cyclohexanoyl-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[[5-butyl-3-(1-oxo-2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-phenyl-1,2,4,-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
5 4'-[(5-butyl-3-phenylmethyl-1,2,4,-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-1,2,4,-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(3-butyl-1,2,4,-triazol-1-yl)methyl][1,1'-biphenyl]-2-
10 carboxylic acid;
4'-[[5-butyl-3-(2-phenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-benzoyl-1,2,4,-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
15 4'-[[5-butyl-3-(1-oxo-2-phenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-butyl-3-(1,1-dimethoxypropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-butyl-3-(1,1-dimethoxybutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
20 4'-[[5-butyl-3-(1,1-dimethoxypentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-butyl-3-(1-oxopropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
25 4'-[[5-butyl-3-(1-oxobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-butyl-3-(1-oxopentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-butyl-3-(1,1-difluoroethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
30 4'-[[5-butyl-3-(1,1-difluoropropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-butyl-3-(1,1-difluorobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
35 4'-[[3-butyl-5-(1,1-difluorobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-butyl-3-(1,1-difluoropentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[(5-propyl-3-ethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(3,5-dipropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 5 4'-[(3,5-isopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-propyl-3-isopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-propyl-3-butyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 10 4'-[(5-propyl-3-secbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-propyl-3-isobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 15 4'-[(5-propyl-3-tertbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-propyl-3-pentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-propyl-3-isopentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 20 4'-[(5-propyl-3-mercapto-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-propyl-3-thiomethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 25 4'-[(5-propyl-3-thioethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-propyl-3-thiopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-propyl-3-hydroxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 30 4'-[(5-propyl-3-methoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-propyl-3-ethoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 35 4'-[(5-propyl-3-propoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-propyl-3-cyclohexyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[(5-propyl-3-cyclohexylmethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
5-[4'-[[5-propyl-3-(2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
5 4'-[(5-propyl-3-cyclohexanoyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
5-[4'-[[5-propyl-3-(1-oxo-2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-propyl-3-phenyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-
10 biphenyl]-2-carboxylic acid;
4'-[(5-propyl-3-phenylmethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-propyl-3-(2-phenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
15 4'-[(5-propyl-3-benzoyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-propyl-3-(1-oxo-2-phenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-propyl-3-(1,1-dimethoxypropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
20 4'-[[5-propyl-3-(1,1-dimethoxybutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-propyl-3-(1,1-dimethoxypentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
25 4'-[[5-propyl-3-(1-oxopropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-propyl-3-(1-oxobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-propyl-3-(1-oxopentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
30 4'-[[5-propyl-3-(1,1-difluoroethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[3,5-bis(heptafluoropropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
35 4'-[[5-propyl-3-(1,1-difluoropropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-propyl-3-(1,1-difluorobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[5-propyl-3-(1,1-difluoropentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(3,5-diethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
5 4'-[(5-ethyl-3-propyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-ethyl-3-isopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-ethyl-3-butyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
10 4'-[(5-ethyl-3-secbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-ethyl-3-isobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
15 4'-[(5-ethyl-3-tertbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-ethyl-3-pentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-ethyl-3-isopentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
20 4'-[(5-ethyl-3-mercapto-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-ethyl-3-thiomethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
25 4'-[(5-ethyl-3-thioethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-ethyl-3-thiopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-ethyl-3-hydroxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
30 4'-[(5-ethyl-3-methoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-ethyl-3-ethoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
35 4'-[(5-ethyl-3-propoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-ethyl-3-cyclohexyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[(5-ethyl-3-cyclohexylmethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-ethyl-3-(2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
5 4'-[(5-ethyl-3-cyclohexanoyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-ethyl-3-(1-oxo-2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-ethyl-3-phenyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
10 biphenyl]-2-carboxylic acid;
4'-[(5-ethyl-3-phenylmethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-ethyl-3-(2-phenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
15 4'-[(5-ethyl-3-benzoyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-ethyl-3-(1-oxo-2-phenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-ethyl-3-(1,1-dimethoxypropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
20 4'-[[5-ethyl-3-(1,1-dimethoxybutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-ethyl-3-(1,1-dimethoxypentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
25 4'-[[5-ethyl-3-(1-oxopropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-ethyl-3-(1-oxobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-ethyl-3-(1-oxopentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
30 4'-[[5-ethyl-3-(1,1-difluoroethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-ethyl-3-(1,1-difluoropropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
35 4'-[[5-ethyl-3-(1,1-difluorobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-ethyl-3-(1,1-difluoropentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[(5-secbutyl-3-ethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-secbutyl-3-propyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 5 4'-[(5-secbutyl-3-isopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-secbutyl-3-butyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(3,5-disecbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 10 4'-[(5-secbutyl-3-isobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-secbutyl-3-tertbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 15 4'-[(5-secbutyl-3-pentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-secbutyl-3-isopentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-secbutyl-3-mercapto-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 20 4'-[(5-secbutyl-3-thiomethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-secbutyl-3-thioethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 25 4'-[(5-secbutyl-3-thiopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-secbutyl-3-hydroxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-secbutyl-3-methoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 30 4'-[(5-secbutyl-3-ethoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-secbutyl-3-propoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 35 4'-[(5-secbutyl-3-cyclohexyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-secbutyl-3-cyclohexylmethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[5-secbutyl-3-(2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[5-secbutyl-3-cyclohexanoyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
5 4'-[5-secbutyl-3-(1-oxo-2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[5-secbutyl-3-phenyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[5-secbutyl-3-phenylmethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
10 4'-[5-secbutyl-3-(2-phenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[5-secbutyl-3-benzoyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
15 4'-[5-secbutyl-3-(1-oxo-2-phenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[5-secbutyl-3-(1,1-dimethoxypropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[5-secbutyl-3-(1,1-dimethoxybutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
20 4'-[5-secbutyl-3-(1,1-dimethoxypentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[5-secbutyl-3-(1-oxopropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
25 4'-[5-secbutyl-3-(1-oxobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[5-secbutyl-3-(1-oxopentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[5-secbutyl-3-(1,1-difluoroethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
30 4'-[5-secbutyl-3-(1,1-difluoropropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[5-secbutyl-3-(1,1-difluorobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
35 4'-[5-secbutyl-3-(1,1-difluoropentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[5-isobutyl-3-ethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[(5-isobutyl-3-propyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-isobutyl-3-isopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
5 4'-[(5-isobutyl-3-butyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-isobutyl-3-secbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(3,5-diisobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-
10 biphenyl]-2-carboxylic acid;
4'-[(5-isobutyl-3-tertbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-isobutyl-3-pentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
15 4'-[(5-isobutyl-3-isopentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-isobutyl-3-mercapto-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-isobutyl-3-thiomethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
20 4'-[(5-isobutyl-3-thioethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-isobutyl-3-thiopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
25 4'-[(5-isobutyl-3-hydroxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-isobutyl-3-methoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-isobutyl-3-ethoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
30 4'-[(5-isobutyl-3-propoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-isobutyl-3-cyclohexyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
35 4'-[(5-isobutyl-3-cyclohexylmethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-isobutyl-3-(2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[(5-isobutyl-3-cyclohexanoyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-isobutyl-3-(1-oxo-2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
5 4'-[(5-isobutyl-3-phenyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-isobutyl-3-phenylmethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-isobutyl-3-(2-phenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
10 4'-[(5-isobutyl-3-benzoyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-isobutyl-3-(1-oxo-2-phenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
15 4'-[[5-isobutyl-3-(1,1-dimethoxypropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-isobutyl-3-(1,1-dimethoxybutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-isobutyl-3-(1,1-dimethoxypentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
20 4'-[[5-isobutyl-3-(1-oxopropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-isobutyl-3-(1-oxobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
25 4'-[[5-isobutyl-3-(1-oxopentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-isobutyl-3-(1,1-difluoroethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-isobutyl-3-(1,1-difluoropropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
30 4'-[[5-isobutyl-3-(1,1-difluorobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-isobutyl-3-(1,1-difluoropentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
35 4'-[(5-tertbutyl-3-ethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-tertbutyl-3-propyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[(5-tertbutyl-3-isopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-tertbutyl-3-butyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
5 4'-[(5-tertbutyl-3-secbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-tertbutyl-3-isobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(3,5-ditertbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-
10 biphenyl]-2-carboxylic acid;
4'-[(5-tertbutyl-3-pentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-tertbutyl-3-isopentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
15 4'-[(5-tertbutyl-3-mercapto-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-tertbutyl-3-thiomethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-tertbutyl-3-thioethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
20 4'-[(5-tertbutyl-3-thiopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-tertbutyl-3-hydroxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
25 4'-[(5-tertbutyl-3-methoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-tertbutyl-3-ethoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-tertbutyl-3-propoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
30 4'-[(5-tertbutyl-3-cyclohexyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-tertbutyl-3-cyclohexylmethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
35 4'-[(5-tertbutyl-3-(2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-tertbutyl-3-cyclohexanoyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[5-tertbutyl-3-(1-oxo-2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[5-tertbutyl-3-phenyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
5 4'-[5-tertbutyl-3-phenylmethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[5-tertbutyl-3-(2-phenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[5-tertbutyl-3-benzoyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
10 4'-[5-tertbutyl-3-(1-oxo-2-phenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[5-tertbutyl-3-(1,1-dimethoxypropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
15 4'-[5-tertbutyl-3-(1,1-dimethoxybutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[5-tertbutyl-3-(1,1-dimethoxypentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[5-tertbutyl-3-(1-oxopropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
20 4'-[5-tertbutyl-3-(1-oxobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[5-tertbutyl-3-(1-oxopentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
25 4'-[5-tertbutyl-3-(1,1-difluoroethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[5-tertbutyl-3-(1,1-difluoropropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[5-tertbutyl-3-(1,1-difluorobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
30 4'-[5-tertbutyl-3-(1,1-difluoropentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[5-pentyl-3-ethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
35 4'-[5-pentyl-3-propyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[5-pentyl-3-isopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[(5-pentyl-3-butyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-pentyl-3-secbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 5 4'-[(5-pentyl-3-isobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-pentyl-3-tertbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(3,5-dipentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 10 4'-[(5-pentyl-3-isopentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-pentyl-3-mercapto-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 15 4'-[(5-pentyl-3-thiomethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-pentyl-3-thioethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-pentyl-3-thiopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 20 4'-[(5-pentyl-3-hydroxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-pentyl-3-methoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 25 4'-[(5-pentyl-3-ethoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-pentyl-3-propoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-pentyl-3-cyclohexyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 30 4'-[(5-pentyl-3-cyclohexylmethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-pentyl-3-(2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 35 4'-[(5-pentyl-3-cyclohexanoyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-pentyl-3-(1-oxo-2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[(5-pentyl-3-phenyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-pentyl-3-phenylmethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 5 4'-[[5-pentyl-3-(2-phenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-pentyl-3-benzoyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-pentyl-3-(1-oxo-2-phenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 10 4'-[[5-pentyl-3-(1,1-dimethoxypropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-pentyl-3-(1,1-dimethoxybutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 15 4'-[[5-pentyl-3-(1,1-dimethoxypentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-pentyl-3-(1-oxopropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-pentyl-3-(1-oxobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 20 4'-[[5-pentyl-3-(1-oxopentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-pentyl-3-(1,1-difluoroethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 25 4'-[[5-pentyl-3-(1,1-difluoropropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-pentyl-3-(1,1-difluorobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-pentyl-3-(1,1-difluoropentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 30 4'-[(5-isopentyl-3-ethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-isopentyl-3-propyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 35 4'-[(5-isopentyl-3-isopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-isopentyl-3-butyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[(5-isopentyl-3-secbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-isopentyl-3-isobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
5 4'-[(5-isopentyl-3-tertbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-isopentyl-3-pentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(3,5-diisopentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-
10 biphenyl]-2-carboxylic acid;
4'-[(5-isopentyl-3-mercapto-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-isopentyl-3-thiomethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
15 4'-[(5-isopentyl-3-thioethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-isopentyl-3-thiopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-isopentyl-3-hydroxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
20 4'-[(5-isopentyl-3-methoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-isopentyl-3-ethoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
25 4'-[(5-isopentyl-3-propoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-isopentyl-3-cyclohexyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-isopentyl-3-cyclohexylmethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
30 4'-[(5-pentyl-3-(2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-isopentyl-3-cyclohexanoyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
35 4'-[(5-isopentyl-3-(1-oxo-2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-isopentyl-3-phenyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[(5-isopentyl-3-phenylmethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-isopentyl-3-(2-phenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
5 4'-[(5-isopentyl-3-benzoyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-isopentyl-3-(1-oxo-2-phenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-isopentyl-3-(1,1-dimethoxypropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
10 4'-[[5-isopentyl-3-(1,1-dimethoxybutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-isopentyl-3-(1,1-dimethoxypentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
15 4'-[[5-isopentyl-3-(1-oxopropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-isopentyl-3-(1-oxobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-isopentyl-3-(1-oxopentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
20 4'-[[5-isopentyl-3-(1,1-difluoroethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-isopentyl-3-(1,1-difluoropropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
25 4'-[[5-isopentyl-3-(1,1-difluorobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-isopentyl-3-(1,1-difluoropentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(1-butenyl)-3-ethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
30 4'-[[5-(1-butenyl)-3-propyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(1-butenyl)-3-isopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
35 4'-[[5-(1-butenyl)-3-butyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(1-butenyl)-3-secbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[[5-(1-butenyl)-3-isobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(1-butenyl)-3-tertbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 5 4'-[[5-(1-butenyl)-3-pentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(1-butenyl)-3-isopentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(1-butenyl)-3-mercapto-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 10 4'-[[5-(1-butenyl)-3-thiomethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(1-butenyl)-3-thioethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 15 4'-[[5-(1-butenyl)-3-thiopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(1-butenyl)-3-hydroxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(1-butenyl)-3-methoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 20 4'-[[5-(1-butenyl)-3-ethoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(1-butenyl)-3-propoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 25 4'-[[5-(1-butenyl)-3-cyclohexyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(1-butenyl)-3-cyclohexylmethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(1-butenyl)-3-(2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 30 4'-[[5-(1-butenyl)-3-cyclohexanoyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(1-butenyl)-3-(1-oxo-2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 35 4'-[[5-(1-butenyl)-3-phenyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(1-butenyl)-3-phenylmethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[[5-(1-butenyl)-3-(2-phenylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(1-butenyl)-3-benzoyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
5 4'-[[5-(1-butenyl)-3-(1-oxo-2-phenylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(1-butenyl)-3-(1,1-dimethoxypropyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(1-butenyl)-3-(1,1-dimethoxybutyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
10 4'-[[5-(1-butenyl)-3-(1,1-dimethoxypentyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(1-butenyl)-3-(1-oxopropyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
15 4'-[[5-(1-butenyl)-3-(1-oxobutyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(1-butenyl)-3-(1-oxopentyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(1-butenyl)-3-(1,1-difluoroethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
20 4'-[[5-(1-butenyl)-3-(1,1-difluoropropyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(1-butenyl)-3-(1,1-difluorobutyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
25 4'-[[5-(1-butenyl)-3-(1,1-difluoropentyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(2-butenyl)-3-ethyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(2-butenyl)-3-propyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
30 4'-[[5-(2-butenyl)-3-isopropyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(2-butenyl)-3-butyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
35 4'-[[5-(2-butenyl)-3-secbutyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(2-butenyl)-3-isobutyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[[5-(2-butenyl)-3-tertbutyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(2-butenyl)-3-pentyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
5 4'-[[5-(2-butenyl)-3-isopentyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(2-butenyl)-3-mercapto-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(2-butenyl)-3-thiomethyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
10 4'-[[5-(2-butenyl)-3-thioethyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(2-butenyl)-3-thiopropyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
15 4'-[[5-(2-butenyl)-3-hydroxy-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(2-butenyl)-3-methoxy-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(2-butenyl)-3-ethoxy-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
20 4'-[[5-(2-butenyl)-3-propoxy-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(2-butenyl)-3-cyclohexyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
25 4'-[[5-(2-butenyl)-3-cyclohexylmethyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(2-butenyl)-3-(2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(2-butenyl)-3-cyclohexanoyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
30 4'-[[5-(2-butenyl)-3-(1-oxo-2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(2-butenyl)-3-phenyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
35 4'-[[5-(2-butenyl)-3-phenylmethyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(2-butenyl)-3-(2-phenylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[[5-(2-butenyl)-3-(1,1-dimethoxypropyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(2-butenyl)-3-(1,1-dimethoxybutyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
5 4'-[[5-(2-butenyl)-3-(1,1-dimethoxypentyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(2-butenyl)-3-(1-oxopropyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(2-butenyl)-3-(1-oxobutyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
10 4'-[[5-(2-butenyl)-3-(1-oxopentyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(2-butenyl)-3-(1,1-difluoroethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
15 4'-[[5-(2-butenyl)-3-(1,1-difluoropropyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(2-butenyl)-3-(1,1-difluorobutyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(2-butenyl)-3-(1,1-difluoropentyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
20 4'-[[5-(3-butenyl)-3-ethyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(3-butenyl)-3-propyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
25 4'-[[5-(3-butenyl)-3-isopropyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(3-butenyl)-3-butyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(3-butenyl)-3-secbutyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
30 4'-[[5-(3-butenyl)-3-isobutyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(3-butenyl)-3-tertbutyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
35 4'-[[5-(3-butenyl)-3-pentyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(3-butenyl)-3-isopentyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[[5-(3-butenyl)-3-mercapto-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(3-butenyl)-3-thiomethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 5 4'-[[5-(3-butenyl)-3-thioethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(3-butenyl)-3-thiopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(3-butenyl)-3-hydroxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 10 4'-[[5-(3-butenyl)-3-methoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(3-butenyl)-3-ethoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 15 4'-[[5-(3-butenyl)-3-propoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(3-butenyl)-3-cyclohexyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(3-butenyl)-3-cyclohexylmethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 20 4'-[[5-(3-butenyl)-3-(2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(3-butenyl)-3-cyclohexanoyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 25 4'-[[5-(3-butenyl)-3-(1-oxo-2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(3-butenyl)-3-phenyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(3-butenyl)-3-phenylmethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 30 4'-[[5-(3-butenyl)-3-(2-phenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(3-butenyl)-3-benzoyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 35 4'-[[5-(3-butenyl)-3-(1-oxo-2-phenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(3-butenyl)-3-(1,1-dimethoxypropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[5-(3-butenyl)-3-(1,1-dimethoxybutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[5-(3-butenyl)-3-(1,1-dimethoxypentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 5 4'-[5-(3-butenyl)-3-(1-oxopropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[5-(3-butenyl)-3-(1-oxobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[5-(3-butenyl)-3-(1-oxopentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 10 4'-[5-(3-butenyl)-3-(1,1-difluoroethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[5-(3-butenyl)-3-(1,1-difluoropropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 15 4'-[5-(3-butenyl)-3-(1-difluorobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[5-(3-butenyl)-3-(1-difluoropentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[5-(1-butyryl)-3-ethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 20 4'-[5-(1-butyryl)-3-propyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[5-(1-butyryl)-3-butyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 25 4'-[5-(1-butyryl)-3-secbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[5-(1-butyryl)-3-isobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[5-(1-butyryl)-3-tertbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 30 4'-[5-(1-butyryl)-3-pentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[5-(1-butyryl)-3-isopentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 35 4'-[5-(1-butyryl)-3-mercapto-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[5-(1-butyryl)-3-thiomethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[[5-(1-butynyl)-3-thioethyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(1-butynyl)-3-thiopropyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
5 4'-[[5-(1-butynyl)-3-hydroxy-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(1-butynyl)-3-methoxy-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(1-butynyl)-3-ethoxy-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
10 4'-[[5-(1-butynyl)-3-propoxy-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(1-butynyl)-3-cyclohexyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
15 4'-[[5-(1-butynyl)-3-cyclohexylmethyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(1-butynyl)-3-(2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(1-butynyl)-3-cyclohexanoyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
20 4'-[[5-(1-butynyl)-3-(1-oxo-2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(1-butynyl)-3-phenyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
25 4'-[[5-(1-butynyl)-3-phenylmethyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(1-butynyl)-3-(2-phenylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(1-butynyl)-3-benzoyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
30 4'-[[5-(1-butynyl)-3-(1-oxo-2-phenylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(1-butynyl)-3-(1,1-dimethoxypropyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
35 4'-[[5-(1-butynyl)-3-(1,1-dimethoxybutyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(1-butynyl)-3-(1,1-dimethoxypentyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[[5-(1-butynyl)-3-(1-oxopropyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(1-butynyl)-3-(1-oxobutyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 5 4'-[[5-(1-butynyl)-3-(1-oxopentyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(1-butynyl)-3-(1,1-difluoroethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(1-butynyl)-3-(1,1-difluoropropyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 10 4'-[[5-(1-butynyl)-3-(1,1-difluorobutyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(1-butynyl)-3-(1,1-difluoropentyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 15 4'-[[5-(2-butynyl)-3-ethyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(2-butynyl)-3-propyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(2-butynyl)-3-butyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 20 4'-[[5-(2-butynyl)-3-secbutyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(2-butynyl)-3-isobutyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 25 4'-[[5-(2-butynyl)-3-tertbutyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(2-butynyl)-3-pentyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(2-butynyl)-3-isopentyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 30 4'-[[5-(2-butynyl)-3-mercapto-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(2-butynyl)-3-thiomethyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 35 4'-[[5-(2-butynyl)-3-thioethyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(2-butynyl)-3-thiopropyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[[5-(2-butynyl)-3-hydroxy-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(2-butynyl)-3-methoxy-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 5 4'-[[5-(2-butynyl)-3-ethoxy-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(2-butynyl)-3-propoxy-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(2-butynyl)-3-cyclohexyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 10 4'-[[5-(2-butynyl)-3-cyclohexylmethyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(2-butynyl)-3-(2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 15 4'-[[5-(2-butynyl)-3-cyclohexanoyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(2-butynyl)-3-(1-oxo-2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(2-butynyl)-3-phenyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 20 4'-[[5-(2-butynyl)-3-phenylmethyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(2-butynyl)-3-(2-phenylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 25 4'-[[5-(2-butynyl)-3-benzoyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(2-butynyl)-3-(1-oxo-2-phenylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(2-butynyl)-3-(1,1-dimethoxypropyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 30 4'-[[5-(2-butynyl)-3-(1,1-dimethoxybutyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(2-butynyl)-3-(1,1-dimethoxypentyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 35 4'-[[5-(2-butynyl)-3-(1-oxopropyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(2-butynyl)-3-(1-oxobutyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[[5-(2-butynyl)-3-(1-oxopentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(2-butynyl)-3-(1,1-difluoroethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 5 4'-[[5-(2-butynyl)-3-(1,1-difluoropropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(2-butynyl)-3-(1,1-difluorobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(2-butynyl)-3-(1,1-difluoropentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 10 4'-[[5-(3-butynyl)-3-ethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(3-butynyl)-3-propyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 15 4'-[[5-(3-butynyl)-3-isopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(3-butynyl)-3-butyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(3-butynyl)-3-secbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 20 4'-[[5-(3-butynyl)-3-isobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(3-butynyl)-3-tertbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 25 4'-[[5-(3-butynyl)-3-isopentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(3-butynyl)-3-pentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(3-butynyl)-3-mercapto-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 30 4'-[[5-(3-butynyl)-3-thiomethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(3-butynyl)-3-thioethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 35 4'-[[5-(3-butynyl)-3-thiopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(3-butynyl)-3-hydroxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[[5-(3-butynyl)-3-methoxy-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(3-butynyl)-3-ethoxy-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
5 4'-[[5-(3-butynyl)-3-propoxy-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(3-butynyl)-3-cyclohexyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(3-butynyl)-3-cyclohexylmethyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
10 4'-[[5-(3-butynyl)-3-(2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(3-butynyl)-3-cyclohexanoyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
15 4'-[[5-(3-butynyl)-3-(1-oxo-2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(3-butynyl)-3-phenyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(3-butynyl)-3-phenylmethyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
20 4'-[[5-(3-butynyl)-3-(2-phenylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(3-butynyl)-3-benzoyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
25 4'-[[5-(3-butynyl)-3-(1-oxo-2-phenylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(3-butynyl)-3-(1,1-dimethoxypropyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(3-butynyl)-3-(1,1-dimethoxybutyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
30 4'-[[5-(3-butynyl)-3-(1,1-dimethoxypentyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(3-butynyl)-3-(1-oxopropyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
35 4'-[[5-(3-butynyl)-3-(1-oxobutyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-(3-butynyl)-3-(1-oxopentyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[[5-(3-butynyl)-3-(1,1-difluoroethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(3-butynyl)-3-(1,1-difluoropropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 5 4'-[[5-(3-butynyl)-3-(1,1-difluorobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-(3-butynyl)-3-(1,1-difluoropentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(3,5-diethoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 10 4'-[(3,5-dipropoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(3,5-diisopropoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 15 4'-[(3,5-dibutoxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(3,5-dithiomethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(3,5-dithioethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 20 4'-[(3,5-dithiopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(3,5-dithioisopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 25 4'-[[3,5-di(1-butenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[3,5-di(2-butenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[3,5-di(3-butenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 30 4'-[[3,5-di(1-butynyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[3,5-di(2-butynyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid; and
- 35 4'-[[3,5-di(3-butynyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid.

Another family of specific angiotensin II antagonist compounds of Formula I, from which a suitable first component of the conjugate may be selected, consists of biphenylmethyl 1H-substituted-1,2,4-triazole compounds listed below having an amino terminal moiety attached at the R¹ or R² positions of Formula I.. Such amino terminal moiety may be reacted directly with the carboxylic acid moiety of a second component of the conjugate, such as a glutamic acid residue to form an enzyme-cleavable bond.

Specific examples of these compounds are listed below:

- 5-[4'-[(5-ethyl-3-amino-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[(5-ethyl-3-aminomethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[(5-ethyl-3-aminoethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[(5-ethyl-3-aminopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[(5-ethyl-3-aminobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-ethyl-5-(4-aminophenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-ethyl-5-(4-aminophenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-ethyl-5-(4-aminophenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-ethyl-5-(4-aminomethylphenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-ethyl-5-(4-aminomethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-ethyl-5-(4-aminoethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-ethyl-5-(4-aminocyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-ethyl-5-(4-aminocyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;

- 5-[4'-[[3-ethyl-5-(4-aminocyclohexylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-ethyl-5-(4-aminomethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5 5-[4'-[[3-ethyl-5-(4-aminomethylcyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-ethyl-5-(4-aminoethylcyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-propyl-3-amino-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
10 5-[4'-[(5-propyl-3-aminomethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-propyl-3-aminoethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
15 5-[4'-[(5-propyl-3-aminopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-propyl-3-aminobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-propyl-5-(4-aminophenyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
20 5-[4'-[[3-propyl-5-(4-aminophenylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-propyl-5-(4-aminophenylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
25 5-[4'-[[3-propyl-5-(4-aminomethylphenylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-propyl-5-(4-aminomethylphenyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-propyl-5-(4-aminoethylphenyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
30 5-[4'-[[3-propyl-5-(4-aminocyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-propyl-5-(4-aminocyclohexylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
35 5-[4'-[[3-propyl-5-(4-aminocyclohexylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;

- 5-[4'-[[3-propyl-5-(4-aminomethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-propyl-5-(4-aminomethylcyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-propyl-5-(4-aminoethylcyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[(5-isopropyl-3-amino-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 10 5-[4'-[(5-isopropyl-3-aminomethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[(5-isopropyl-3-aminoethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[(5-isopropyl-3-aminopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 15 5-[4'-[(5-isopropyl-3-aminobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-isopropyl-5-(4-aminophenyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 20 5-[4'-[[3-isopropyl-5-(4-aminophenylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-isopropyl-5-(4-aminophenylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-isopropyl-5-(4-aminomethylphenylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 25 5-[4'-[[3-isopropyl-5-(4-aminomethylphenyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-isopropyl-5-(4-aminoethylphenyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 30 5-[4'-[[3-isopropyl-5-(4-aminocyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-isopropyl-5-(4-aminocyclohexylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-isopropyl-5-(4-aminocyclohexylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 35 5-[4'-[[3-isopropyl-5-(4-aminomethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;

- 5-[4'-[[3-isopropyl-5-(4-aminomethylcyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isopropyl-5-(4-aminoethylcyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5 5-[4'-[(5-butyl-3-amino-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-butyl-3-aminomethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-butyl-3-aminoethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
10 5-[4'-[(5-butyl-3-aminopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-butyl-3-aminobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
15 5-[4'-[[3-butyl-5-(4-aminophenyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminophenylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminophenylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
20 5-[4'-[[3-butyl-5-(4-aminomethylphenylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminomethylphenyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
25 5-[4'-[[3-butyl-5-(4-aminoethylphenyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminocyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminocyclohexylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
30 5-[4'-[[3-butyl-5-(4-aminocyclohexylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminomethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
35 5-[4'-[[3-butyl-5-(4-aminomethylcyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminoethylcyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;

- 5-[4'-[(5-secbutyl-3-amino-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-secbutyl-3-aminomethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5 5-[4'-[(5-secbutyl-3-aminoethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-secbutyl-3-aminopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-secbutyl-3-aminobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
10 5-[4'-[[3-secbutyl-5-(4-aminophenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-secbutyl-5-(4-aminophenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-secbutyl-5-(4-aminophenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
15 5-[4'-[[3-secbutyl-5-(4-aminomethylphenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-secbutyl-5-(4-aminomethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-secbutyl-5-(4-aminocyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
20 5-[4'-[[3-secbutyl-5-(4-aminocyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-secbutyl-5-(4-aminocyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-secbutyl-5-(4-aminomethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
25 5-[4'-[[3-secbutyl-5-(4-aminomethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-secbutyl-5-(4-aminocyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
30 5-[4'-[[3-secbutyl-5-(4-aminocyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-secbutyl-5-(4-aminomethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
35 5-[4'-[(5-isobutyl-3-amino-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;

- 5-[4'-[(5-isobutyl-3-aminomethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-isobutyl-3-aminoethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5 5-[4'-[(5-isobutyl-3-aminopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-isobutyl-3-aminobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isobutyl-5-(4-aminophenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
10 5-[4'-[[3-isobutyl-5-(4-aminophenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isobutyl-5-(4-aminophenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
15 5-[4'-[[3-isobutyl-5-(4-aminomethylphenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isobutyl-5-(4-aminomethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isobutyl-5-(4-aminoethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
20 5-[4'-[[3-isobutyl-5-(4-aminocyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isobutyl-5-(4-aminocyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
25 5-[4'-[[3-isobutyl-5-(4-aminocyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isobutyl-5-(4-aminomethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
30 5-[4'-[[3-isobutyl-5-(4-aminomethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isobutyl-5-(4-aminoethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-tertbutyl-3-amino-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
35 5-[4'-[(5-tertbutyl-3-aminomethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;

- 5-[4'-[(5-tertbutyl-3-aminoethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-tertbutyl-3-aminopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5 5-[4'-[(5-tertbutyl-3-aminobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-tertbutyl-5-(4-aminophenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-tertbutyl-5-(4-aminophenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
10 5-[4'-[[3-tertbutyl-5-(4-aminophenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-tertbutyl-5-(4-aminomethylphenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
15 5-[4'-[[3-tertbutyl-5-(4-aminomethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-tertbutyl-5-(4-aminoethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-tertbutyl-5-(4-aminocyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
20 5-[4'-[[3-tertbutyl-5-(4-aminocyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-tertbutyl-5-(4-aminocyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
25 5-[4'-[[3-tertbutyl-5-(4-aminomethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-tertbutyl-5-(4-aminomethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
30 5-[4'-[[3-tertbutyl-5-(4-aminoethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-pentyl-3-amino-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-pentyl-3-aminomethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
35 5-[4'-[(5-pentyl-3-aminoethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;

- 5-[4'-[(5-pentyl-3-aminopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-pentyl-3-aminobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5 5-[4'-[[3-pentyl-5-(4-aminophenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-pentyl-5-(4-aminophenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-pentyl-5-(4-aminophenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
10 5-[4'-[[3-pentyl-5-(4-aminomethylphenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-pentyl-5-(4-aminomethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
15 5-[4'-[[3-pentyl-5-(4-aminoethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-pentyl-5-(4-aminocyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-pentyl-5-(4-aminocyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
20 5-[4'-[[3-pentyl-5-(4-aminocyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-pentyl-5-(4-aminomethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
25 5-[4'-[[3-pentyl-5-(4-aminomethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-pentyl-5-(4-aminoethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
30 5-[4'-[(5-isopentyl-3-amino-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-isopentyl-3-aminomethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-isopentyl-3-aminoethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
35 5-[4'-[(5-isopentyl-3-aminopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;

- 5-[4'-[(5-isopentyl-3-aminobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-isopentyl-5-(4-aminophenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5 5-[4'-[[3-isopentyl-5-(4-aminophenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-isopentyl-5-(4-aminophenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-isopentyl-5-(4-aminomethylphenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 10 5-[4'-[[3-isopentyl-5-(4-aminomethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-isopentyl-5-(4-aminomethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-isopentyl-5-(4-aminoethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 15 5-[4'-[[3-isopentyl-5-(4-aminocyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-isopentyl-5-(4-aminocyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-isopentyl-5-(4-aminocyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 20 5-[4'-[[3-isopentyl-5-(4-aminomethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-isopentyl-5-(4-aminomethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 25 5-[4'-[[3-isopentyl-5-(4-aminoethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[(5-hexyl-3-amino-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 30 5-[4'-[(5-hexyl-3-aminomethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[(5-hexyl-3-aminoethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[(5-hexyl-3-aminopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 35 5-[4'-[(5-hexyl-3-aminobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;

- 5-[4'-[[3-hexyl-5-(4-aminophenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-hexyl-5-(4-aminophenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-hexyl-5-(4-aminophenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-hexyl-5-(4-aminomethylphenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-hexyl-5-(4-aminomethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-hexyl-5-(4-aminoethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-hexyl-5-(4-aminocyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-hexyl-5-(4-aminocyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-hexyl-5-(4-aminocyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-hexyl-5-(4-aminomethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-hexyl-5-(4-aminomethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole; and
5-[4'-[[3-hexyl-5-(4-aminoethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole.

25

Another family of specific angiotensin II antagonist compounds of Formula I, from which a suitable first component of the conjugate may be selected, consists of biphenylmethyl 1H-substituted-1,2,4-triazole compounds listed below having a terminal carboxylic acid moiety attached at the R¹ or R² positions of Formula I.. Those compounds having a terminal carboxylic moiety may be reacted with one of the aforementioned linker groups, such as a hydrazine or a piperazine linker, to provide an amino terminal moiety which can then be reacted with the carboxylic acid moiety of a second component of the conjugate, such as a glutamic acid residue to form an

enzyme-cleavable bond. Specific examples of these compounds are listed below:

- 5- [4'-[(5-ethyl-3-carboxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5- [4'-[(5-ethyl-3-carboxymethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5- [4'-[(5-ethyl-3-carboxyethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 10 5- [4'-[(5-ethyl-3-carboxypropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5- [4'-[(5-ethyl-3-carboxybutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5- [4'-[[3-ethyl-5-(4-carboxyphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 15 5- [4'-[[3-ethyl-5-(4-carboxyphenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5- [4'-[[3-ethyl-5-(4-carboxyphenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 20 5- [4'-[[3-ethyl-5-(4-carboxymethylphenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5- [4'-[[3-ethyl-5-(4-carboxymethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5- [4'-[[3-ethyl-5-(4-carboxyethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 25 5- [4'-[[3-ethyl-5-(4-carboxycyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5- [4'-[[3-ethyl-5-(4-carboxycyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 30 5- [4'-[[3-ethyl-5-(4-carboxycyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5- [4'-[[3-ethyl-5-(4-carboxymethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 35 5- [4'-[[3-ethyl-5-(4-carboxymethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5- [4'-[[3-ethyl-5-(4-carboxyethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;

- 5-[4'-[(5-propyl-3-carboxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-propyl-3-carboxymethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5 5-[4'-[(5-propyl-3-carboxyethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-propyl-3-carboxypropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-propyl-3-carboxybutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
10 5-[4'-[[3-propyl-5-(4-carboxyphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-propyl-5-(4-carboxyphenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
15 5-[4'-[[3-propyl-5-(4-carboxyphenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-propyl-5-(4-carboxymethylphenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-propyl-5-(4-carboxymethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
20 5-[4'-[[3-propyl-5-(4-carboxyethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-propyl-5-(4-carboxycyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
25 5-[4'-[[3-propyl-5-(4-carboxycyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-propyl-5-(4-carboxycyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-propyl-5-(4-carboxymethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
30 1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-propyl-5-(4-carboxymethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-propyl-5-(4-carboxyethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
35 5-[4'-[(5-isopropyl-3-carboxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;

- 5-[4'-[(5-isopropyl-3-carboxymethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-isopropyl-3-carboxyethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5 5-[4'-[(5-isopropyl-3-carboxypropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-isopropyl-3-carboxybutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isopropyl-5-(4-carboxyphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
10 5-[4'-[[3-isopropyl-5-(4-carboxyphenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isopropyl-5-(4-carboxyphenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
15 5-[4'-[[3-isopropyl-5-(4-carboxymethylphenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isopropyl-5-(4-carboxymethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
20 5-[4'-[[3-isopropyl-5-(4-carboxyethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isopropyl-5-(4-carboxycyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isopropyl-5-(4-carboxycyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
25 5-[4'-[[3-isopropyl-5-(4-carboxycyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isopropyl-5-(4-carboxymethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
30 5-[4'-[[3-isopropyl-5-(4-carboxymethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isopropyl-5-(4-carboxyethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
35 5-[4'-[(5-butyl-3-carboxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-butyl-3-carboxymethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;

- 5-[4'-[(5-butyl-3-carboxyethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[(5-butyl-3-carboxypropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5 5-[4'-[(5-butyl-3-carboxybutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-butyl-5-(4-carboxyphenyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-butyl-5-(4-carboxyphenylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 10 5-[4'-[[3-butyl-5-(4-carboxyphenylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-butyl-5-(4-carboxymethylphenylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 15 5-[4'-[[3-butyl-5-(4-carboxymethylphenyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-butyl-5-(4-carboxyethylphenyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-butyl-5-(4-carboxycyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 20 5-[4'-[[3-butyl-5-(4-carboxycyclohexylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-butyl-5-(4-carboxycyclohexylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 25 5-[4'-[[3-butyl-5-(4-carboxymethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[[3-butyl-5-(4-carboxymethylcyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 30 5-[4'-[[3-butyl-5-(4-carboxyethylcyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[(5-secbutyl-3-carboxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 5-[4'-[(5-secbutyl-3-carboxymethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 35 5-[4'-[(5-secbutyl-3-carboxyethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;

- 5-[4'-[(5-secbutyl-3-carboxypropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-secbutyl-3-carboxybutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5 5-[4'-[[3-secbutyl-5-(4-carboxyphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-secbutyl-5-(4-carboxyphenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-secbutyl-5-(4-carboxyphenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
10 5-[4'-[[3-secbutyl-5-(4-carboxymethylphenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-secbutyl-5-(4-carboxymethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
15 5-[4'-[[3-secbutyl-5-(4-carboxyethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-secbutyl-5-(4-carboxycyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
20 5-[4'-[[3-secbutyl-5-(4-carboxycyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-secbutyl-5-(4-carboxycyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-secbutyl-5-(4-carboxymethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
25 5-[4'-[[3-secbutyl-5-(4-carboxymethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-secbutyl-5-(4-carboxyethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
30 5-[4'-[(5-isobutyl-3-carboxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-isobutyl-3-carboxymethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
35 5-[4'-[(5-isobutyl-3-carboxyethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-isobutyl-3-carboxypropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;

- 5-[4'-[(5-isobutyl-3-carboxybutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isobutyl-5-(4-carboxyphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5 5-[4'-[[3-isobutyl-5-(4-carboxyphenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isobutyl-5-(4-carboxyphenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isobutyl-5-(4-carboxymethylphenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
10 1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isobutyl-5-(4-carboxymethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isobutyl-5-(4-carboxyethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
15 triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isobutyl-5-(4-carboxycyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isobutyl-5-(4-carboxycyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
20 5-[4'-[[3-isobutyl-5-(4-carboxycyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isobutyl-5-(4-carboxymethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isobutyl-5-(4-carboxymethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
25 5-[4'-[[3-isobutyl-5-(4-carboxyethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-tertbutyl-3-carboxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
30 5-[4'-[(5-tertbutyl-3-carboxymethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-tertbutyl-3-carboxyethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
35 5-[4'-[(5-tertbutyl-3-carboxypropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-tertbutyl-3-carboxybutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;

- 5-[4'-[[3-tertbutyl-5-(4-carboxyphenyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-tertbutyl-5-(4-carboxyphenylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5 5-[4'-[[3-tertbutyl-5-(4-carboxyphenylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-tertbutyl-5-(4-carboxymethylphenylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
10 5-[4'-[[3-tertbutyl-5-(4-carboxymethylphenyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-tertbutyl-5-(4-carboxyethylphenyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-tertbutyl-5-(4-carboxycyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
15 5-[4'-[[3-tertbutyl-5-(4-carboxycyclohexylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-tertbutyl-5-(4-carboxycyclohexylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
20 5-[4'-[[3-tertbutyl-5-(4-carboxymethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-tertbutyl-5-(4-carboxymethylcyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
25 5-[4'-[[3-tertbutyl-5-(4-carboxyethylcyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-pentyl-3-carboxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-pentyl-3-carboxymethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
30 5-[4'-[(5-pentyl-3-carboxyethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-pentyl-3-carboxypropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
35 5-[4'-[(5-pentyl-3-carboxybutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-pentyl-5-(4-carboxyphenyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;

- 5-[4'-[[3-pentyl-5-(4-carboxyphenylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-pentyl-5-(4-carboxyphenylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5 5-[4'-[[3-pentyl-5-(4-carboxymethylphenylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-pentyl-5-(4-carboxymethylphenyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-pentyl-5-(4-carboxyethylphenyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
10 5-[4'-[[3-pentyl-5-(4-carboxycyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-pentyl-5-(4-carboxycyclohexylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-pentyl-5-(4-carboxycyclohexylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
15 5-[4'-[[3-pentyl-5-(4-carboxymethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-pentyl-5-(4-carboxymethylcyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
20 5-[4'-[[3-pentyl-5-(4-carboxyethylcyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[5-isopentyl-3-carboxy-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
25 5-[4'-[[5-isopentyl-3-carboxymethyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[5-isopentyl-3-carboxyethyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
30 5-[4'-[[5-isopentyl-3-carboxypropyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[5-isopentyl-3-carboxybutyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isopentyl-5-(4-carboxyphenyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
35 5-[4'-[[3-isopentyl-5-(4-carboxyphenylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;

- 5-[4'-[[3-isopentyl-5-(4-carboxyphenylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isopentyl-5-(4-carboxymethylphenylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5 [4'-[[3-isopentyl-5-(4-carboxymethylphenyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isopentyl-5-(4-carboxyethylphenyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
10 5-[4'-[[3-isopentyl-5-(4-carboxycyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isopentyl-5-(4-carboxycyclohexylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isopentyl-5-(4-carboxycyclohexylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
15 5-[4'-[[3-isopentyl-5-(4-carboxymethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-isopentyl-5-(4-carboxymethylcyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
20 5-[4'-[[3-isopentyl-5-(4-carboxyethylcyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-hexyl-3-carboxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
25 5-[4'-[(5-hexyl-3-carboxymethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-hexyl-3-carboxyethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-hexyl-3-carboxypropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
30 5-[4'-[(5-hexyl-3-carboxybutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-hexyl-5-(4-carboxyphenyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
35 5-[4'-[[3-hexyl-5-(4-carboxyphenylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-hexyl-5-(4-carboxyphenylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;

- 5-[4'-[[3-hexyl-5-(4-carboxymethylphenylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
 5-[4'-[[3-hexyl-5-(4-carboxymethylphenyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
 5 5-[4'-[[3-hexyl-5-(4-carboxyethylphenyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
 5-[4'-[[3-hexyl-5-(4-carboxycyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
 5-[4'-[[3-hexyl-5-(4-carboxycyclohexylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
 10 5-[4'-[[3-hexyl-5-(4-carboxycyclohexylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
 5-[4'-[[3-hexyl-5-(4-carboxymethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
 15 5-[4'-[[3-hexyl-5-(4-carboxymethylcyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole; and
 5-[4'-[[3-hexyl-5-(4-carboxyethylcyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole.
 20

A family of specific angiotensin II antagonist compounds of highest interest within Formula I from which a suitable first component of the conjugate may be selected, consists of amino-terminated biphenylmethyl 1H-substituted-
 25 1,2,4-triazole compounds as listed below. Such compounds would be suitable to form a conjugate with a carboxylic moiety of a second component of the conjugate, such as a glutamic acid residue, to form an enzyme-cleavable amide bond by direct reaction or by reaction through a diamino-
 30 containing linker of the type mentioned above. These Formula I angiotensin II antagonist compounds of highest interest are as follows:

- 35 methyl 4'-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylate;
 4'-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid, hydrazide;
- 4'-[(5-butyl-3-chloro-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 5 4'-[(3-butyl-5-chloro-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-butyl-3-propyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-butyl-3-isopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 10 4'-[(5-butyl-3-secbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-butyl-3-isobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 15 4'-[(5-butyl-3-tertbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-butyl-3-pentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-butyl-3-isopentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 20 4'-[(5-butyl-3-cyclohexyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-butyl-3-cyclohexylmethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 25 4'-[(5-butyl-3-(2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-butyl-3-cyclohexanoyl-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-butyl-3-(1-oxo-2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 30 4'-[(5-butyl-3-phenyl-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-butyl-3-phenylmethyl-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 35 4'-[(5-butyl-3-(2-phenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(5-butyl-3-benzoyl-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[[5-butyl-3-(1-oxo-2-phenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-butyl-3-(1,1-dimethoxypropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 5 4'-[[5-butyl-3-(1,1-dimethoxybutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-butyl-3-(1-oxopropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-butyl-3-(1-oxobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 10 4'-[[5-butyl-3-(1-oxopentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-butyl-3-(1,1-difluoroethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 15 4'-[[5-butyl-3-(1,1-difluoropropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[5-butyl-3-(1,1-difluorobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[[3-butyl-5-(1,1-difluorobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 20 4'-[[5-butyl-3-(1,1-difluoropentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(3,5-dipropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 25 4'-[(3,5-isopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(3,5-disecbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(3,5-diisobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 30 4'-[(3,5-ditertbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 4'-[(3,5-dipentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 35 4'-[(3,5-diisopentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 5-[4'-[(5-butyl-3-amino-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;

- 5-[4'-[(5-butyl-3-aminomethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-butyl-3-aminoethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5 5-[4'-[(5-butyl-3-aminopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-butyl-3-aminobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminophenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
10 5-[4'-[[3-butyl-5-(4-aminophenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminophenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
15 5-[4'-[[3-butyl-5-(4-aminomethylphenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminomethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminoethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
20 5-[4'-[[3-butyl-5-(4-aminocyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminocyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
25 5-[4'-[[3-butyl-5-(4-aminocyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminomethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminomethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
30 5-[4'-[[3-butyl-5-(4-aminoethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-butyl-3-carboxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
35 5-[4'-[(5-butyl-3-carboxymethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-butyl-3-carboxyethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;

- 5-[4'-[(5-butyl-3-carboxypropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-butyl-3-carboxybutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5 5-[4'-[[3-butyl-5-(4-carboxyphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-carboxyphenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-carboxyphenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
10 5-[4'-[[3-butyl-5-(4-carboxymethylphenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-carboxymethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
15 5-[4'-[[3-butyl-5-(4-carboxyethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-carboxycyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-carboxycyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
20 5-[4'-[[3-butyl-5-(4-carboxycyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-carboxymethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
25 tetrazole;
5-[4'-[[3-butyl-5-(4-carboxymethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole; and
5-[4'-[[3-butyl-5-(4-carboxyethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole.

General Synthetic Procedures


Conjugates of the invention are synthesized by reaction between precursors of the first and second residues. One of such precursors must contain a reactive acid moiety, and the other precursor must contain a reactive amino moiety, so that a conjugate is formed having a cleavable bond. Either precursor of the first and second residues may contain such reactive acid or amino moieties. Preferably, the precursors of the first residue are angiotensin II antagonists and will contain a reactive amino moiety or a moiety convertible to a reactive amino moiety. Inhibitor compounds lacking a reactive amino moiety may be chemically modified to provide such reactive amino moiety. Chemical modification of these inhibitor compounds lacking a reactive amino group may be accomplished by reacting an acid or an ester group on an AII antagonist compound with an amino compound having at least one reactive amino moiety and another reactive hetero atom selected from O, S and N. A suitable amino compound would be a diamino compound such as hydrazine or urea. Hydrazine, for example, may be reacted with a carboxylic acid or ester moiety of an AII antagonist compound to form a hydrazide derivative of such AII antagonist compound.

The AII antagonist compound 4'-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid may be used as a model compound to illustrate the chemical modification of a carboxylic acid-containing compound to make a reactive amino-containing precursor for synthesizing a conjugate of the invention. In the following General Synthetic Procedures, there are described firstly, in Schemes I-VI, methods for making suitable angiotensin II antagonists of Formula I for selection as the first component of the conjugate. Then, in Schemes VII-XII, there are described general methods for making a conjugate by reacting a first component AII antagonist of Formula I

with a cleavable second component represented by N-acetyl- γ -glutamic acid.

Conjugates of the invention may be prepared using precursors of highly active angiotensin II antagonists of Formula I. Examples of lesser active, suitable precursors are acid chloride, esters and amides of angiotensin II antagonists of Formula I. For example, ester precursors of more active angiotensin II antagonists, such as ester-type AII antagonists of Example Nos. 1, 34, 35, 67 and 68, as well as AII antagonists containing carboxylic acid terminal moieties, of Example Nos. 2, 6, 8, 18-20, 36, 52, 63, 64, 71-72 and 75-78, may be reacted with hydrazine to provide an intermediate bearing an amino terminal moiety which then can be reacted with a glutamic acid derivative to form a conjugate of the invention. Such precursors or intermediates themselves may be relatively strong, relatively weak, or inactive as AII antagonists. Also, conjugates of the invention may be prepared using angiotensin II antagonists lacking a reactive terminal acidic or amino moiety. Such angiotensin II antagonists, as shown in Example Nos. 3, 5, 7, 9-16, 21-33, 37-40, 42-51, 53-62, 65, 66, 69-70, 73 and 74, lack a terminal amino moiety. These AII antagonist compounds may be modified to contain a terminal amino moiety which then may be connected to a glutamyl residue through a diamino-terminated linker group, such as shown in Tables I-III.


Scheme I



$R = H, CH_3, \text{ or } C_2H_5$

- Synthetic Scheme I shows the preparation of 1H-1,2,4-triazoles 1 from ethyl iminoesters 2 and the corresponding hydrazide 3 via the general procedure outlined by H. Paul, G. Hilgetog and G. Jahnchen, Chem. Ber., 101, 2033 (1968). The free, iminoesters 2 can be prepared from the corresponding iminoester hydrochlorides 4, which in turn can be prepared from the corresponding nitrile 5; the procedures for the preparation of 2 and 4 from 5 are outlined by P.
- 10 Reynaud and R. D. Moreau, Bull. Soc. Chim. France, 2997 (1964). The hydrazides 3 can be either purchased or prepared from the corresponding alkyl esters 6 or lactones 6a and hydrazine.


Scheme II



Synthetic Scheme II shows the preparation of the alkylating agent 7 from the corresponding precursor 8.


When R⁵ equals CO₂CH₃, 8 was purchased from Chemo Dynamics Inc.

Scheme III




Synthetic Scheme III shows the preparation of the alkylating agent precursor 8 where R^5 equal $CN_4C(C_6H_5)_3$ from the corresponding methyl ester 8 ($R^5=CO_2CH_3$). In step 1, the methyl ester is converted to the corresponding acid ($R^5=CO_2H$) by the action of sodium hydroxide/hydrochloric acid. In step 2, the acid is converted to the corresponding acid chloride ($R^5=COCl$) by the action of oxalyl chloride. In step 3, the acid chloride is converted to the corresponding primary amide ($R^5=CONH_2$) by the action of ammonia. In step 4, the amide is converted to the corresponding nitrile 9 by the action of thionyl chloride at reflux. The nitrile 9 is reacted with trimethyltinazide in toluene at reflux to give the corresponding trimethyltin protected tetrazole 10; deprotection with acetic acid/water and reprotection with triphenylmethyl chloride/triethylamine gives the N-trityl tetrazole 8 ($R^5=CN_4C(C_6H_5)_3$).

Scheme IV




Synthetic Scheme IV shows the coupling reaction of the 1H-1,2,4-triazole 1 with the appropriate alkylating reagent 7. In the first step, 1 is treated with a base, such as sodium hydride, to generate the corresponding anion 11. Anion 11 is reacted with an alkylating agent 7 to give a mixture of regioisomers 12a and 12b. The isomer mixture may be converted to mixtures of the corresponding acids 13a and 13b or tetrazoles 14a and 14b by treatment with the appropriate reagent. Or, the isomers 12a and 12b may be separated by chromatographic methods, and each isomer may be reacted with the appropriate reagent to provide the acid- or tetrazole-substituted end product.

Scheme V




Synthetic Scheme V shows the regioselective synthesis of isomer 13a or 14a from Scheme IV. In the first step of the reaction, an alkylating agent 7 is
5 reacted with an appropriate hydrazide 3 to provide substituted hydrazide 15. An imidate 2 is reacted with hydrazide 15 to provide intermediate 16 which cyclizes upon heating to provide the corresponding product compound 13a or 14a.

Scheme VI




Synthetic Scheme VI shows the regioselective synthesis of isomer 13b or 14b from Scheme IV. In the first step of the reaction, imidate 2 is reacted with hydrazine to give amidazone 17. This intermediate is reacted with alkylating agent 7 to give intermediate 18 which is then cyclized in the presence of heat and an appropriate orthoester 19 to yield the corresponding product compound 13b or 14b.

Scheme VII



Synthetic Scheme VII shows the preparation of renal-selective angiotensin II antagonists by coupling γ -glutamic acid with an angiotensin II antagonist; the biphenyl R⁵ acid moiety of the AII antagonist is coupled to the γ -acid moiety of glutamic acid via an hydrazine linker. In step 1, the methyl ester of the AII antagonist 12b is converted to the hydrazide 20 by the action of hydrazine. In step 2, the hydrazide 20 is first reacted with the symmetrical anhydride of the protected γ -glutamic acid 21 and subsequently reacted with trifluoroacetic acid (TFA) to give the deprotected coupled material 22. In step 3, the free amino group of 22 is acetylated with acetic anhydride in the presence of base to give the renal-selective angiotensin II antagonists 23.


Scheme VIII



Synthetic Scheme VIII shows the preparation of angiotensin II antagonists 24 which have an amino moiety incorporated in R¹. In step 1, a protected amino acid 6 is first reacted with dicyclohexylcarbodiimide (DCC) and subsequently reacted with hydrazine to give the hydrazide 3. In step 2, the hydrazide 3 is reacted with the iminoester 2 and subsequently cyclized to the corresponding 1H-1,2,4-triazole 1. In step 3, the anion of 1 is reacted with the appropriate alkylating agent 7 to give a mixture of regioisomers 12a and 12b which can be separated by chromatography. In step 4, the desired isomer 12b is deprotected with TFA to give the free amino AII antagonist 24.


128

Scheme IX

 $\text{R} = \text{CH}_3, \text{C}_2\text{H}_5$


Synthetic Scheme IX shows the preparation of angiotensin II antagonists 25 and 26 which have a carbomethoxy moiety and an amino moiety, respectively, incorporated in R¹. In step 1, the dialkylacetal alkyl ester 6 is reacted with hydrazine to give the hydrazide 3. In step 2, the hydrazide 3 is reacted with the iminoester 2 and subsequently cyclized to the corresponding 1H-1,2,4-triazole 1. In step 3, the anion of 1 is reacted with the appropriate alkylating agent 7 to give a mixture of regioisomers which can be separated by chromatography prior to the generation of the free aldehyde 12b with aqueous acid. In step 4, the aldehyde 12b is oxidized to the corresponding acid with KMnO₄ and subsequently converted to the methyl ester 25 by SOCl₂/CH₃OH at -10° or reduced to the corresponding aminomethyl analog 26 (homolog of 24) by NH₄OAc/NaBH₃CN.

Scheme X



Synthetic Scheme X shows an alternate preparation of angiotensin II antagonists 25 which have a carbomethoxy moiety incorporated in R¹. In step 1, the lactone 6a is reacted with hydrazine to give the hydrazide 3. In step 2, the hydrazide 3 is reacted with the iminoester 2 and subsequently cyclized to the corresponding 1H-1,2,4-triazole 1. In step 3, the anion of 1 is reacted with the appropriate alkylating agent 7 to give a mixture of regioisomers which can be separated by chromatography to give 12b. In step 4, the primary alcohol 12b is oxidized to the corresponding acid with KMnO₄ and subsequently converted to the methyl ester 25 by SOCl₂/CH₃OH at -10°.

Scheme XI




Synthetic Scheme XI shows the preparation of renal-selective angiotensin II antagonists by coupling amino containing AII antagonists 24 with γ -glutamic acid. In step 1, the AII antagonist is reacted with the

5 symmetrical anhydride of the protected γ -glutamic acid 21 to give 27. In step 2, the protected material 27 is reacted with TFA to give the deprotected coupled material 28. In step 3, the free amino compound 28 is acetylated with acetic anhydride in the presence of base to give the


10 renal-selective angiotensin II antagonists 29.

Scheme XII



Synthetic Scheme XII shows the preparation of renal-selective angiotensin II antagonists by coupling γ -glutamic acid with an angiotensin II antagonist; the triazole R¹ acid moiety of the AII antagonist is coupled to the γ -acid moiety of glutamic acid via an hydrazine linker. In step 1, the methyl ester of the AII antagonist 25 is converted to the hydrazide 30 by the action of the hydrazine. In step 2, the hydrazide 30 is first reacted with the symmetrical anhydride of the protected γ -glutamic acid 21 and subsequently reacted with TFA to give the deprotected coupled material 31. In step 3, the free amino group of 31 is acetylated with acetic anhydride in the presence of base to give the renal-selective angiotensin II antagonists 32.

The following Examples 1-78 are detailed descriptions of the methods of preparation of specific angiotensin II antagonist compounds within Formula I. These detailed preparations fall within the scope of, and serve to exemplify, the above described General Synthetic Procedures which form part of the invention. These Examples #1-#78, as well as methods described in other preparatory examples which follow, are presented for illustrative purposes only and are not intended as a restriction on the scope of the invention. All parts are by weight and temperatures are given in centigrade degrees, unless otherwise indicated.

Example 1

5 methyl 4'-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylate

Step 1: Preparation of 4-bromomethyl-2'-methoxycarbonylbiphenyl.

10 A 47.46 g (210 mmol) sample of methyl 2-(p-tolyl)benzoate (Chemo Dynamics Inc.) was dissolved in 3L of carbon tetrachloride and treated with 37.33 g (209 mmol) of N-bromosuccinimide (NBS) and 1.17 g (7.13 mmol) of azobisisobutyronitrile (AIBN) at reflux under nitrogen for
15 24 hours. The reaction mixture was treated again with 1.0 g (6.1 mmol) of AIBN and stirred at reflux for an additional 24 hours. The reaction was filtered and the solvent removed in in vacuo. Purification by silica gel chromatography (Waters Prep-500A) using ethyl
20 acetate/hexane (5:95) as eluent provided 50.0 g (78%) of a colorless solid: mp 48-51°C; NMR (CDCl₃) δ 3.64 (s, 3H), 4.54 (s, 2H), 7.23-7.63 (m, 7H), 7.81-7.89 (m, 1H). NMR indicated that this material was only 91% pure; it contained 9% of the corresponding dibromocompound (δ 6.70);
25 however, no further attempts at purification were made and this mixture was used in all subsequent alkylation reactions.

Step 2: Preparation of 3,5-dibutyl-1H-1,2,4-triazole.


A solution of 64.5 g (0.50 mol) of ethyl
5 iminovalerate [P. Reynaud and R. C. Moreau, Bull. Soc.
Chim. France, 2997 (1964)] in 100 mL of methanol was added
slowly to 58.0 g (0.50 mol) of valeric acid hydrazide
(Lancaster Synthesis) in 400 mL of methanol at 0°C under a
nitrogen atmosphere. After the addition was complete, the
10 reaction was allowed to warm to ambient temperature and
then stir at reflux for 2 days. The solvent was removed in
vacuo; purification by silica gel chromatography (Waters
Prep-500A) using ethyl acetate/hexane (80:20) gave 78.9 g
(93%) of a colorless solid: mp 50.5-51.5°C; NMR (CDCl₃) δ
15 0.88 (t, J =7 Hz, 6H), 1.28-1.33 (m, 4H), 1.63-1.77 (m, 4H),
2.72 (t, J =7 Hz, 4H); MS (FAB) m/e (rel intensity) 183
(12%), 182 (100), 181 (3), 180 (6), 152 (8), 139 (4); HRMS.
Calcd for M+H: 182.1657. Found: 182.1661.

20 Step 3: Preparation of methyl 4'-[(3,5-dibutyl-1H-1,2,4-
triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylate.

Under a static nitrogen atmosphere, 2.01 g (11.0
mmol) of solid 3,5-dibutyl-1H-1,2,4-triazole was added in
25 small portions to 12 mmol of sodium hydride in 50 ml of
dimethylformamide (DMF); stirring was continued until
hydrogen evolution had ceased. The anion solution was
cooled to -10°C (ice/methanol) and treated with a solution
of 3.37 g (11.0 mmol) of 4-bromomethyl-2'-methoxycarbonyl-
30 biphenyl in 20 ml of dry DMF. The reaction was allowed to
warm to ambient temperature and stir overnight. Methanol
(10 ml) was added to destroy any unreacted sodium hydride
and the DMF was removed in vacuo. The residue was
dissolved in ethyl acetate, washed with water, and dried
35 (MgSO₄). Silica gel chromatography (Waters Prep-500A)
using 40% ethyl acetate/hexane gave 2.0 g (41%) of compound
as an oil: NMR (CDCl₃) δ 0.90 (t, J = 7 HZ, 3H), 0.94 (t,
 J = 7 Hz, 3H), 1.28-1.47 (m, 4H), 1.62-1.80 (m, 4H), 2.63-

2.75 (m, 4H), 3.63 (s, 3H), 5.27 (s, 2H), 7.13-7.18 (m, 2H), 7.25-7.35 (m, 3H), 7.37-7.44 (m, 1H), 7.48-7.55 (m, 1H), 7.80-7.85 (m, 1H) .

Example 2



4'-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid


5

A 2.0 g (4.9 mmol) sample of the methyl ester product compound from Example 1 was dissolved in 80 ml of ethanol and treated with 80 ml of 10% NaOH at ambient temperature for 3 days. The ethanol was removed in vacuo and the aqueous phase acidified to pH 1 with hydrochloric acid which caused the product to precipitate; filtration and drying in vacuo gave 1.65 g (86%) of colorless compound: mp 134-135°C; NMR (DMSO-d₆) δ 0.85 (t, J = 7 Hz, 3H), 0.90 (t, J = 7 Hz, 3H), 1.23-1.39 (m, 4H), 1.53-1.68 (m, 4H), 2.59 (t, J = 7 Hz, 2H), 2.78 (t, J = 7 Hz, 2H), 5.37 (s, 2H), 5.37 (s, 2H), 7.18-7.26 (m, 2H), 7.28-7.37 (m, 3H), 7.42-7.48 (m, 1H), 7.53-7.60 (m, 1H), 7.70-7.75 (m, 1H).

10

15

Example 3



5 5-[4'-[3-aminomethyl-5-butyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

Step 1: Preparation of N-Triphenylmethyl-5-[2-(4'-bromomethylbiphen-2-yl)]tetrazole.

10 A 542.5 g (2.4 mol) sample of methyl 2-(p-tolyl)benzoate (Chemo Dynamics Inc.) was dissolved in 5.5 L of ethanol and treated with 3 L (7.5 mol) of 2.5 N sodium hydroxide. The reaction was stirred overnight at ambient temperature and treated with an additional 480 ml (6.0 mol)

15 of sodium hydroxide; stirring was continued for an additional 24 h and the ethanol removed in vacuo. The remaining solution was cooled in ice and acidified to pH 1 with hydrochloric acid which caused the product to precipitate; filtration and drying in vacuo gave 510 g

20 (100%) of crude 2-(p-tolyl)benzoic acid: mp 145.0-147.5°C; NMR (CDCl₃) δ 2.40 (s, 3H), 7.17-7.28 (m, 4H), 7.35-7.45 (m, 2H), 7.51-7.59 (m, 1H), 7.90-7.97 (m, 1H). The crude acid was suspended in 1 L of toluene and slowly treated with 400 g (3.15 mol) of oxalyl chloride under nitrogen.

25 The reaction was allowed to stir at ambient temperature for 4.5 h and concentrated in vacuo to remove excess oxalyl chloride. The residue was redissolved in 2L of toluene and treated with 92.8 g (5.46 mol) of anhydrous ammonia. The

reaction was filtered and the filtrate concentrated in vacuo producing 424 g (84%) of crude 2-(p-tolyl)benzamide: mp 128-130°C; NMR (CDCl₃) δ 2.40 (s, 3H), 5.28 (br s, 1H), 5.77 (br s, 1H), 7.21-7.53 (m, 7H), 7.76-7.83 (m, 1H). The

5 crude amide was treated with 1420 ml (19.5 mol) of thionyl chloride at reflux for 3.5 h. The reaction was filtered and the thionyl chloride removed in vacuo. The residue was dissolved in 800 ml of toluene and reconcentrated in vacuo. On standing overnight, the residue crystallized. The

10 crystals were collected and washed with hexane to give 296 g (64%) of 2-(p-tolyl)benzonitrile: mp 50.5-52.0°C; NMR (CDCl₃) δ 2.42 (s, 3H), 7.22-7.34 (m, 2H), 7.37-7.52 (m, 3H), 7.58-7.66 (m, 1H), 7.72-7.78 (m, 1H). A 286 g (1.48 mol) sample of the crude nitrile was dissolved in 1630 mL

15 to toluene and treated with 377 g (1.8 mol) of trimethyltinazide at reflux for 24 h. The reaction was cooled; filtration gave 600 g of crude N-trimethylstannyl-5-[2-(4'-methylbiphen-2-yl)]tetrazole: mp 271-272°C (dec.); NMR (DMSO-d₆) δ 0.36 (br t, J=34 Hz, 9H), 2.24 (s, 3H),

20 6.89-7.06 (m, 4H), 7.35-7.55 (m, 4H). The crude N-trimethylstannyl tetrazole was suspended in 4270 mL of toluene and 287 mL of anhydrous tetrahydrofuran (THF) and treated with 63.4 g (173 mol) of anhydrous hydrogen chloride at ambient temperature under nitrogen with

25 stirring. The reaction was allowed to stand overnight and filtered; recrystallization from toluene gave 217 g (62%) of 5-[2-(4'-methylbiphen-2-yl)]tetrazole as a solid: mp 149-152°C; NMR (DMSO-d₆) δ 2.28 (s, 3H), 6.94-7.02 (m, 2H), 7.08-7.15 (m, 2H), 7.50-7.59 (m, 2H), 7.62-7.72 (m, 2H). A

30 200 g (0.85 mol) sample of the tetrazole was suspended in 3.3 L of dichloromethane and treated with 262 g (0.91 mol) of triphenylmethyl chloride and 141 mL (1.0 mol) of anhydrous triethylamine. The reaction was stirred at reflux for 3 h under nitrogen, washed with water, dried

35 (MgSO₄), and concentrated in vacuo. Recrystallization gave 338 g (83%) of N-triphenylmethyl-5-[2-(4'-methylbiphen-2-yl)]tetrazole as a colorless solid: mp 170-173°C; NMR (CDCl₃) δ 2.27 (s, 3H), 6.86-6.96 (m, 8H), 6.98-7.04 (m,

2H), 7.09-7.52 (m, 12H), 7.86-7.94 (m, 1H). The N-triphenylmethyl tetrazole was dissolved in 4260 mL of carbon tetrachloride and treated with 126.4 g (0.71 mol) of N-bromosuccinimide (NBS) and 11.9 g (49 mmol) of benzoyl peroxide at reflux for 3.5 h. The reaction was filtered and the solvent removed in vacuo. Recrystallization from toluene gave 277 g (59%) of N-triphenylmethyl-5-[2-(4'-bromophenylbiphen-2-yl)]tetrazole as a colorless solid: mp 140-141 °C; NMR (CDCl₃) δ 4.39 (s, 2H), 6.85-6.95 (m, 7H), 7.06-7.15 (m, 4H), 7.22-7.43 (m, 9H), 7.45-7.55 (m, 2H), 7.94-8.01 (m, 1H). NMR indicated that this material was only 85% pure; it contained 7% of corresponding dibromo-compound (δ 6.50) and 8% of starting material (δ 2.27); however, no further attempts at purification were made and this mixture was used in all subsequent alkylation reactions.


Step 2: Preparation of 5-butyl-3-tert-butoxycarbonylamino-methyl-1H-1,2,4-triazole.

20

Under nitrogen, a solution of 5.0 g (15.6 mmol) of anhydrous hydrazine in 100 mL of methanol is cooled to 0°C and treated with a solution of 25.0 g (13.2 mmol) of N-BOC glycine methyl ester (Chemical Dynamics Corporation) in 50 mL of methanol. The reaction is allowed to warm to ambient temperature and stirred overnight. Concentration in vacuo gives the crude hydrazide which is purified either by recrystallization or by silica gel chromatography (Waters Prep-500A). A 18.9 g (100 mmol) sample of hydrazide is dissolved in 100 mL of methanol and is treated with 12.9 g (100 mmol) of ethyl iminovalerate under nitrogen. The reaction is stirred at reflux overnight and concentrated in vacuo. Purification by silica gel chromatography (Waters Prep-500A) gives 5-butyl-3-tert-butoxycarbonylamino-methyl-1H-1,2,4-triazole.

Step 3: Preparation of 5-[4'-[(3-aminomethyl-5-butyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole.

5 Under a static nitrogen atmosphere, 3.81 g (15.0 mmol) of 3-butyl-5-tert-butoxycarbonylamino-1H-1,2,4-triazole is added slowly to 15 mmol of sodium hydride in 50 ml of dimethylformamide (DMF); stirring is continued until hydrogen evolution has ceased. The anion solution is
10 cooled to -10°C (ice/methanol) and treated with a solution of 8.45 g (15.0 mmol) of N-triphenylmethyl-5-[2-(4'-bromomethylbiphen-2-yl)]tetrazole in 20 ml of dry DMF. The reaction is allowed to warm to ambient temperature and stir overnight. Methanol (10 ml) is added to destroy any
15 unreacted sodium hydride and the DMF is removed in vacuo. The residue is dissolved in ethyl acetate, washed with water, and dried (MgSO₄). Purification by silica gel chromatography gives the desired isomer which is dissolved in 100 mL of TFA and is stored at ambient temperature
20 overnight. The reaction is cooled to 0°C, 50 mL of water is added, and is allowed to stir at ambient temperature overnight. The solvent is removed in vacuo and the residue is dissolved in acetonitrile/water. Purification by reverse phase chromatography (Waters Delta Prep.-3000)
25 provides pure 5-[4'-[(3-aminomethyl-5-butyl-1H-1,2,4-triazol-1-yl)[1,1'-biphenyl]-2-yl]-1H-tetrazole as the TFA salt.

Example 4


4'-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-
biphenyl]-2-carboxylic acid, hydrazide

5

A 7.10 g (17.5 mmol) sample of the methyl ester product compound of Example 1 was dissolved in 150 ml of methanol and treated with 22 ml (22.2 g 695 mmol) of anhydrous hydrazine under a static nitrogen atmosphere.

- 10 The reaction was stirred at reflux for 2 days and concentrated in vacuo to give 7.03 g (99%) of compound which was a colorless glass: NMR (CDCl₃) δ 0.88 (t, J = 7 Hz, 3H), 0.94 (t, J = 7 Hz, 3H), 1.28-1.47 (m, 4H), 1.62-1.78 (m, 4H), 2.62-2.73 (m, 4H), 3.5-4.1 (br s, 2H), 5.26
15 (s, 2H), 6.53 (s, 1H), 7.13-7.63 (m, 8H).

Example 5



5-[4'-[[3-(3-aminopropyl)-5-butyl-1H-1,2,4-triazol-1-yl]methyl]]-1,1'-biphenyl-2-yl]-1H-tetrazole.

5


Step 1: Preparation of 5-butyl-3-(3-tert-butoxycarbonylaminoethyl)-1H-1,2,4-triazole

Under nitrogen, 60.0 g (0.30 mol) of N-BOC-γ-aminobutyric acid (BACHEM) is dissolved in 1200 mL of methylene chloride and treated with 30.5 g (0.15 mol) of dicyclohexylcarbodiimide (DCC). The reaction is allowed to stir for 2 hours and filtered under nitrogen. The anhydride solution is then added to a solution of 1.58 g (98.5 mmol) of anhydrous hydrazine in 100 mL of methylene chloride at 0°C. The reaction is allowed to warm to ambient temperature and stirred overnight. Concentration in vacuo gives the crude hydrazide which is purified either by recrystallization or by silica gel chromatography (Waters Prep-500A). A 21.7 g (100 mmol) sample of hydrazide is dissolved in 100 mL of methanol and treated with 12.9 g (100 mmol) of ethyl iminovalerate under nitrogen. The reaction is stirred at reflux overnight and concentrated in vacuo. Purification by silica gel chromatography (Waters Prep-500A) gives 5-butyl-3-(3-tert-butoxycarbonylaminoethyl)-1H-1,2,4-triazole.

Step 2: Preparation of 5-[4'-[[3-(3-aminopropyl)-5-butyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

- 5 Under a static nitrogen atmosphere, 14.4 g (46.0 mmol) of 3-(3-N-BOC-aminopropyl)-5-butyl-1H-1,2,4-triazole is added in small portions to 50 mmol of sodium hydride in 250 ml of dimethylformamide (DMF); stirring is continued until hydrogen evolution has ceased. The anion
- 10 solution is cooled to -10°C (ice/methanol) and is treated with a solution of 25.5 g (46.0 mmol) of N-triphenylmethyl-5-[2-(4'-bromomethylbiphen-2-yl)]tetrazole (from Step 1 of Example 3) in 100 ml of dry DMF. The reaction is allowed to warm to ambient temperature and stir overnight.
- 15 Methanol (10 ml) is added to destroy any unreacted sodium hydride and the DMF is removed in vacuo; the residue is dissolved in ethyl acetate, washed with water, and dried (MgSO₄). Purification by silica gel chromatography gives the desired isomer which is dissolved in 200 mL of TFA and
- 20 is stirred at ambient temperature overnight. The reaction is cooled to 0°C, 100 mL of water is added, and is allowed to stir at ambient temperature overnight. The solvent is removed in vacuo and the residue is dissolved in acetonitrile/water. Purification by reverse phase
- 25 chromatography (Waters Delta Prep-3000) provided pure 5-[4'-[[3-(3-aminopropyl)-5-butyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole as the TFA salt.

Example 6



4'-[5-butyl-3-chloro-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid


5

Step 1: Preparation of 3-butyl-5-chloro-1H-1,2,4-triazole.

A 10.0 g (80 mmol) sample of 3-butyl-1H-1,2,4-triazole [H. Paul, G. Hilgetag, and G. Jahnchen, Chem. Ber., 101, 2033 (1968)] was dissolved in 320 mL of water containing 7.0 g (177 mmol) of sodium hydroxide. With stirring, the solution was cooled to 0°C and chlorine was introduced over 3 h. The reaction was purged with nitrogen overnight and the solution extracted with chloroform. The extracts were combined, dried (MgSO₄), and concentrated in vacuo to give 16.8 g of a colorless oil which was placed in 200 mL of water and treated twice with 8.0 g (80 mmol) of sodium metabisulfite. The pH of the reaction medium was adjusted to 6 with 1M sodium carbonate prior to extraction with chloroform; the extracts were dried (MgSO₄) and concentrated in vacuo to give 14.9 g of crude product. Purification by silica gel chromatograph (Waters Prep-500A) using chloroform/methanol (95:5) gave 9.53 g (75%) of a colorless solid: mp 104-105°C; NMR (CDCl₃) δ 0.94 (t, J=7 Hz, 3H), 1.33-1.47 (m, 2H), 1.68-1.83 (m, 2H), 2.80 (t, J=7 Hz, 2H); MS (FAB) m/e (rel intensity) 162 (28), 160 (100), 158 (10), 130 (5), 126 (10), 117 (5); HRMS. Calcd for M+H: 160.0642. Found: 160.0651.

Step 2: Preparation of 4'-[(5-butyl-3-chloro-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid.

5 Under a static nitrogen atmosphere, 5.0 g (31.3 mmol) of solid 5-butyl-3-chloro-1H-1,2,4-triazole was added in small portions to 32 mmol of sodium hydride in 50 ml of dimethylformamide (DMF); stirring was continued until hydrogen evolution had ceased. The anion solution was
10 cooled to -10°C (ice/methanol) and treated with a solution of 9.55 g (31.3 mmol) of 4-bromomethyl-2'-methoxycarbonylbiphenyl in 20 ml of dry DMF. The reaction was allowed to warm to ambient temperature and stir overnight. Methanol (10 ml) was added to destroy any
15 unreacted sodium hydride and the DMF was removed in vacuo. The residue was dissolved in ethyl acetate, washed with water, and dried (MgSO₄) to give 12.2 g of crude material was obtained which was a clear golden oil. A 4.81 g sample of this material was dissolved in 250 ml of methanol and
20 treated with 250 ml of 10% NaOH at ambient temperature for 2 days. A portion of the isomer mixture of acids was separated by reverse phase chromatography (Waters Delta Prep-3000) using isocratic 45% acetonitrile/water (0.05% TFA). The faster moving isomer (250 mg) was identified as
25 the 3-chloro isomer: NMR (DMSO-d₆) δ 0.86 (t, J = 7 Hz, 3H), 1.23-1.36 (m, J = 7 Hz, 2H), 1.54-1.65 (m, J = 7 Hz, 1H), 2.80 (t, J = 7Hz, 2H), 5.40 (s, 2H), 7.23-7.58 (m, 7H), 7.72-7.77 9m, 1H).

Example 7

5-[4'-[5-butyl-3-carboxymethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

5


Step 1: Preparation of 5-butyl-3-(2,2-diethoxyethyl)-1H-1,2,4-triazole.

- Under nitrogen, a solution of 38.4 g (1.2 mol) of anhydrous hydrazine in 500 mL of methanol is cooled to 0°C and treated with a solution of 190.0 g (1.0 mol) of ethyl 3,3-diethoxypropionate (Aldrich) in 200 mL of methanol. The reaction is allowed to warm to ambient temperature and is stirred overnight at reflux.
- 15 Concentration in vacuo gives the crude hydrazide which is purified either by recrystallization or by silica gel chromatography (Waters Prep-500A). A 17.6 g (100 mmol) sample of hydrazide is dissolved in 100 mL of methanol and is treated with 12.9 g (100 mmol) of ethyl iminovalerate under nitrogen. The reaction is stirred at reflux
- 20 overnight and concentrated in vacuo. Purification by silica gel chromatography (Waters Prep-500A) gives 5-butyl-3-(2,2-di-ethoxyethyl)-1H-1,2,4-triazole.

Step 2: Preparation of 5-[4'-[(5-butyl-3-carboxymethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole.

- 5 Under a static nitrogen atmosphere, 11.1 g (46.0 mmol) of 5-butyl-3-(2,2-diethoxyethyl)-1H-1,2,4-triazole is added in small portions to 50 mmol of sodium hydride in 250 ml of dimethylformamide (DMF); stirring is continued until hydrogen evolution has ceased. The anion solution is
- 10 cooled to -10°C (ice/methanol) and is treated with a solution of 25.5 g (9.2 mmol) of N-triphenylmethyl-5-[2-(4'-bromomethylbiphen-2-yl)]tetrazole (from Step 1 of Example 3) in 100 ml of dry DMF. The reaction is allowed to warm to ambient temperature and stir overnight.
- 15 Methanol (10 ml) is added to destroy any unreacted sodium hydride and the DMF is removed in vacuo; the residue is dissolved in ethyl acetate, washed with water, and dried (MgSO₄). Purification by silica gel chromatography gives the desired isomer which is treated with 3N HCl/methanol
- 20 (1:1) at reflux for 4 hours. The methanol is removed in vacuo and the pH is adjusted to 9 with NaOH. The solution is extracted with ethyl acetate twice to remove triphenylmethanol. The pH is readjusted to 3 with 6N HCl and the solution is extracted 3 times with ethyl acetate,
- 25 the extracts are combined, washed with water, and dried (MgSO₄). Concentration in vacuo gives 5-[4'-[(5-butyl-3-formalmethyl-1H-1,2,4-triazole-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole. A 4.01 g (10.0 mmol) sample of the free aldehyde is dissolved in 500 mL of acetone/water (1:1)
- 30 and is treated with 1.65 g (10.4 mmol) of solid KMnO₄ over 6 hours at ambient temperature. The reaction is allowed to stir overnight and excess KMnO₄ is destroyed by the addition of 100 mL of methanol. The reaction is filtered and the acetone is removed in vacuo; the pH is adjusted to
- 35 3 with 6N HCl and the product is extracted with ethyl acetate. The extracts are combined, washed with water, and dried (MgSO₄). Concentration in vacuo gives the crude product which was dissolved in acetonitrile/water;

purification by reverse phase chromatography (Waters Delta Prep-3000) provides pure 5-[4'-[5-butyl-3-carboxymethyl-1H-1,2,4-triazol-1-yl)methyl][1,1-biphenyl]-2-yl]-1H-tetrazole.


Example 8

4'-[(3-butyl-5-chloro-1H-1,2,4-triazol-1-yl)methyl][1,1'-
biphenyl]-2-carboxylic acid

5

The slower moving isomer (120 mg) isolated in
Example 6 was identified as the 5-chloro isomer: NMR
(DMSO- d_6) δ 0.88 (t, J = 7 Hz, 3H), 1.25-1.35 (m, J = 7 Hz,
2H), 1.56-1.67 (m, J = 7 Hz, 2H), 2.49 (t, J = 7 Hz, 2H),
10 5.37 (s, 2H), 7.23-7.58 (m, 7H), 7.72-7.77 (m, 1H).

Example 9



5-[4'-[1,1'-biphenyl]-2-yl]-1H-tetrazole-1-ylmethyl-1,2,4-triazole-3-carboxylate

Step 1: Preparation of 5-butyl-3-(3-hydroxypropyl)-1H-1,2,4-triazole.


Under nitrogen, a solution 38.4 g (1.2 mol) of anhydrous hydrazine in 500 mL of methanol is cooled to 0°C and is treated with a solution of 86.0 g (1.0 mol) of γ -butyrolactone (Aldrich) in 100 mL of methanol. The reaction is allowed to warm to ambient temperature and stir overnight at reflux. Concentration in vacuo gives the crude hydrazide which is purified either by recrystallization or by silica gel chromatography (Waters Prep-500A). A 11.8 g (100 mmol) sample of hydrazide is dissolved in 100 mL of methanol and is treated with 12.9 g (100 mmol) of ethyl iminovalerate under nitrogen. The reaction is stirred at reflux overnight and concentrated in vacuo. Purification by silica gel chromatography (Waters Prep-500A) gives 5-butyl-3-(3-hydroxypropyl)-1H-1,2,4-triazole.

Step 2: Preparation of 5-[4'-[[5-butyl-3-(2-carboxyethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole.

5 Under a static nitrogen atmosphere, 8.4 g (46 mmol) of 5-butyl-3-(3-hydroxypropyl)-1H-1,2,4-triazole is added in small portions to 50 ml of sodium hydride in 250 ml of dimethylformamide (DMF); stirring is continued until hydrogen evolution has ceased. The anion solution is
10 cooled to -10°C (ice/methanol) and treated with a solution of 25.5 g (46 mmol) of N-triphenylmethyl-5-[2-(4'-bromomethylbiphen-2-yl)]tetrazole (from Step 1 of Example 3) in 100 ml of dry DMF. The reaction is allowed to warm to ambient temperature and stir overnight. Methanol (10
15 ml) is added to destroy any unreacted sodium hydride and the DMF is removed in vacuo; the residue is dissolved in ethyl acetate, washed with water, and dried (MgSO₄). Purification by silica gel chromatography gives a mixture of isomers which is treated with 3N HCl/methanol (1:1) at
20 reflux for 4 hours. The methanol is removed in vacuo and the pH is adjusted to 9 with NaOH. The solution is extracted with ethyl acetate twice to remove triphenylmethanol. The pH is readjusted to 3 with 6 N HCl and the solution is extracted 3 times with ethyl acetate;
25 the extracts are combined, washed with water, and dried (MgSO₄). Concentration in vacuo gives a mixture of 5-butyl-3-(3-hydroxypropyl)- and 3-butyl-3-(3-hydroxypropyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole isomers. A 4.17 g (10.0 mmol) sample of the
30 isomeric mixture of alcohols is dissolved in 500 mL of acetone/water (1:1) and is treated with 3.30 g (20.8 mmol) of solid KMnO₄ over 6 hours at ambient temperature. The reaction is allowed to stir overnight and excess KMnO₄ is destroyed by the addition of 100 mL of methanol. The
35 reaction is filtered and the acetone is removed in vacuo; the pH is adjusted to 3 with 6N HCl and the product is extracted with ethyl acetate. The extracts are combined, washed with water, and dried (MgSO₄). Concentration in

vacuo give sthe isomeric mixture of acids; purification by reverse phase chromatography (Waters Delta Prep-3000) provides pure 5-[4'-[[5-butyl-3-(2-carboxyethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole.

Example 10



5-[4'-[[5-butyl-3-(1,1-difluorobutyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid

5

Step 1: Preparation of 3-butyl-5-(1,1-difluorobutyl)-1H-1,2,4-triazole.

Under nitrogen, a stirred solution of 56.9 g (0.65 mol) of N-tert-butyl-N-methylamine (Fluka) and 66.1 g (0.65 mol) of triethylamine in 1 L of methylene chloride was cooled to 0°C and treated with neat difluoroacetic anhydride [E. Sawicki, *J. Org. Chem.*, 21, 376 (1956)] at such a rate as to maintain the reaction temperature below 10°C. The reaction was allowed to warm to ambient temperature and stir overnight. All volatiles were removed in vacuo (bath temperature < 35°C) and the residue redissolved in methylene chloride; the solution was washed with saturated sodium bicarbonate, dried (MgSO₄), and concentrated to give 94 g (89%) of a yellow liquid. Vacuum distillation gave 86 g (81%) of colorless N-tert-butyl-N-methyldifluoroacetamide: bp 87-88°C (22 mm); ¹H NMR (CDCl₃) δ 1.37 (s, 9H), 2.93 (t, J=3 Hz, 3H), 5.97 (t, J=57 Hz, 1H); ¹⁹F NMR (CDCl₃) δ -122.20 (d, J=57 Hz, 2F). A 17.0 g (103 mmol) sample of the amide was dissolved in 50 mL of dry THF and added slowly to a solution of 145 mmol of

lithium diisopropylamine (LDA) in 450 mL of dry THF at -78°C. The reaction was allowed to stir for 1 h at -78°C prior to the addition of 17 mL (175 mmol) of 1-iodopropane by syringe. Stirring at -78°C was continued for 1 hr and then the reaction was allowed to warm to ambient temperature overnight. Methanol (10 mL) was added and the reaction was concentrated in vacuo; the residue was dissolved in methylene chloride and washed with 1N hydrochloric acid, dried (MgSO₄) and reconcentrated to give 17.6 g (83%) of crude product. Purification by vacuum distillation gave 13.3 g (62%) of colorless N-tert-butyl-N-methyl-2,2-difluoro-valeramide: bp 125-130°C (84 mm); ¹H NMR (CDCl₃) δ 0.94 (t, J=7 Hz, 3H), 1.37 (s, 9H), 1.40-1.52 (m, 2H), 1.95-2.15 (m, 2H); ¹⁹F NMR (CDCl₃) δ -100.29 (t, J = 20Hz, 2F). The difluorovaleramide was dissolved in 30 mL of trifluoroacetic acid (TFA) and stirred at reflux overnight under nitrogen. The solvent was removed in vacuo and the residue dissolved in methylene chloride; the solution was washed with water, dried (MgSO₄) and concentrated to give 9.7 g (100%) of crude N-methyl-2,2-difluorovaleramide: ¹H NMR (CDCl₃) δ 0.92 (t, J = 7 Hz, 3H), 1.36-1.51 (m, 2H), 1.90-2.11 (m, 2H), 2.84 (d, J = 6 Hz, 3H), 6.60-6.85 (br s, 1H); ¹⁹F NMR (CDCl₃) δ -106.98 (t, J = 19 Hz, 2F). The crude N-methyl amide was dissolved in 40 mL of 6N hydrochloric acid and stirred at reflux for 24 hr. The reaction was cooled to ambient temperature and extracted with methylene chloride; the extracts were combined, dried (MgSO₄), and concentrated to give 8.0 g (56%) of crude 2,2-difluorovaleric acid: ¹H NMR (CDCl₃) δ 1.00 (t, J = 7 Hz, 3H), 1.46-1.63 (m, 2H), 1.97-2.17 (m, 2H); ¹⁹F NMR (CDCl₃) δ -107.16 (t, J = 18 Hz, 2F). A 4.83 g (35 mmol) sample of the crude acid was dissolved in 25 mL (35.2 g, 174 mmol) of phthaloyl chloride in a flask equipped with a reflux condenser and stirred under nitrogen in a 110°C oil bath for 6 hrs. The condenser was replaced with a distillation head and 3.22 g (60%) of colorless 2,2-difluorovaleryl chloride was collected: bp 96°; ¹H NMR (CDCl₃) δ 1.03 (t, J = 7 Hz, 3H), 1.48-1.65 (m, 2H), 2.03-

2.23 (m, 2H); ^{19}F NMR (CDCl_3) δ -102.41 (t, J = 18 Hz, 2F). The 2,2-difluorovaleryl chloride (20.6 mmol) was dissolved in 10 mL of methylene chloride and dropwise to a solution of 135 g (42 mmol) of anhydrous hydrazine in 20 mL of methylene chloride at 0°C. After the addition was complete, the reaction was stirred at ambient temperature for 1 h, washed with water, dried (MgSO_4), and concentrated to give 3.12 g (91%) of 2,2-difluorovaleric acid hydrazide: NMR (CDCl_3) δ 0.96 (t, J = 7 Hz, 3H), 1.40-1.56 (m, 2H), 1.96-2.17 (m, 2H), 3.93 (br s, 2H), 7.67 (br s, 1H). A 2.84 g (18.7 mmol) sample of the crude hydrazide was dissolved in 50 mL of methanol and treated with 2.41 g (18.7 mmol) of ethyl iminovalerate. Under nitrogen, the reaction was stirred at reflux for 3 days and concentrated in vacuo. Purification by silica gel chromatography (Waters Prep-500A) using ethyl acetate/hexane (60:40) gave 3.0 g (74%) of 3-butyl-5-(1,1-difluorobutyl)-1H-1,2,4-triazole as a colorless solid: mp 92-93°C; ^1H NMR (CDCl_3) δ 0.97 (t, J = Hz, 3H), 0.92 (t, J = 7 Hz, 3H), 1.30-1.45 (m, 2H), 1.47-1.62 (m, 2H), 1.66-1.81 (m, 2H), 2.18-2.36 (m, 2H), 2.83 (t, J = 7 Hz, 2H); ^{19}F NMR (CDCl_3) δ -97.27 (t, J = 18 Hz, 2F); MS (FAB) m/e (rel intensity) 218 (100), 198 (8), 188 (5), 178 (8), 170 (5); HRMS. Calcd for $\text{M}+\text{H}$: 218.1469. Found: 218.1461.

25

Step 2: Preparation of 5-[4'-[[5-butyl-3-(1,1-difluorobutyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid

30


Under a static nitrogen atmosphere, 2.0 g (9.2 mmol) of solid 3-butyl-5-(1,1-difluorobutyl)-1H-1,2,4-triazole is added in small portions to 10 mmol of sodium hydride in 50 ml of dimethylformamide (DMF); stirring is continued until hydrogen evolution has ceased. The anion solution is cooled to -10°C (ice/methanol) and is treated with a solution of 3.0 g (9.2 mmol) of 4-bromomethyl-2'-methoxycarbonylbiphenyl (from Step 1 of Example 1) in 20 ml of dry DMF. The reaction is allowed to warm to ambient

35

temperature and stir overnight. Methanol (10 ml) is added to destroy any unreacted sodium hydride and the DMF is removed in vacuo; the residue is dissolved in ethyl acetate, washed with water, and dried (MgSO_4). Silica

5 gel chromatography separates the two isomers; the more abundant isomer is dissolved in 80 ml of ethanol and is treated with 80 ml of 10% NaOH at ambient temperature for 3 days. The ethanol is removed in vacuo and the aqueous phase is acidified to pH 1 with hydrochloric acid which

10 causes the product to precipitate; filtration and drying in vacuo gives 4'-[(5-butyl-3-(1,1-difluorobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid.


Example 11

4'-[[3-butyl-5-(1,1-difluorobutyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-carboxylic acid

5

The less abundant isomer isolated in Example 10 is dissolved in 40 ml of ethanol and is treated with 40 ml of 10% NaOH at ambient temperature for 3 days. The ethanol is removed in vacuo and the aqueous phase is acidified to pH 1 with hydrochloric acid which causes the product to precipitate; filtration and drying in vacuo gives 4'-[(3-butyl-5-(1,1-difluorobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid.

Example 12



5-[4'-[(3-aminomethyl-5-propyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

5

Step 1: Preparation of 5-propyl-3-tert-butoxycarbonylamino-methyl-1H-1,2,4-triazole.

Under nitrogen, a solution of 18.9 g (100 mmol) of N-Boc glycine hydrazide (from step 2 of Example 3) in 100 mL of methanol is treated with 11.5 g (100 mmol) of ethyl iminobutyrate [P. Reynaud and R. C. Moreau, Bull. Soc. Chim. France, 2997 (1964)]. The reaction is stirred at reflux overnight and concentrated in vacuo. Purification by silica gel chromatography (Waters Prep-500A) gives 5-propyl-3-tert-butoxycarbonylamino-methyl-1H-1,2,4-triazole.


Step 2: Preparation of 5-[4'-[(3-aminomethyl-5-propyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole.

Under a static nitrogen atmosphere, 3.81 g (15.0 mmol) of 3-propyl-5-tert-butoxy-carbonylamino-methyl-1H-1,2,4-triazole is added to 15 mmol of sodium hydride in 50 mL of DMF; stirring is continued until hydrogen evolution has ceased. The anion solution is cooled to -10°C

(ice/methanol) and treated with a solution of 8.45 g (15 mmol) of N-triphenylmethyl-5-[2-(4'-bromomethylbiphen-2-yl)]tetrazole in 20 mL of dry DMF. The reaction is allowed to warm to ambient temperature and stir overnight.

- 5 Methanol (10 mL) is added to destroy any unreacted sodium hydride and the DMF is removed in vacuo. The residue is dissolved in ethyl acetate, washed with water, and dried (MgSO₄). Purification by silica gel chromatography gives the desired isomer which is dissolved in 100 mL of TFA and
- 10 is stirred at ambient temperature overnight. The reaction is cooled to 0°C, 50 mL of water is added and is allowed to stir at ambient temperature overnight. The solvent is removed in vacuo and the residue is dissolved in acetonitrile/water. Purification by reverse phase
- 15 chromatography (Waters Delta Prep-3000) provides pure 5-[4'-[(3-aminomethyl-5-propyl-1H-1,2,4-triazole-1-yl)][1,1'-biphenyl]-2-yl]-1H-tetrazole as the TFA salt.

Example 13



5-[4'-[(3-aminomethyl-5-pentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

5

Step 1: Preparation of 5-pentyl-3-tert-butoxycarbonylamino-methyl-1H-1,2,4-triazole.

- Under nitrogen, a solution of 18.9 g (100 mmol) of N-Boc glycine hydrazide (from step 2 of Example 3) in 100 mL of methanol is treated with 14.3 g (100 mmol) of ethyl iminocaproate [P. Reynaud and R. C. Moreau, Bull. Soc. Chim. France, 2997 (1964)]. The reaction is stirred at reflux overnight and concentrated in vacuo.
- Purification by silica gel chromatography (Waters Prep-500A) gives 5-pentyl-3-tert-butoxycarbonylamino-methyl-1H-1,2,4-triazole.


Step 2: Preparation of 5-[4'-[(3-aminomethyl-5-pentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole.

- Under a static nitrogen atmosphere, 4.02 g (15.0 mmol) of 3-pentyl-5-tert-butoxy-carbonylamino-methyl-1H-1,2,4-triazole is added to 15 mmol of sodium hydride in 50 mL of DMF; stirring is continued until hydrogen evolution has ceased. The anion solution is cooled to -10°C

(ice/methanol) and treated with a solution of 8.45 g (15 mmol) of N-triphenylmethyl-5-[2-(4'-bromomethylbiphen-2-yl)]tetrazole in 20 mL of dry DMF. The reaction is allowed to warm to ambient temperature and stir overnight.

- 5 Methanol (10 mL) is added to destroy any unreacted sodium hydride and the DMF is removed in vacuo. The residue is dissolved in ethyl acetate, washed with water, and dried (MgSO₄). Purification by silica gel chromatography gives the desired isomer which is dissolved in 100 mL of TFA and
- 10 is stirred at ambient temperature overnight. The reaction is cooled to 0°C, 50 mL of water is added and is allowed to stir at ambient temperature overnight. The solvent is removed in vacuo and the residue is dissolved in acetonitrile/water. Purification by reverse phase
- 15 chromatography (Waters Delta Prep-3000) provides pure 5-[4'-[(3-aminomethyl-5-pentyl-1H-1,2,4-triazole-1-yl)][1,1'-biphenyl]-2-yl]-1H-tetrazole as the TFA salt.

Example 14



4'-[5-butyl-3-propyl-1H-1,2,4-triazol-1-yl)methyl]-[1,1'-biphenyl]-2-carboxylic acid


5

Step 1: Preparation of 5-butyl-3-propyl-1H-1,2,4-triazole.

A 3.95 g (38.7 mmol) sample of butyric acid hydrazide was dissolved in 30 mL of methanol and treated with 5.0 g (38.8 mmol) of ethyl iminovalerate under nitrogen. The reaction was stirred at reflux for 3 days and concentrated in vacuo. Purification by silica gel chromatography (Waters Prep-500A) using ethyl acetate/hexane (80:20) gave 5.51 g (85%) of 5-butyl-3-propyl-1H-1,2,4-triazole as a colorless solid: mp 48.5-50.0°C; NMR (CDCl₃) δ 0.92 (t, J = 7 Hz, 3H), 0.97 (t, J = 7 Hz, 3H), 1.31-1.46 (m, 2H), 1.66-1.84 (m, 4H), 2.72 (t, J = 7 Hz, 2H), 2.75 (t, J = 7 Hz, 2H); MS (FAB) m/e (rel intensity) 168 (100), 166 (4), 152 (3), 138 (3), 125 (3); HRMS. Calcd for M+H: 168.1501. Found: 168.1534.


Step 2: Preparation of 4'-[(5-butyl-3-propyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid.

Under a static nitrogen atmosphere, 2.0 g (12 mmol) of solid 5-butyl-3-propyl-1H-1,2,4-triazole is added in small portions to 12 mmol of sodium hydride in 50 ml of dimethylformamide (DMF); stirring is continued until hydrogen evolution has ceased. The anion solution is cooled to -10°C (ice/methanol) and treated with a solution of 3.8 g (12 mmol) of 4-bromomethyl-2'-methoxycarbonylbiphenyl (from Step 1 of Example 1) in 20 ml of dry DMF. The reaction is allowed to warm to ambient temperature and stir overnight. Methanol (10 ml) is added to destroy any unreacted sodium hydride and the DMF is removed in vacuo. The residue is dissolved in ethyl acetate, washed with water, and dried (MgSO₄). Silica gel chromatography produces a mixture of the two isomers which is dissolved in 80 ml of ethanol and is treated with 80 ml of 10% NaOH at ambient temperature for 3 days. The ethanol is removed in vacuo and the aqueous phase is acidified to pH 1 with hydrochloric acid which causes the mixture of product isomers to precipitate. Purification by reverse phase chromatography (Waters Delta Prep-3000) gives the 5-butyl-3-propyl isomer.

Example 15

4'-[(3-butyl-5-propyl-1H-1,2,4-triazol-1-yl)methyl]-[1,1'-
5 biphenyl]-2-carboxylic acid

The other isomer for Example 14 is isolated in an identical manner and provides the 3-butyl-5-propyl isomer.

Example 16

4'-[(3-butyl-5-isopentyl-1H-1,2,4-triazol-1-yl)methyl]-
[1,1'-biphenyl]-2-carboxylic acid

5

Step 1: Preparation of 3-butyl-5-isopentyl-1H-1,2,4-
triazole.

10

Under nitrogen, a solution of 53.8 g (1.68 mol) of anhydrous hydrazine in 300 mL of methanol was cooled to 0°C and treated with 186 g (1.4 mol) of methyl 4-methylvalerate. The reaction was allowed to warm to ambient temperature and stir overnight prior to stirring at reflux for 4 h. The reaction was concentrated in vacuo giving 166 g (93%) of 4-methylvaleric acid hydrazide as a colorless solid: 49-51°C; NMR (CDCl₃) δ 0.89 (d, J = 7 Hz, 6H), 1.48-1.64 (m, 3H), 2.15 (t, J = 7 Hz, 2H), 3.91 (br s, 2H), 6.94 (br s, 1H). A 7.86 g (60 mmol) sample of the

15

hydrazide was dissolved in 50 mL of methanol and treated with 7.8 g (60 mmol) of ethyl iminovalerate under nitrogen. The reaction was stirred at reflux overnight and concentrated in vacuo. Purification by silica gel chromatography (Waters Prep-500A) using ethyl acetate/hexane (40:60) gave 9.6 g (82%) of 3-butyl-5-isopentyl-1H-1,2,4-triazole as a colorless solid which melts close to ambient temperature: NMR (CDCl₃) δ 0.81-0.90 (m, 9H), 1.25-1.40 (m, 2H), 1.47-1.74 (m, 5H), 2.70 (t, J = 7 Hz, 4H); MS (FAB) m/e (rel intensity) 196 (100); HRMS. Calcd for M+H: 196.1814. Found: 196.1832.

Step 2: Preparation of 4'-[(3-butyl-5-isopentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid.

15


Under a static nitrogen atmosphere, 2.0 g (10.3 mmol) of solid 3-butyl-5-isopentyl-1H-1,2,4-triazole is added in small portions to 11 mmol of sodium hydride in 50 ml of dimethylformamide (DMF); stirring is continued until hydrogen evolution has ceased. The anion solution is cooled to -10°C (ice/methanol) and is treated with a solution of 3.27 g (10.2 mmol) of 4-bromomethyl-2'-methoxycarbonylbiphenyl (from Step 1 of Example 1) in 20 ml of dry DMF. The reaction is allowed to warm to ambient

20

temperature and stir overnight. Methanol (10 ml) is added to destroy any unreacted sodium hydride and the DMF is removed in vacuo. The residue is dissolved in ethyl acetate, washed with water, and dried (MgSO_4). The crude

5 material is dissolved in 80 ml of ethanol and is treated with 80 ml of 10% NaOH at ambient temperature for 3 days. The ethanol is removed in vacuo and the aqueous phase is acidified to pH 1 with hydrochloric acid which causes the mixture of product isomers to precipitate purification by

10 reverse phase chromatography (Waters Delta Prep-3000) give sthe 3-butyl-5-isopentyl isomer.


Example 17

4'-[(5-butyl-3-isopentyl-1H-1,2,4-triazol-1-yl)methyl]-
[1,1'-biphenyl]-2-carboxylic acid

5

The other isomer from Example 16 is isolated in an identical manner and provides the 5-butyl-3-isopentyl isomer.


Example 18



5 4'-[(5-butyl-1H-1,2,4-triazol-1-yl)methyl]-[1,1'-biphenyl]-
2-carboxylic acid

Following General Procedure A, 5.0 g (39.9 mmol)
 10 of 5-butyl-1H-1,2,4-triazole was coupled with 12.2 g (39.9
 mmol) of the alkylating reagent prepared in step 1 of
 Example 1 to give 3.1 g (22%) of a faster moving isomer:
 NMR (CDCl₃) δ 0.90 (t, J = 8 Hz, 3H), 1.29-1.45 (m, 2H),
 1.63-1.76 (m, 2H), 2.71 (t, J = 8 Hz, 2H), 3.62 (s, 3H),
 15 5.34 (s, 2H), 7.14-7.20 (m, 2H), 7.24-7.33 (m, 3H), 7.40
 (dt, J = 8 and 2 Hz, 1H), 7.50 (dt, J = 8 and 2 Hz, 1H),
 7.82 (dd, J = 8 and 2 Hz, 1H), 7.85 (s, 1H) and 3.7 g
 (26%) of a slower moving isomer: NMR (CDCl₃) δ 0.92 (t, J
 = 8 Hz, 3H), 1.31-1.46 (m, 2H), 1.66-1.79 (m, 2H), 2.73 (t,
 20 J = 8 Hz, 2H), 3.62 (s, 3H), 5.28 (s, 2H), 7.21-7.34 (m,
 5H), 7.39 (dt, J = 8 and 2 Hz, 1H), 7.50 (dt, J = 8 and 2
 Hz, 1H), 7.83 (dd, J = 8 and 2 Hz, 1H), 7.94 (s, 1H). The
 faster moving isomer was hydrolyzed to give 2.75 g (92%) of
 25 4'-[(5-butyl-1H-1,2,4-triazol-1-yl)methyl]-[1,1'-biphenyl]-
 2-carboxylic acid as a colorless solid: NMR (DMSO-d₆) δ
 0.85 (t, J = 8 Hz, 3H), 1.23-1.38 (m, 2H), 1.52-1.65 (m,
 2H), 2.76 (t, J = 7 Hz, 2H), 5.41 (s, 2H), 7.16-7.24 (m,
 2H), 7.27-7.37 (m, 3H), 7.44 (dt, J = 8 and 2 Hz, 1H), 7.56


(dt, $J = 8$ and 2 Hz, $1H$), 7.72 (dd, $J = 8$ and 2 Hz, $1H$),
 7.86 (s, $1H$); MS (FAB) m/e (rel intensity) 336 (100), 307
(7), 289 (7), 224 (8), 211 (100), 193 (9).

Example 19

5 4'-[(3-butyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-
2-carboxylic acid

The slower moving isomer from Example 18 was
hydrolyzed to give 2.3 g (67%) of 4'-[(3-butyl-1H-1,2,4-
10 triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid as a
colorless solid: NMR (DMSO-d₆) δ 0.87 (t, J = 8 Hz, 3H),
1.23-1.38 (m, 2H), 1.54-1.67 (m, 2H), 2.58 (t, J = 8 Hz,
2H), 5.36 (s, 2H), 7.23-7.38 (m, 5H), 7.44 (dt, J = 8 and
2 Hz, 1H), 7.56 (dt, J = 8 and 2 Hz, 1H), 7.72 (dd, J = 8
15 and 2 Hz, 1H), 8.51 (s, 1H); MS (FAB) m/e (rel intensity)
336 (85), 211 (100), 165 (24), 126 (52).

Example 20




5

4'-[(3,5-di-perfluoropropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid

Following General Procedure A, 11.0 g (27.2 mmol) of 3,5-bis(perfluoropropyl)-1H-1,2,4-triazole was coupled with 8.30 g (27.2 mmol) of the alkylating reagent prepared in step 1 of Example 1 to give 8.7 g (51%) of a colorless oil: NMR (CDCl₃) δ 3.62 (s, 3H), 5.63 (s, 2H), 7.25-7.37 (m, 5H), 7.44 (dt, $J = 8$ and 2 Hz, 1H), 7.54 (dt, $J = 8$ and 2 Hz, 1H), 7.87 (dd, $J = 8$ and 2 Hz, 1H). A 8.5 g (13.5 mmol) sample of this material was hydrolyzed to give 6.91 g (81%) of 4'-[(3,5-bis(1,1,2,2,3,3,3-heptafluoropropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid as a colorless solid: NMR (CDCl₃) δ 5.62 (s, 2H), 7.27-7.37 (m, 5H), 7.44 (dt, $J = 8$ and 2 Hz, 1H), 7.58 (dt, $J = 8$ and 2 Hz, 1H), 7.98 (dd, $J = 8$ and 2 Hz, 1H); MS (TSP) $M+NH_4$ (rel intensity) 633 (100), 211 (12); HRMS. Calc'd for $M+Li$: 622.0788. Found: 622.0759.

Example 21



5


5-[4'-[(3-phenyl-5-butyl-1H-1,2,4-triazol-1-yl)methyl]](1,1'-biphenyl)-2-yl]-1H-tetrazole

Following General Procedure A, 2.0 g (9.9 mmol) of 5-butyl-3-phenyl-1H-1,2,4 triazole was coupled with 5.5 g (9.9 mmol) of the alkylating reagent prepared in step 1 of Example 3 to give 5.2 g (77%) of a faster moving isomer: NMR (CDCl₃) δ 0.90 (t, J = 8 Hz), 1.30-1.44 (m, 2H), 1.63-1.77 (m, 2H), 2.67 (t, J = 8 Hz, 2H), 5.24 (s, 2H), 6.88-7.01 (m, 8H), 7.08-7.53 (m, 17H), 7.95 (dd, J = 8 and 2 Hz, 1H), 8.12 (dd, J = 8 and 2 Hz, 2H) and 420 mg (6.3%) of a slower moving isomer: NMR (CDCl₃) δ 0.96 (t, J = 8 Hz, 3H), 1.38-1.52 (m, 2H), 1.75-1.89 (m, 2H), 2.84 (t, J = 8 Hz, 2H), 5.27 (s, 2H), 6.88-6.97 (m, 8H), 7.13 (d, J = 8 Hz, 2H), 7.17-7.54 (m, 15H), 7.57 (d, J = 8 Hz, 2H), 7.93 (dd, J = 8 and 2 Hz, 1H). A 4.8 g (7.1 mmol) sample of the faster moving isomer was deprotected to give 1.64 g (53%) of 5-[4'-[(3-phenyl-5-butyl-1H-1,2,4-triazol-1-yl)methyl]](1,1'-biphenyl-2-yl)-1H-tetrazole as a colorless solid: mp 113 °C (dec); NMR (CDCl₃) δ 0.87 (t, J = 8 Hz, 3H), 1.24-1.39 (m, 2H), 1.55-1.69 (m, 2H), 2.60 (t, J = 8 Hz, 2H), 5.19 (s, 2H), 7.04 (d, J = 8 Hz, 2H), 7.11 (d, J = 8

Hz, 2H), 7.27-7.39 (m, 4H), 7.47-7.60 (m, 2H), 7.87 (dd, $J = 8$ and 2 Hz, 2H), 7.95 (dd, $J = 8$ and 2 Hz, 1H); MS (FAB) m/e (rel intensity) 436 (100), 393 (8), 207 (60), 192 (20), 178 (13), 165 (8); HRMS. Calc'd for M+H: 436.2249.

5 Found: 436.2240.

Example 22




5

5-[4'-[(3-butyl-5-phenyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

The slower moving isomer from Example 21 was
10 deprotected to give 251 mg (60%) of 5-[4'-[(3-butyl-5-phenyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole as a colorless solid: mp 205 °C (dec); NMR (CDCl₃) δ 0.88 (t, J = 8 Hz, 3H), 1.23-1.38 (m, 2H), 1.55-1.68 (m, 2H), 2.52 (t, J = 8 Hz, 2H), 5.32 (s, 2H), 6.99
15 (d, J = 8 Hz, 2H), 7.08 (d, J = 8 Hz, 2H), 7.31-7.61 (m, 8H), 7.90 (dd, J = 8 and 2 Hz, 1H); MS⁺ (FAB) m/e (rel intensity) 436 (100), 393 (9), 374 (7), 277 (7), 247 (15), 207 (45); HRMS. Calc'd for M+H: 436.2249. Found: 436.2201.

Example 23



5

5-[4'-[[[3-butyl-5-(dimethoxymethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole


10

Following General Procedure A, 2.0 g (10 mmol) of 5-butyl-3-(dimethoxymethyl)-1H-1,2,4-triazole was coupled with 5.6 g (10 mmol) of the alkylating reagent prepared in step 1 of Example 3 to give 1.65 g (24%) of a faster moving isomer: NMR (CDCl₃) δ 0.92 (t, J = 8 Hz, 3H), 1.31-1.45 (m, 2H), 1.66-1.78 (m, 2H), 2.69 (t, J = 8 Hz, 2H), 3.33 (s, 6H), 5.30 (s, 2H), 5.42 (s, 1H), 6.90-6.96 (m, 5H), 7.02-7.12 (m, 4H), 7.21-7.38 (m, 11H), 7.41-7.52 (m, 2H), 7.86-7.92 (m, 1H) and 3.65 g (54.0%) of a slower moving isomer: NMR (CDCl₃) δ 0.85 (t, J = 8 Hz, 3H), 1.22-1.36 (m, 2H), 1.57-1.70 (m, 2H), 2.59 (t, J = 8 Hz, 2H), 3.45 (s, 6H), 5.20 (s, 2H), 5.55 (s, 1H), 6.89-6.97 (m, 5H), 7.08-7.19 (m, 4H), 7.21-7.37 (m, 11H), 7.41-7.52 (m, 2H), 7.90-7.95 (m, 1H). The faster moving isomer was deprotected to give 737 mg (70%) of 5-[4'-[[[3-butyl-5-(dimethoxymethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole as a colorless solid: mp 152.5-153.5 °C; NMR (CDCl₃) δ 0.87 (t, J = 8 Hz, 3H), 1.20-1.30 (m, 2H), 1.53-1.65 (m, 2H), 2.46 (t, J = 8 Hz, 2H), 3.30 (s, 6H), 5.37 (s, 2H), 5.40 (s, 1H), 7.05-8.05 (m,

25

4H), 7.45 (dd, $J = 8$ and 2 Hz, 1H), 7.51-7.65 (m, 2H), 8.02
(d, $J = 8$ Hz, 1H); MS (FAB) m/e (rel intensity) 434 (65),
402 (40), 370 (11), 342 (46), 249 (85), 235 (18), 207
(100), 192 (89), 168 (70); HRMS. Calc'd for M+H:
5 434.2304. Found: 434.2332.

Example 24



5


5-[4'-[5-butyl-3-(dimethoxymethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

10

A 3.65 g (5.4 mmol) sample of the slower moving isomer from Example 23 was deprotected to give 108 mg of colorless 5-[4'-[5-butyl-3-(dimethoxymethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole after lyophilization: NMR (CDCl₃) δ 0.86 (t, J = 8 Hz, 3H), 1.24-1.39 (m, 2H), 1.56-1.70 (m, 2H), 2.66 (t, J = 8 Hz, 2H), 3.30 (s, 6H), 5.23 (s, 2H), 5.35 (s, 1H), 7.02 (d, J = 8 Hz, 2H), 7.10 (d, J = 8 Hz, 2H), 7.40 (dd, J = 8 and 2 Hz, 1H), 7.51 (dt, J = 8 and 2 Hz, 1H), 7.59 (dt, J = 8 and 2 Hz, 1H), 7.89 (dd, J = 8 and 2 Hz, 1H); MS (FAB) m/e (rel intensity) 434 (10), 402 (40), 249 (100), 207 (36), 192 (54), 168 (12); HRMS. Calc'd for M+H: 434.2304. Found: 434.2271.

20

Example 25



5

5-[4'-[[3-butyl-5-phenylethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl-2-yl]-1H-tetrazole

10

Following General Procedure A, 2.3 g (10 mmol) of 3-butyl-5-phenylethyl-1H-1,2,4-triazole was coupled with 6.6 g (10 mmol) of the alkylating reagent prepared in step 1 of Example 3 to give 2.67 g (38%) of a faster moving isomer: NMR (CDCl₃) δ 0.97 (t, J = 8 Hz, 3H), 1.36-1.50 (m, 2H), 1.71-1.84 (m, 2H), 2.73 (t, J = 8 Hz, 2H), 2.78-2.88 (m, 2H), 2.92-3.01 (m, 2H), 4.82 (s, 2H), 6.80-6.95 (m, 9H), 7.02-7.11 (m, 5H), 7.16-7.36 (m, 11H), 7.41-7.52 (m, 2H), 7.91-7.97 (m, 1H) and 2.94 g (42%) of a slower moving isomer: NMR (CDCl₃) δ 0.88 (t, J = 8 Hz, 3H), 1.23-1.38 (m, 2H), 1.57-1.70 (m, 2H), 2.58 (t, J = 8 Hz, 2H), 2.97-3.13 (m, 4H), 5.11 (s, 2H), 6.86 (d, J = 8 Hz, 2H), 6.89-6.96 (m, 6H), 7.10 (d, J = 8 Hz, 2H), 7.14-7.39 (m, 15H), 7.42-7.54 (m, 2H), 7.93 (dd, J = 8 and 2 Hz, 1H). The faster moving isomer was deprotected to give 1.6 g (88%) of

25


5-[4'-[[3-butyl-5-(2-phenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl-2-yl]-1H-tetrazole a colorless

solid: NMR (CDCl₃) δ 0.86 (t, J = 8 Hz, 3H), 1.17-1.31 (m, 2H), 1.48-1.51 (m, 2H), 2.42 (t, J = 8 Hz, 2H), 2.76 (t, J = 8 Hz, 2H), 2.92 (t, J = 8 Hz, 2H), 4.80 (s, 2H), 6.83 (d,

$J = 8$, 2H), 6.92-7.00 (m, 2H), 7.08 (d, $J = 8$ Hz, 2H),
7.17-7.32 (m, 3H), 7.42 (dd, $J = 8$ and 2 Hz, 1H), 7.51-7.65
(m, 2H), 7.96 (dd, $J = 8$ and 2 Hz, 1H); MS (FAB) m/e (rel
intensity) 464 (37), 230 (25), 207 (100), 178 (17); HRMS.

5 Calc'd for M+H: 464.2563. Found: 464.2532.

Example 26




5

5-[4'-[(5-butyl-3-phenylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl-2-yl]-1H-tetrazole

The slower moving isomer from Example 25 was
 10 deprotected to give 1.5 g (77%) of 5-[4'-[(5-butyl-3-(2-phenylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl-2-yl]-1H-tetrazole as a colorless solid: mp 156.0-157.2°C;
 NMR (CDCl₃) δ 0.84 (t, J = 8 Hz, 3H), 1.19-1.34 (m, 2H),
 1.47-1.61 (m, 2H), 2.52 (t, J = 8 Hz, 2H), 2.66-2.77 (m,
 15 2H), 2.85-2.95 (m, 2H), 5.15 (s, 2H), 6.88 (d, J = 8 Hz, 2H), 7.05-7.11 (m, 4H), 7.12-7.28 (m, 5H), 7.41 (dd, J = 8 and 2 Hz, 1H), 7.50 (dt, J = 8 and 2 Hz, 1H), 7.59 (dt, J = 8 and 2 Hz, 1H), 7.92 (dd, J = 8 and 2 Hz, 1H); MS (FAB) m/e (rel intensity) 464 (83), 230 (22), 207 (92); HRMS.
 20 Calc'd for M+H: 464.2563. Found: 464.2560.

Example 27




5

5-butyl-1-[2'-(1H-tetrazol-5-yl)][1,1'-biphenyl]-4-ylmethyl-1H-1,2,4-triazole-3-methanamine

A 1.5 g (3.5 mmol) sample of the diacetal
10 compound of Example 24 was dissolved in 40 ml of ethanol
and 40 ml of 3N HCl. The reaction was stirred at ambient
temperature overnight and then at reflux for 50 min, the
course of the reaction was followed by analytical reverse
phase HPLC. The solvent was removed in vacuo. The residue
15 was dissolved in ethyl acetate and extracted with saturated
sodium bicarbonate solution until the water layer stayed
basic. The water layer was acidified and extracted with an
ethyl acetate/methylene chloride mixture. The organic
layer was dried (MgSO₄) and removed in vacuo yielding 0.96
20 g of crude aldehyde: NMR (CDCl₃) δ 0.66 (t, J = 8 Hz, 3H),
1.06-1.20 (m, 2H), 1.42-1.54 (m, 2H), 2.53 (t, J = 8 Hz,
2H), 5.17 (s, 2H), 6.80-6.95 (m, 4H), 7.17-7.38 (m, 3H),
7.42-7.48 (m, 1H), 9.70 (s, 1H). Under a static nitrogen
atmosphere, 0.91 g (2.3 mmol) of this material was
25 dissolved in 10 ml of methanol and 1.77 g (23 mmol)
ammonium acetate was added followed by 0.10 g (1.61 mmol)
sodium cyanoborate. After stirring at ambient temperature
for 3 days, analytical reverse phase HPLC indicated that
the reaction was complete. Purification was accomplished

by reverse phase chromatography (Waters Delta Prep 3000) using 25% acetonitrile/water (0.05% TFA). The solvent from the pure fractions was removed in vacuo and the residue dissolved in methanol and 3N HCl. After stirring for 30 min, the solvent was removed in vacuo and the residue lyophilized from acetonitrile/water providing 100 mg (10%) of 5-butyl-1-[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]-1H-1,2,4-triazole-3-methanamine as the colorless hydrochloride salt: NMR (DMSO-d₆) δ 0.85 (t, J = 8 Hz, 3H), 1.22-1.38 (m, 2H), 1.51-1.64 (m, 2H), 2.77 (t, J = 8 Hz, 2H), 4.00-4.09 (m, 2H), 5.40 (s, 2H), 7.01-7.25 (m, 4H), 7.39 (s, 1H), 7.48-7.73 (m, 3H); MS (FAB) m/e (rel intensity) 389 (100), 207 (65), 192 (17); HRMS. Calc'd for M+H: 389.2202. Found: 389.2170.

Example 28



5

5-[4'-[(3-butyl-5-octyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

10 Following General Procedure A, 2.0 g (8.4 mmol) of 3-butyl-5-octyl-1H-1,2,4-triazole was coupled with 5.5 g (8.4 mmol) of the alkylating reagent prepared in step 1 of Example 3. Silica gel chromatography (Waters Prep 500-A) using 25% ethyl acetate/hexane produced 4.5 g (75.4%) of a


15 mixture of the two isomers which was dissolved in 40 ml of 10% water/acetic acid and stirred at ambient temperature overnight. The solvent was removed in vacuo. Purification of a sample of the isomeric product mix by reverse phase chromatography (Waters Delta Prep-3000) using 45%

20 acetonitrile/water (0.05% TFA) for 30 minutes followed by 50% acetonitrile/water (0.05% TFA) provided two isomers. The faster moving isomer compound was dissolved in dilute base, the water was acidified (pH 4-5) with 1N HCl, and the product extracted with ethyl acetate. The ethyl acetate

25 was dried (MgSO₄) and removed in vacuo yielding a solid which was recrystallized from acetonitrile providing 92 mg of 5-[4'-[(3-butyl-5-octyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole as a colorless solid: mp 136.5-138.0°C; NMR (CDCl₃) δ 0.82 (t,

$J = 8$ Hz, 3H), 0.86 (t, $J = 8$ Hz, 3H), 1.11-1.33 (m, 12H),
1.40-1.60 (m, 4H), 2.27 (t, $J = 8$ Hz, 2H), 2.44 (t, $J = 8$
Hz, 2H), 5.16 (s, 2H), 6.87 (d, $J = 8$ Hz, 2H), 7.07 (d, J
= 8 Hz, 2H), 7.44 (dd, $J = 8$ and 2 Hz, 1H), 7.50-7.66 (m,
5 2H), 7.90 (dd, $J = 8$ and 2 Hz, 1H); MS (FAB) m/e (rel
intensity) 472 (5), 207 (7); HRMS. Calc'd for M+H:
472.3189. Found: 472.3180.

Example 29




5

5-[4'-[(5-butyl-3-octyl-1H-1,2,4-triazol-1-yl)methyl]][1,1'-biphenyl]-2-yl]-1H-tetrazole

The slower moving isomer from Example 28 was
10 isolated in an identical manner and the product lyophilized
from acetonitrile/water providing 53 mg of 5-[4'-[(5-butyl-
3-octyl-1H-1,2,4-triazol-1-yl)methyl]][1,1'-biphenyl]-2-yl]-
1H-tetrazole as a colorless solid: NMR (CDCl₃) δ 0.83 (t,
 J = 8 Hz, 3H), 0.86 (t, J = 8 Hz, 3H), 1.12-1.37 (m, 12H),
15 1.46-1.63 (m, 4H), 2.35 (t, J = 8 Hz, 2H), 2.51 (t, J = 8
Hz, 2H), 5.19 (s, 2H), 6.92 (d, J = 8 Hz, 2H), 7.09 (d, J =
8 Hz, 2H), 7.40-7.48 (m, 1H), 7.51-7.66 (m, 2H), 7.91 (dd,
 J = 8 and 2 Hz, 1H); MS (FAB) m/e (rel intensity) 472 (6),
207 (8); HRMS. Calc'd for M+H: 472.3189. Found:
20 472.3230.

Example 30



5

1-[[5-butyl-1-[2'-(1H-tetrazol-5-yl)][1,1'-biphenyl]-4-ylmethyl]-1H-1,2,4-triazol-3-yl]methyl-1-butanone

10

Under a static nitrogen atmosphere, 30.0 g (150 mmol) 5-butyl-3-dimethoxy-1H-1,2,4-triazole (from Step 1 of Example 23) was added in small portions to 165 mmol of sodium hydride in 300 ml anhydrous THF; stirring was continued for 1 h. The anion solution was cooled to -10 °C

15

(methanol/ice) and treated with Sem-Cl dropwise. The reaction was allowed to warm to ambient temperature and stir overnight. The solvent was removed in vacuo. The residue was dissolved in methylene chloride, washed with water, dried (MgSO₄), and the solvent removed again in

20

vacuo. Silica gel chromatography using 20% ethyl acetate/hexane followed by ethyl acetate provided 13.6 g of the faster moving isomer as an oil: NMR (CDCl₃) δ - 0.07 (s, 9H), 0.82-0.92 (m, 5H), 1.26-1.40 (m, 2H), 1.62-1.74 (m, 2H), 2.67 (t, J = 8 Hz, 2H), 3.38 (s, 6H), 3.53-3.62 (m, 2H), 5.47 (s, 2H), 5.55 (s, 1H). Under a static


25

nitrogen atmosphere, 12.6 g (38.2 mmol) of the faster moving Sem-protected triazole isomer from above was dissolved in 630 ml of anhydrous THF, cooled to -78 °C, and

45.8 mmol of sec-butyl lithium was added dropwise. The solution was stirred for 1 h and then the anion was quenched with 4.5 ml (45.8 mmol) of n-propyl iodide. The solution was stirred at -78 °C for 5 h and then allowed to warm to ambient temperature overnight. The solvent was removed in vacuo. The residue was dissolved in methylene chloride, washed with water, dried (MgSO₄) and the solvent removed providing 12.4 g of crude Sem-protected 5-butyl-3-(1,1-dimethoxybutane)-1H-1,2,4-triazole: NMR (CDCl₃) δ -0.04 (s, 9H), 0.75-0.93 (m, 8H), 0.98-1.11 (m, 2H), 1.25-1.41 (m, 2H), 1.62-1.74 (m, 2H), 1.99-2.08 (m, 2H), 2.68 (t, J = 8 Hz, 2H), 3.20 (s, 6H), 3.62 (t, J = 8 Hz, 2H), 5.56 (s, 2H). A 1.0 g (2.7 mmol) sample of the crude dimethoxybutyl compound was dissolved in 5 ml of ethanol and 5 ml of 3M HCl and stirred at reflux for 2.5 h. The solvent was removed in vacuo providing 560 mg of the crude HCl salt of 5-butyl-3-(1-butanone)-1H-1,2,4-triazole: NMR (DMSO-d₆) δ 0.83-0.94 (m, 6H), 1.21-1.36 (m, 2H), 1.55-1.75 (m, 4H), 2.83 (t, J = 8 Hz, 2H), 2.97 (t, J = 8 Hz, 2H). The crude HCl salt of the triazole was dissolved in 20 ml of fresh methanol, 3 Å molecular sieves were added, and the mixture stirred at reflux under nitrogen overnight. The solution was filtered through celite and the solvent removed in vacuo. The residue was dissolved in ethyl acetate and washed with saturated sodium bicarbonate solution. The solvent was dried (MgSO₄) and removed in vacuo providing 390 mg of crude 5-butyl-3-(1,1-dimethoxybutyl)-1H-1,2,4-triazole: NMR (CDCl₃) δ 0.90 (t, J = 8 Hz, 3H), 0.99 (t, J = 8 Hz, 3H), 1.29-1.46 (m, 2H), 1.66-1.84 (m, 4H), 2.80 (t, J = 8 Hz, 2H), 3.06 (t, J = 8 Hz, 2H), 3.18 (s, 6H). Under a static nitrogen atmosphere, 380 mg (1.6 mmol) of this material in 5 ml of anhydrous dimethylformamide (DMF) was added to 1.9 mmol sodium hydride in 5 ml of DMF; stirring was continued for 1 h. The anion solution was cooled to 0 °C and 890 mg (1.6 mmol) of the alkylating reagent prepared in from step 1 of Example 3 was added as a solid. The reaction was allowed

to warm to ambient temperature and stir overnight.
Methanol (1 ml) was added to destroy any unreacted sodium
hydride and the DMF was removed in vacuo. The residue was
dissolved in ethyl acetate, washed with water, and dried
5 (MgSO₄). The solvent was removed in vacuo providing 1.2 g
of crude material which was dissolved in 10 ml ethanol and
10 ml 3N HCl and stirred at reflux for 2 h. The solvent
was removed in vacuo. Purification by reverse phase
chromatography (Water Delta Prep 3000) using isocratic 37%
10 acetonitrile/water (0.05% TFA) for 40 minutes followed by
50% acetonitrile/water (0.05% TFA) provided the TFA salt.
The salt was dissolved in basic water (pH 9-10), the water
was acidified to pH 4 with 3N HCl, and the product
15 extracted with ethyl acetate. The ethyl acetate was dried
(MgSO₄) and removed in vacuo yielding a solid which was
recrystallized from acetonitrile providing 146 mg (21%) of
1-[[5-butyl-1-[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-
ylmethyl]-1H-1,2,4-triazol-3-yl]methyl]-1-butanone mp
150.5-152.5°C; NMR (CDCl₃) δ 0.85 (t, J = 8 Hz, 3H), 0.96
20 (t, J = 8 Hz, 3H), 1.23-1.39 (m, 2H), 1.54-1.77 (m, 4H),
2.67 (t, J = 8 Hz, 2H), 2.97 (t, J = 8 Hz, 2H), 5.28 (s,
2H), 7.02 (d, J = 8 Hz, 2H), 7.10 (d, J = 8 Hz, 2H), 7.37
(d, J = 8 Hz, 1H), 7.43-7.62 (m, 2H), 7.86 (d, J = 8 Hz,
1H); MS (FAB) m/e (rel intensity) 430 (6), 207 (12); HRMS.
25 Calc'd for M+H: 430.2355. Found: 430.2404.

Example 31




5

N-[[5-butyl-1-[2'-(1H-tetrazol-5-yl)][1,1'-biphenyl]-4-ylmethyl]-1H-1,2,4-triazol-3-yl]methylacetamide

A 92.9 mg (0.22 mmol) sample of 5-[4'-[(5-butyl-3-methylamine-1H-1,2,4-triazol-1-yl)methyl]-(1,1'-biphenyl-2-yl)]-1H-tetrazole hydrochloride from Example 27 was dissolved in 5 ml of water and the pH was adjusted to 9 with 1M potassium carbonate (K_2CO_3). The solution was cooled to 0 °C and 0.22 ml of 1M K_2CO_3 was added followed by 0.22 mmol of acetic anhydride. Additional K_2CO_3 was added as needed to maintain a pH of 9. At 30 min intervals, this addition was repeated until analytical reverse phase chromatography showed that all starting material has been consumed. The pH was adjusted to three with 1N HCl and extracted with ethyl acetate. The ethyl acetate was removed in vacuo and the product lyophilized from acetonitrile/water providing 36 mg (39%) of N-[[5-butyl-1-[2'-(1H-tetrazol-5-yl)][1,1'-biphenyl]-4-ylmethyl]-1H-1,2,4-triazol-3-yl]methylacetamide as a colorless solid: NMR ($CDCl_3$) δ 0.80 (t, J = 8 Hz, 3H), 1.18-1.37 (m, 2H), 1.47-1.64 (m, 2H), 1.94 (s, 3H), 2.63 (t, J = 8 Hz, 2H), 4.22 (d, J = 8 Hz, 2H), 5.18 (s, 2H), 6.95 (d, J = 8

Hz, 2H), 7.04-7.21 (m, 2H), 7.44 (d, $J = 8$ Hz, 1H), 7.50-7.66 (m, 2H), 7.90 (dd, $J = 8$ and 2 Hz, 1H); MS (FAB) m/e (rel intensity) 431 (42), 207 (100); HRMS. Calc'd for M+H: 431.2308. Found: 431.2271.

Example 32



5

5-[4'-[(5-butyl-3-methyl-1H-1,2,4-triazol-1-yl)methyl]-
[1,1'-biphenyl]-2-yl]-1H-tetrazole


10

Following General Procedure A, 700 mg (5.0 mmol) of 5-butyl-3-methyl-1H-1,2,4-triazole was coupled with 2.8 g (5.0 mmol) of the alkylating reagent prepared in step 1 of Example 3. Silica gel chromatography using ethyl acetate/toluene (35:65) produced 2.75 g (90%) of a mixture of the two isomers which was dissolved in 30 ml 10% water/acetic acid and stirred at ambient temperature for 3 days. The solvent was removed in vacuo and the residue dissolved in dilute base, washed with toluene, acidified to pH 4 with 1N HCl and the product extracted with ethyl acetate. Purification of the isomeric product mixture by reverse phase chromatography (Water Delta Prep-3000) using isocratic acetonitrile/water (23:77) (0.05% TFA) provided two isomers. The faster moving isomer compound was dissolved in dilute base, acidified to pH 3-4 and extracted with ethyl acetate. The solvent was removed in vacuo; recrystallization from acetonitrile gave 199 mg (12%) of 5-[4'-[(5-butyl-3-propyl-1H-1,2,4-triazol-1-yl)methyl]-[1,1'-

25

biphenyl]-2-yl]-1H-tetrazole as a colorless solid: mp
170.0-170.5°C; NMR (CDCl₃) δ 0.79 (t, J = 8 Hz, 3H), 1.18-
1.32 (m, 2H), 1.44-1.56 (m, 2H), 1.99 (s, 3H), 2.49 (t, J =
8 Hz, 2H), 5.15 (s, 2H), 6.87 (d, J = 8 Hz, 2H), 7.08 (d, J
5 = 8 Hz, 2H), 7.44 (dd, J = 8 and 2 Hz, 1H), 7.51-7.65 (m,
2H), 7.90 (dd, J = 8 and 2 Hz, 1H); MS (FAB) m/e (rel
intensity) 374 (100), 235 (13), 207 (100), 140 (29); HRMS.
Calc'd for M+H: 374.2093. Found: 374.2062.

Example 33




5

5-[4'-[(3-butyl-5-methyl-1H-1,2,4-triazol-1-yl)methyl]-[1,1'-biphenyl]-1-yl]-1H-tetrazole

The slower moving isomer from Example 32 was dissolved in dilute base, acidified to pH 3-4, and extracted with ethyl acetate. The solvent was removed in vacuo and the product lyophilized from acetonitrile/water which gave 536 mg. (33 %) of 5-[4'-[(3-butyl-5-methyl-1H-1,2,4-triazol-1-yl)methyl]-[1,1'-biphenyl]-1-yl]-1H-tetrazole as a colorless solid: NMR (CDCl₃) δ 0.82 (t, J = 8 Hz, 3H), 1.21-1.35 (m, 2H), 1.53-1.66 (m, 2H), 2.35 (s, 3H), 2.56 (t, J = 8 Hz, 2H), 5.23 (s, 2H), 7.03 (d, J = 8 Hz, 2H), 7.11 (d, J = 8 Hz, 2H), 7.42 (dd, J = 8 and 2 Hz, 1H), 7.47-7.63 (m, 2H), 7.88 (dd, J = 8 and 2 Hz, 1H); MS (FAB) m/e (rel intensity) 374 (65), 331 (12), 235 (13), 207 (100), 192 (38), 140 (56); HRMS. Calc'd for M+H: 374.2093. Found: 374.2071.

Example 34



5

1,1-dimethylethyl [3-butyl-[1-2'-(1H-tetrazol-5-yl)[1,1'-
biphenyl]-4-ylmethyl]-1H-1,2,4-triazol-5-ylmethyl]carbamate

10

Following General Procedure A, 6.0 g (25 mmol) of [(5-butyl-1H-1,2,4-triazol-3-yl)methyl]carbamate was coupled with 16.4 g (25 mmol) of the alkylating reagent prepared in step 1 of Example 3. Silica gel chromatography (Waters Prep 500A) using ethyl acetate/hexane (2:3) gave


15 4.29 g (24%) of a faster moving isomer: NMR (CDCl₃) δ 0.94 (t, J = 8 Hz, 3H), 1.33-1.46 (m, 2H), 1.43 (s, 9H), 1.66-1.78 (m, 2H), 2.69 (t, J = 8 Hz, 2H), 4.26 (d, J = 8 Hz, 2H), 5.24 (s, 2H), 6.87-6.93 (m, 7H), 7.01 (d, J = 8 Hz, 2H), 7.11 (d, J = 8 Hz, 2H), 7.21-7.28 (m, 5H), 7.30-7.37

20 (m, 4H), 7.42-7.52 (m, 2H), 7.94 (dd, J = 8 and 2 Hz, 1H) and 9.74 g (54%) of a slower moving isomer: NMR (CDCl₃) δ 0.87 (t, J = 8 Hz, 3H), 1.24-1.36 (m, 2H), 1.44 (s, 9H), 1.57-1.69 (m, 2H), 2.58 (t, J = 8 Hz, 2H), 4.38 (d, J = 8 Hz, 2H), 5.12 (s, 2H), 6.91 (d, J = 8 Hz, 8H), 7.11 (d, J =

25 8 Hz, 2H), 7.21-7.28 (m, 6H), 7.30-7.37 (m, 4H), 7.42-7.52 (m, 2H), 7.91-7.96 (m, 1H). The faster moving isomer was dissolved in 30 ml of water/acetic acid (1:4) and stirred at ambient temperature overnight. The solvent was removed

in vacuo; the residue dissolved in dilute base and washed with toluene. The water layer was acidified to pH 3-4 and extracted with ethyl acetate. Recrystallization from acetonitrile gave 2.53 g (88%) of 1,1-dimethylethyl[3-butyl-5-[1-[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-ylmethyl]-1H-1,2,4-triazol-5-yl]methylcarbamate as a colorless solid: mp 144-147°C; NMR (CDCl₃) δ 0.88 (t, J = 8 Hz, 3H), 1.22-1.45 (m, 2H), 1.36 (s, 9H), 1.54-1.66 (m, 2H), 2.56 (t, J = 8 Hz, 2H), 4.05 (d, J = 8 Hz, 2H), 5.38 (s, 2H), 5.50 (s, 1H), 7.06 (d, J = 8 Hz, 2H), 7.12 (d, J = 8 Hz, 2H), 7.45 (dd, J = 8 and 2 Hz, 1H), 7.50-7.63 (m, 2H), 7.99 (d, J = 8 Hz, 1H); MS (FAB) m/e (rel intensity) 495 (12), 395 (38), 367 (15), 207 (100), 178 (42); HRMS. Calc'd for M+Li: 495.2808. Found: 495.2771.

Example 35



5

1,1-dimethylethyl[5-butyl-[1-[2'-(1H-tetrazol-5-yl)][1,1'-biphenyl]-4-ylmethyl]-1H-1,2,4-triazol-3-yl]methylcarbamate

10


The slower moving isomer from Example 34 was dissolved in 70 ml of water/acetic acid (1:4) and stirred at ambient temperature overnight. The solvent was removed in vacuo; the residue dissolved in dilute base and washed with toluene. The water layer was acidified to pH 4 with 1N HCl and extracted with ethyl acetate. The ethyl acetate was dried (MgSO₄) and removed in vacuo. Lyophilization

15

from acetonitrile/water gave 5.0 g (76%) of 1,1-dimethylethyl[5-butyl-[1-[2'-(1H-tetrazol-5-yl)][1,1'-biphenyl]-4-ylmethyl]-1H-1,2,4-triazol-3-yl]methylcarbamate as a colorless solid: NMR (CDCl₃) δ 0.77 (t, J = 8 Hz, 3H), 1.16-1.30 (m, 2H), 1.40 (s, 9H), 1.42-1.56 (m, 2H), 2.57 (t, J = 8 Hz, 2H), 4.12 (d, J = 8 Hz, 2H), 5.19 (s, 2H), 5.59 (s, 1H), 6.94 (d, J = 8 Hz, 2H), 7.08 (d, J = 8 Hz, 2H), 7.43 (dd, J = 8 and 2 Hz, 1H), 7.50-7.63 (m, 2H), 7.92 (dd, J = 8 and 2 Hz, 1H); MS (FAB) m/e (rel intensity) 495 (29), 367 (38), 338 (21), 207 (100), 178 (48); HRMS: Calc'd for M+Li: 495.2808. Found: 495.2800.

25

Example 36




5

3-butyl-1-[2'-(1H-tetrazol-5-yl)][1,1'-biphenyl]-4-ylmethyl-1H-1,2,4-triazole-5-methanamine

Under nitrogen, 219 mg (0.45 mmol) of 1,1-dimethylethyl [3-butyl-[1-[2'-(1H-tetrazol-5-yl)][1,1'-biphenyl]-4-ylmethyl]-1H-1,2,4-triazol-5-yl] methylcarbamate from Example 34 was suspended in 10 ml of 4N HCl in dioxane. After 30 minutes, 10 ml of methylene chloride was added. The mixture was stirred at ambient temperature overnight. The solvent was removed in vacuo. Lyophilization from acetonitrile/water gave 201 mg (100%) of 3-butyl-1-[2'-(1H-tetrazol-5-yl)][1,1'-biphenyl]-4-ylmethyl-1H-1,2,4-triazole-5-methanamine as the colorless hydrochloride salt: NMR (CDCl₃) δ 0.82 (t, J = 8 Hz, 3H), 1.21-1.35 (m, 2H), 1.59-1.72 (m, 2H), 2.72 (t, J = 8 Hz, 2H), 4.97 (s, 2H), 5.69 (s, 2H), 6.93 (d, J = 8 Hz, 2H), 7.23 (d, J = 8 Hz, 2H), 7.32 ((d, J = 8 Hz, 1H), 7.39 (dt, J = 8 and 2 Hz, 1H), 7.51 (dt, J = 8 and 2 Hz, 1H), 7.74 (dd, J = 8 and 2 Hz, 1H); MS (FAB) m/e (rel intensity) 389 (55), 207 (100), 192 (39), 178 (26); HRMS. Calc'd for M+H: 389.2202. Found: 389.2170.

Example 37



5

5-[4'-[[3-butyl-5-(2,2-diethoxyethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

10

Following General Procedure A, 4.7 g (19.5 mmol) of 5-butyl-3-(2,2-diethoxyethyl)-1H-1,2,4-triazole was coupled with 19.5 mmol of the alkylating reagent prepared in step 1 of Example 3. Silica gel chromatography (Waters Prep 500A) using ethyl acetate/hexane (2:3) gave 5.98 g (43%) of a faster moving isomer: NMR (CDCl₃) δ 0.94 (t, J = 8 Hz, 3H), 1.13 (t, J = 8 Hz, 6H), 1.33-1.48 (m, 2H), 1.67-1.79 (m, 2H), 2.70 (t, J = 8 Hz, 2H), 2.94 (d, J = 8 Hz, 2H), 3.39-3.51 (m, 2H), 3.63-3.75 (m, 2H), 4.81 (t, J = 7 Hz, 1H), 5.22 (s, 2H), 6.87-6.98 (m, 8H), 7.09 (d, J = 8 Hz, 2H), 7.21-7.29 (m, 6H), 7.30-7.37 (m, 4H), 7.41-7.52 (m, 2H), 7.91-7.96 (m, 1H) and 6.3 g (45%) of a slower moving isomer: NMR (CDCl₃) δ 0.88 (t, J = 8 Hz, 3H), 1.15 (t, J = 7 Hz, 6H), 1.24-1.37 (m, 2H), 1.56-1.68 (m, 2H), 2.57 (t, J = 8 Hz, 2H), 3.06 (d, J = 7 Hz, 2H), 3.46-3.60 (m, 2H), 3.66-3.78 (m, 2H), 5.05 (t, J = 7 Hz, 1H), 5.14 (s, 2H), 6.87-6.95 (m, 8H), 7.10 (d, J = 8 Hz, 2H), 7.21-7.29 (m, 6H), 7.31-7.37 (m, 4H), 7.42-7.53 (m, 2H), 7.90-7.96 (m, 1H). The faster moving isomer was dissolved in 50 ml of acetic acid/water (4:1) and stirred at ambient


temperature for 3 days. The solvent was removed in vacuo; the residue dissolve in dilute base and washed with toluene. The water layer was cooled to 0 °C, acidified to pH 4-5 and extracted with ethyl acetate. Lyophilization

5 from acetonitrile/water gave 3.8 g (98%) of 5-[4'-[[3-butyl-5-(2,2-diethoxyethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole as a colorless solid: NMR (CDCl₃) δ 0.80 (t, J = 8 Hz, 3H), 1.10 (t, J = 8 Hz, 6H), 1.10-1.24 (m, 2H), 1.37-1.49 (m,

10 2H), 2.31 (t, J = 8 Hz, 2H), 2.66 (d, J = 7 Hz, 2H), 3.34-3.47 (m, 2H), 3.59-3.71 (m, 2H), 4.75 (t, J = 7 Hz, 1H), 5.28 (s, 2H), 6.88 (d, J = 8 Hz, 2H), 7.06 (d, J = 8 Hz, 2H), 7.46 (dd, J = 8 and 2 Hz, 1H), 7.55 (dt, J = 8 and 2 Hz, 1H), 7.62 (dt, J = 8 and 2 Hz, 1H), 7.91 (dd, J = 8 and

15 2 Hz, 1H); MS (FAB) m/e (rel intensity) 476 (20), 430 (28), 356 (13), 235 (15), 207 (100), 192 (67); HRMS. Calc'd for M+H: 476.2773. Found: 476.2723.

Example 38




5

5-[4'-[[5-butyl-3-(2,2-diethoxyethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

The 6.2 g (8.6 mmol) of the slower moving isomer from Example 37 was dissolved in 50 ml of acetic acid/water (4:1) and stirred at ambient temperature for 3 days. The solvent was removed *in vacuo*; the residue dissolved in dilute base, and washed with toluene. The water layer was cooled to 0 °C, acidified with 1N HCl to pH 4-5, and extracted with ethyl acetate. Lyophilization from acetonitrile/water gave 3.7 g (92%) of 5-[4'-[[5-butyl-3-(2,2-diethoxyethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole as colorless solid: NMR (CDCl₃) δ 0.85 (t, J = 8 Hz, 3H), 1.09 (t, J = 8 Hz, 6H), 1.24-1.38 (m, 2H), 1.51-1.63 (m, 2H), 2.55 (t, J = 8 Hz, 2H), 2.75 (d, J = 7 Hz, 2H), 3.38-3.50 (m, 2H), 3.54-3.66 (m, 2H), 4.87 (t, J = 7 Hz, 1H), 5.19 (s, 2H), 6.99 (d, J = 8 Hz, 2H), 7.12 (d, J = 8 Hz, 2H), 7.42 (dd, J = 8 and 2 Hz, 1H), 7.50-7.65 (m, 2H), 7.97 (dd, J = 8 and 2 Hz, 1H); MS (FAB) m/e (rel intensity) 476 (3), 430 (28), 235 (10), 207 (100), 192 (61); HRMS. Calc'd for M+H: 476.2773. Found: 476.2760.

Example 39

5




5-[4'-[(5-butyl-3-ethyl-1H-1,2,4-triazol-1-yl)methyl]]-1,1'-biphenyl]-2-yl]-1H-tetrazole

10

Following General Procedure A, 770 mg (5 mmol) of 5-butyl-3-ethyl-1H-1,2,4-triazole was coupled with 2.8 g (5 mmol) of the alkylating reagent prepared in step 1 of Example 3. Silica gel chromatography (Waters Prep-500A) using ethyl acetate/hexane (2:3) gave 2.24 g (72%) of a mixture of the two isomers which was dissolved in 30 ml of acetic acid/water (9:1) and stirred at ambient temperature for 4 days. The solvent was removed in vacuo; the residue dissolved in dilute base, and washed with toluene. The water layer was acidified to pH 4 and extracted with ethyl acetate. The extracts were combined, dried (MgSO₄) and concentrated in vacuo. Purification of a 600 mg sample of the isomeric product mixture by reverse phase chromatography (Waters Delta Prep-3000) using isocratic acetonitrile/water (25:75) (0.05% TFA) provided two isomers. The faster moving isomer was dissolved in dilute base, acidified, extracted with ethyl acetate, dried (MgSO₄) and the ethyl acetate removed in vacuo.

Lyophilization from acetonitrile/water gave 209 mg of 5-[4'-[(5-butyl-3-ethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole as a colorless solid: NMR (CDCl₃) δ 0.88 (t, J = 8 Hz, 3H), 1.33-1.40 (m, 2H), 1.38 (t, J = 8 Hz, 3H), 1.58-1.70 (m, 2H), 2.67-2.85 (m, 4H), 5.26 (s, 2H), 7.10 (s, 4H), 7.41 (dd, J = 8 and 2 Hz, 1H), 7.74-7.62 (m, 2H), 7.86 (dd, J = 8 and 2 Hz, 1H); MS (FAB) m/e (rel intensity) 388 (58), 207 (100), 192 (35); HRMS. Calc'd for M+H: 388.2249. Found: 388.2218.

Example 40




5

5-[4'-[(3-butyl-5-ethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

The slower moving isomer from Example 39 was dissolved in dilute base, acidified to pH 3-4, extracted with ethyl acetate, dried (MgSO₄), and the ethyl acetate removed in vacuo. Lyophilization from acetonitrile/water gave 188 mg of 5-[4'-[(3-butyl-5-ethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole as a colorless solid: NMR (CDCl₃) δ 0.87 (t, J = 8 Hz, 3H), 1.16 (t, J = 8 Hz, 3H), 1.21-1.35 (m, 2H), 1.52-1.65 (m, 2H), 2.50 (t, J = 8 Hz, 2H), 2.67 (q, J = 8 Hz, 2H), 5.23 (s, 2H), 7.02 (d, J = 8 Hz, 2H), 7.12 (d, J = 8 Hz, 2H), 7.43 (dd, J = 8 and 2 Hz, 1H), 7.50-7.65 (m, 2H), 7.89 (dd, J = 8 and 2 Hz, 1H); MS (FAB) m/e (rel intensity) 388 (35), 207 (100), 192 (47); HRMS. Calc'd for M+H: 388.2249. Found: 388.2222.

Example 41



5

N2-acetyl-N-[[5-butyl-1-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-ylmethyl]-1H-1,2,4-triazol-3-yl]methyl]-L-glutamine, 1,1-dimethylethyl ester


10

Under nitrogen, 30 ml of TFA was added dropwise to a solution of 4.9 g (10 mmol) of 1,1dimethylethyl [5-butyl-[1-[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-ylmethyl]-1H-1,2,4-triazol-3-yl]methyl]carbamate from Example 35 in 30 ml of methylene chloride at -10°C (methanol/ice). The mixture was allowed to warm to ambient temperature and stir overnight. The solvent was removed in vacuo giving the TFA salt of the free amine. The TFA salt was dissolved in 35 ml of anhydrous DMF along with 8.7 ml (50 mmol) of anhydrous diisopropyl ethyl amine and treated with 20 mmol of the symmetrical anhydride of N-Boc-γ-glutamic acid t-butyl ester in 55 ml of anhydrous DMF. The reaction was stirred at ambient temperature overnight. The DMF was removed in vacuo; the residue dissolved in ethyl acetate, washed with cold 1M K₂CO₃ and water. The ethyl acetate was dried (MgSO₄) and the solvent removed in vacuo. Under nitrogen, 30 ml of TFA was added dropwise to a solution of this material in 30 ml of methylene chloride at -10 °C

25

(methanol/ice). The reaction was allowed to warm to ambient temperature and stir overnight. The solvent was removed in vacuo giving the TFA salt of the free amine. The TFA salt was dissolved in 30 ml of water and cooled to 0 °C. The pH was adjusted to nine with 1M K₂CO₃. The solution was cooled to 0 °C, 0.94 ml (10 mmol) of acetic anhydride was added followed by 5 ml of 1M K₂CO₃, and the pH was adjusted to 9 with additional 1M K₂CO₃. At 30 minute intervals, this addition was repeated until 5 additions had been made. The pH was adjusted to 4 with 6N HCl and the solution extracted with ethyl acetate. The ethyl acetate was dried (MgSO₄) and removed in vacuo. Purification by silica gel chromatography (Waters Prep 500A) using isopropanol/acetic acid/chloroform (20:5:75) followed by lyophilization from acetonitrile/water gave 2.5 g (41%) of N2-acetyl-N-[[5-butyl-1-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl)methyl]-1H-1,2,4-triazol-3-yl)methyl]-L-glutamine, 1,1-dimethylethyl ester as a colorless solid: NMR (CDCl₃) δ 0.89 (t, J = 8 Hz, 3H), 1.27-1.41 (m, 2H), 1.52-1.71 (m, 2H), 1.56 (s, 9H), 2.00 (s, 3H), 2.05-2.29 (m, 2H), 2.31-2.55 (m, 2H), 2.73 (t, J = 8 Hz, 2H), 4.48-4.58 (m, 3H), 5.24 (s, 2H), 7.09 (d, J = 8 Hz, 2H), 7.17 (d, J = 8 Hz, 2H), 7.36-7.42 (m, 1H), 7.44-7.55 (m, 2H), 7.90 (dd, J = 8 and 2 Hz, 1H); MS (FAB) m/e (rel intensity) 616 (20), 263 (23), 235 (26), 207 (100); HRMS. Calc'd for M+H: 616.3360. Found: 616.3353.

Example 42



5

5-[4'-[[3-butyl-5-(2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole


Following General Procedure A, 1.2 g (5 mmol) of 5-butyl-3-(2-cyclohexylethyl)-1H-1,2,4-triazole was coupled
 10 with 5 mmol of the alkylating reagent prepared in step 1 of Example 3. Silica gel chromatography (Waters Prep 500A) using ethyl acetate/hexane (1:3) gave 3.6 g (100%) of a mixture of the two isomers which was dissolved in 50 ml of
 15 acetic acid/water (9:1) and stirred at ambient temperature overnight. The solvent was removed in vacuo; the residue dissolved in dilute base, acidified and washed with ethyl acetate. The ethyl acetate was removed in vacuo.
 Purification of a small sample of the isomer product
 20 mixture by reverse phase chromatography (Water Delta Prep-3000) using isocratic acetonitrile/water (41:59) (0.05% TFA) provided two isomers. The faster moving isomer was dissolved in dilute base, acidified to pH 3-4, extracted with ethyl acetate, dried (MgSO₄) and the ethyl acetate
 25 removed in vacuo. Lyophilization from acetonitrile/water gave 58 mg of 5-[4'-[[3-butyl-5-(2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole as a colorless solid: NMR (CDCl₃) δ 0.78-0.95 (m, 5H), 1.07-1.40 (m, 6H), 1.48-1.74 (m, 9H), 2.66 (t, J =

211

8 Hz, 2H), 2.78 (t, J = 8 Hz, 2H), 5.24 (s, 2H), 7.10 (q, J = 8 Hz, 4H), 7.41 (dd, J = 8 and 2 Hz, 1H), 7.48-7.63 (m, 2H), 7.89 (dd, J = 8 and 2 Hz, 1H); MS (FAB) m/e (rel intensity) 470 (45), 236 (35), 207 (100), 192 (33); HRMS.

5 Calc'd for $M+H$: 470.3032. Found: 470.2990.

Example 43




5

5-[4'-[[5-butyl-3-(2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

The slower moving isomer from Example 42 was dissolved in dilute base, acidified to pH 3-4, extracted with ethyl acetate, dried (MgSO₄), and the ethyl acetate removed in vacuo. Lyophilization from acetonitrile/water gave 29 mg of 5-[4'-[[5-butyl-3-(2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole as a colorless solid: NMR (CDCl₃) δ 0.95-1.00 (m, 5H), 1.10-1.43 (m, 6H), 1.59-1.77 (m, 9H), 2.83 (t, $J = 8$ Hz, 2H), 2.92 (t, $J = 8$ Hz, 2H), 5.31 (s, 2H), 7.10-7.18 (m, 4H), 7.40 (dd, $J = 8$ and 2 Hz, 1H), 7.47-7.61 (m, 2H), 7.89 (dd, $J = 8$ and 2 Hz, 1H); MS (FAB) m/e (rel intensity) 470 (35), 236 (33), 207 (100), 192 (33); HRMS. Calc'd for M+H: 470.3032. Found: 470.2994.

Example 44



5

5-[4'-[[5-butyl-3-(phenylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl-2-yl]-1H-tetrazole

10 Under nitrogen, 0.74 g (2.2 mmol) of N2-[4-[(2-cyanophenyl)phenyl]methyl] phenyl acetic acid hydrazide was dissolved in 5 ml of absolute ethanol and treated with 0.28 g (2.2 mmol) of ethyl iminovalerate. The reaction was stirred at reflux for 3 days. The solvent was removed in vacuo, 5 ml of xylene was added and the reaction stirred at reflux under nitrogen for an additional 3 days. The solvent was removed in vacuo. Purification by silica gel chromatography (Chromatatron, 4 mm plate) using a step gradient of ethyl acetate/chloroform gave 410 g (31%) of 5-[4'-[[5-butyl-3-(phenylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl-2-yl]nitrile: NMR (CDCl₃) δ 0.88 (t, J = 8 Hz, 3H), 1.27-1.41 (m, 2H), 1.59-1.72 (m, 2H), 2.67 (t, J = 8 Hz, 2H), 4.06 (s, 2H), 5.30 (s, 2H), 7.14-7.37 (m, 8H), 7.39-7.50 (m, 3H), 7.53 (dt, J = 8 and 2 Hz, 1H), 7.75 (dd, J = 8 and 2 Hz, 1H). Under nitrogen, 230 mg (1.13 mmol) of trimethyl tin azide was added to a solution of the nitrile in 5 ml of xylene. The reaction was stirred at reflux for 3 days. The solvent was removed in vacuo and

the residue dissolved in 10 ml of acetic acid/water (9:1) and stirred at ambient temperature overnight. The solvent was removed in vacuo. Reverse phase chromatography (Delta Prep 3000) using acetonitrile/water (35-45:65-55) gave the


5 TFA salt. The salt was dissolved in dilute base, acidified to pH 3-4, extracted with ethyl acetate, dried (MgSO₄) and the ethyl acetate removed in vacuo. Lyophilization from acetonitrile/water gave 182 mg (45%) of 5-[4'-[[5-butyl-3-(phenylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-

10 biphenyl]-2-yl]-1H-tetrazole as a colorless solid: NMR (CDCl₃) δ 0.84 (t, J = 8 Hz, 3H), 1.22-1.36 (m, 2H), 1.51-1.64 (m, 2H), 2.59 (t, J = 8 Hz, 2H), 3.85 (s, 2H), 5.20 (s, 2H), 7.02 (d, J = 8 Hz, 2H), 7.14 (d, J = 8 Hz, 2H), 7.17-7.28 (m, 5H), 7.43 (dd, J = 8 and 2 Hz, 1H), 7.51-7.65

15 (m, 2H), 7.96 (dd, J = 8 and 2 Hz, 1H); MS (FAB) m/e (rel intensity) 450 (56), 207 (100), 192 (30), 178 (22); HRMS. Calc'd for M+H: 450.2406. Found: 450.2434.

215

Example 45




5

5-butyl-1-[2'-(1H-tetrazol-5-yl)][1,1'-biphenyl]-4-ylmethyl]-1H-1,2,4-triazole-3-propanol

Following General Procedure A, 920 mg (5.0 mmol) of 5-butyl-1H-1,2,4-triazole-3-propanol was coupled with 5.0 mmol of the alkylating reagent prepared in step 1 of Example 3. Silica gel chromatography (Waters Prep 500A) using isopropanol/chloroform (5:95) gave a mixture of the two pure isomers which was dissolved in 40 ml acetic acid/water (9:1) and stirred at ambient temperature for 3 days. The solvent was removed in vacuo. Purification of an aliquot by reverse phase chromatography (Waters Delta Prep 3000) using acetonitrile/water (26:74) (0.05% TFA) provided two isomers. Lyophilization of the faster moving isomer from acetonitrile/ water gave 124 mg of 5-butyl-1-[2'-(1H-tetrazol-5-yl)][1,1'-biphenyl]-4-ylmethyl]-1H-1,2,4-triazole-3-propanol as the colorless FTA salt: NMR (CDCl₃) δ 0.71 (t, J = 8 Hz, 3H), 1.10-1.24 (m, 2H), 1.42-1.54 (m, 2H), 1.70-1.81 (m, 2H), 2.49 (t, J = 8 Hz, 2H), 2.60 (t, J = 8 Hz, 2H), 3.46 (t, J = 6 Hz, 2H), 5.02 (s, 2H), 6.92 (q, J = 8 Hz, 4H), 7.24-7.34 (m, 2H), 7.40 (dt, J = 8 and 2 Hz, 1H), 7.52 (dd, J = 8 and 2 Hz, 1H); MS (FAB) m/e (rel intensity) 418 (100), 207 (48), 184 (10); HRMS. Calc'd for M+H: 418.2355. Found: 418.2370.

Example 46




5

3-butyl-1-[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-1-ylmethyl]-1H-1,2,4-triazole-5-propanol

Lyophilization of the slower moving isomer from
 10 Example 45 gave 114 mg of 3-butyl-1-[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-1-ylmethyl]-1H-1,2,4-triazole-5-propanol as the colorless TFA salt: NMR (CDCl₃) δ 0.75 (t, J = 8 Hz, 3H), 1.13-1.27 (m, 2H), 1.48-1.60 (m, 2H), 1.64-1.75 (m, 2H), 2.52 (t, J = 8 Hz, 2H), 2.66 (t, J = 8 Hz, 2H),
 15 3.39 (t, J = 6 Hz, 2H), 5.12 (s, 2H), 6.94 (s, 4H), 7.25-7.35 (m, 2H), 7.40 (dt, J = 8 and 2 Hz, 1H), 7.52 (d, J = 8 Hz, 1H); MS (FAB) m/e (rel intensity) 418 (100), 207 (30); HRMS. Calc'd for M+H: 418.2355. Found: 418.2344.

Example 47




5

5-[4'-[[5-butyl-3-(difluoromethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

10 Following General Procedure A, 470 mg (2.7 mmol)
of 5-butyl-3-(difluoromethyl)-1H-1,2,4-triazole was coupled
with 2.7 mmol of the alkylating reagent prepared in step 1
of Example 3. Purification by reverse phase chromatography
(Waters Delta Prep 3000) using acetonitrile/water (36:64)
15 (0.05% TFA) gave two isomers. The faster moving isomer was
dissolved in dilute base, acidified to pH 3-4, extracted
with ethyl acetate, dried (MgSO₄), and the ethyl acetate
removed in vacuo. Lyophilization from acetonitrile/water
gave 471 mg (27%) of 5-[4'-[[5-butyl-3-(difluoromethyl)-1H-
20 1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-
tetrazole as a colorless solid: NMR (CDCl₃) δ 0.88 (t, J =
8 Hz, 3H), 1.28-1.43 (m, 2H), 1.60-1.73 (m, 2H), 2.72 (t, J
= 8 Hz, 2H), 5.32 (s, 2H), 6.61 (t, J = 54 Hz, 1H), 7.12
(d, J = 8 Hz, 2H), 7.18 (d, J = 8 Hz, 2H), 7.42 (dd, J = 8
and 2 Hz, 1H), 7.51-7.65 (m, 2H), 8.00 (dd, J = 8 and 2 Hz,
25 1H); MS (FAB) m/e (rel intensity) 410 (45), 235 (55), 207
(100), 192 (57); HRMS. Calc'd for M+H: 410.1905. Found:

410.1903.

Example 48




5

5-[4'-[[[3-butyl-5-(difluoromethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

The slower moving isomer from Example 47 was dissolved in dilute base, acidified, extracted with ethyl acetate, dried (MgSO₄), and the ethyl acetate removed in vacuo. Lyophilization from acetonitrile/water gave 104 mg (6%) of 5-[4'-[[[3-butyl-5-(difluoromethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole as a colorless solid: NMR (CDCl₃) δ 0.91 (t, J = 8 Hz, 3H), 1.25-1.41 (m, 2H), 1.60-1.73 (m, 2H), 2.64 (t, J = 8 Hz, 2H), 5.43 (s, 2H), 6.81 (t, J = 52 Hz, 1H), 7.20 (d, J = 8 Hz, 2H), 7.28 (d, J = 8 Hz, 2H), 7.43 (dd, J = 8 and 2 Hz, 1H), 7.52-7.65 (m, 2H), 8.10 (dd, J = 8 and 2 Hz, 1H); MS (FAB) m/e (rel intensity) 410 (87), 235 (100), 207 (83); HRMS. Calc'd for M+H: 410.1905. Found: 410.1903.

Example 49




5

5-butyl-1-[2'-(1H-tetrazol-5-yl)][1,1'-biphenyl]-4-ylmethyl]-1H-1,2,4-triazole-3-carboxaldehyde

A 92 mg sample of 5-[4'-[[5-butyl-3-(dimethoxymethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole from Example 24 was dissolved in 20 ml of ethanol and 2 ml of 3N HCl. The reaction was stirred at ambient temperature for 3 days. The solvent was removed in vacuo. Lyophilization from acetonitrile/water gave 70 mg (81%) of 5-butyl-1-[2'-(1H-tetrazol-5-yl)][1,1'-biphenyl]-4-ylmethyl]-1H-1,2,4-triazole-3-carboxaldehyde as the colorless hydrochloride salt: NMR (CDCl₃) δ 0.89 (t, J = 8 Hz, 3H), 1.30-1.44 (m, 2H), 1.66-1.78 (m, 2H), 2.78 (t, J = 8 Hz, 2H), 5.38 (s, 2H), 7.17 (q, J = 8 Hz, 4H), 7.40 (d, J = 8 Hz, 1H), 7.50-7.64 (m, 2H), 8.02 (d, J = 8 Hz, 1H); MS (FAB) m/e (rel intensity) 388 (36), 235 (42), 207 (100), 192 (50); HRMS. Calc'd for M+Li: 395.2046. Found: 395.2049.

Example 50




5

5-[4'-[5-butyl-3-(phenylacetyl)-1H-1,2,4-triazol-1-yl]methyl]biphenyl-2-yl-1H-tetrazole

Under nitrogen, 7.3 mmol of sec-butyl lithium was added dropwise to 2.0 g (6.1 mmol) of the Sem-protected 5-butyl-3-(dimethoxymethyl)-1H-1,2,4-triazole from Example 30 in 100 ml of anhydrous THF at -78 °C. The reaction was stirred for 1 h and then quenched with 0.87 ml (7.3 mmol) of benzyl bromide. The reaction was allowed to warm to ambient temperature and stir overnight. The THF was removed in vacuo; the residue dissolved in methylene chloride, washed with water and dried (MgSO₄). The methylene chloride was removed in vacuo. The residue was dissolved in 10 ml of ethanol and 10 ml of 3M HCl and stirred at reflux for 2.5 h. The solvents were removed in vacuo. The residue was dissolved in ethyl acetate, washed with saturated sodium bicarbonate solution, and dried (MgSO₄). Silica gel chromatography (Waters Prep 500A) using ethyl acetate/hexane (35:65) gave 460 mg (31%) of 5-butyl-3-(phenylacetyl)-1H-1,2,4-triazole: NMR (CDCl₃) δ 0.92 (t, J = 8 Hz, 3H), 1.31-1.55 (m, 2H), 1.70-1.83 (m, 2H), 2.88 (t, J = 8 Hz, 2H), 4.39 (s, 2H), 7.21-7.38 (m,

5H). Following General Procedure A, 440 mg (1.8 mmol) of this material was reacted with 1.8 mmol of the alkylating reagent prepared in step 1 of Example 3. The crude product was dissolved in 10 ml of acetic acid/water (9:1) and stirred at ambient temperature for 4 days. The solvent was removed in vacuo; the residue dissolved in dilute base and washed with toluene. The water was acidified and extracted with ethyl acetate. Purification by reverse phase chromatography (Waters Delta Prep 3000) using acetonitrile/water (45:55) gave 50 mg (5%) of 5-[4'-[[5-butyl-3-(phenylacetyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole as the colorless TFA salt: NMR (DMSO-d₆) δ 0.86 (t, J = 8 Hz, 3H), 1.23-1.38 (m, 2H), 1.52-1.65 (m, 2H), 2.80 (t, J = 8 Hz, 2H), 4.32 (s, 2H), 5.52 (s, 2H), 7.11 (d, J = 8 Hz, 2H), 7.19 (d, J = 8 Hz, 2H), 7.23-7.35 (m, 5H), 7.52-7.63 (m, 2H), 7.65-7.74 (m, 2H); MS (FAB) m/e (rel intensity) 478 (100), 244 (8), 235 (8), 207 (34), 192 (12); HRMS. Calc'd for M+H: 478.2355. Found: 478.2414.

Example 51



5

5-[4'-[[5-butyl-3-(1,1-difluoro-2-phenylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-1-yl]-1H-tetrazole

10 Under nitrogen, 8.5 g (48 mmol) of 5-butyl-3-(difluoromethyl)-1H-1,2,4-triazole from Example 47 was added in portions to 53 mmol of sodium hydride in 85 ml of anhydrous THF; stirring was continued for 1 h. The anion solution was cooled to 0 °C and treated dropwise with 8.9
15 ml (50 mmol) of Sem-Cl. The reaction was allowed to warm to ambient temperature and stir overnight. The THF was removed in vacuo; the residue dissolved in methylene chloride, washed with water, and dried (MgSO₄).

Purification by silica gel chromatography (Waters Prep
20 500A) using ethyl acetate/hexane (1:9) gave 4.9 g 34% of the Sem-protected triazole as an oil: NMR (CDCl₃) δ -0.04 (s, 9H), 0.86-0.95 (m, 5H), 1.39-1.43 (m, 2H), 1.65-1.76 (m, 2H), 2.71 (t, J = 8 Hz, 2H), 3.62 (t, J = 8 Hz, 2H), 5.53 (s, 2H), 6.82 (t, J = 52 Hz, 1H). Under nitrogen,
25 14.2 mmol of sec-butyllithium was added dropwise to 3.6 g (11.8 mmol) of the Sem-protected triazole in 200 ml of anhydrous THF at -78 °C. The reaction was allowed to stir for 1 h prior to the addition of 1.7 ml (14.2 mmol) of


benzyl bromide. After stirring at -78 °C for 2 h, the reaction was allowed to warm to ambient temperature and stir overnight. The solvent was removed in vacuo; the residue dissolved in methylene chloride, washed with water, and dried (MgSO₄). Silica gel chromatography (Waters Prep 500A) using ethyl acetate/hexane (5:95) gave 600 mg (10%) of the Sem protected 5-butyl-3-(1,1-difluoro-2-phenylethyl)-1H-1,2,4-triazole: NMR (CDCl₃) δ -0.04 (s, 9H), 0.85 (t, J = 8Hz, 2H), 0.94 (t, J = 8 Hz, 3H), 1.30-1.44 (m, 2H), 1.67-1.79 (m, 2H), 2.73 (t, J = 8 Hz, 2H), 3.54 (t, J = 8 Hz, 2H), 3.70 (t, J = 16 Hz, 2H), 5.34 (s, 2H), 7.21-7.31 (m, 5H). The Sem protected triazole was dissolved in 5 ml of ethanol and 5 ml of 3M HCl and allowed to stir at reflux for 3 h. The ethanol was removed in vacuo and the pH adjusted to nine with dilute sodium hydroxide. The resulting solution was extracted with methylene chloride; the extracts were combined, dried (MgSO₄), and concentrated in vacuo to provide 5-butyl-3-(1,1-difluoro-2-phenethyl-1H-1,2,4-triazole. Following General Procedure A, 1.4 mmol of this material was coupled with 1.4 mmol of the alkylating reagent prepared in step 1 of Example 3. The crude product was dissolved in 10 mL of acetic acid/water (9:1) and stirred at ambient temperature for 3 days. The solvent was removed in vacuo; the residue dissolved in dilute base and washed with toluene. The water was acidified to pH 3-4, extracted with ethyl acetate, and dried (MgSO₄). Purification by reverse phase chromatography (Waters Delta Prep 3000) using acetonitrile/water (47:53) (0.05% TFA) gave the TFA salt. The salt was dissolved in dilute base, acidified to pH 3-4, extracted with ethyl acetate, and dried (MgSO₄). Lyophilization from acetonitrile/water gave 290 mg (42%) of 5-[4'-[[5-butyl-3-(1,1-difluoro-2-phenylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-1-yl]-1H-tetrazole: NMR (CDCl₃) δ 0.86 (t, J = 8 Hz, 3H), 1.21-1.36 (m, 2H), 1.54-1.68 (m, 2H), 2.64 (t, J = 8 Hz, 2H), 3.57 (t, J = 16 Hz, 2H), 5.23 (s, 2H), 6.92 (d, J = 8 Hz, 2H), 7.10 (d, J =

225

8 Hz, 2H), 7.16-7.25 (m, 5H), 7.39 (dd, $J = 8$ and 2 Hz, 1H), 7.50-7.65 (m, 2H), 7.99 (dd, $J = 8$ and 2 Hz, 1H); MS (FAB) m/e (rel intensity) 500(100), 266 (6), 235 (8), 207 (25), 192 (8); HRMS. Calc'd for $M+H$: 500.2374. Found:

5 500.2358.

Example 52




5

5-butyl-1-[2'-(1H-tetrazol-5-yl)][1,1'-biphenyl]-4-ylmethyl-1H-1,2,4-triazole-3-carboxylic acid

A 3.36 g (7.1 mmol) sample of 5-[4'-[[5-butyl-3-(2,2-diethoxyethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole from Example 38 was dissolved in 15 ml of ethanol and 15 ml of 3M HCl. The reaction was stirred at ambient temperature overnight and then at reflux for 2 h. The solvent was removed in vacuo, the residue dissolved in 15 ml of THF and 15 ml of 3M HCl and stirred for 3 days at ambient temperature. The solvent was removed in vacuo; the residue dissolved in dilute base, acidified and extracted with ethyl acetate and methylene chloride. The solvents were combined and dried (MgSO₄). The solvents were removed in vacuo. The residue was dissolved in 70 ml of acetone and 35 ml of water. Potassium permanganate was added in 4 equal portions of 190 mg after each preceding portion had reacted. The mixture was filtered through celite and the solvent removed in vacuo. Purification by reverse phase chromatography (Waters Delta Prep 3000) gave 88 mg (7%) of 5-butyl-1-[2'-(1H-tetrazol-5-yl)][1,1'-biphenyl]-4-ylmethyl-1H-1,2,4-triazole-3-carboxylic acid

as the colorless TFA salt after lyophilization from
acetonitrile/water: NMR (DMSO-d₆) δ 0.85 (t, J = 8 Hz,
3H), 1.23-1.38 (m, 2H), 1.52-1.62 (m, 2H), 2.78 (t, J = 8
Hz, 2H), 5.46 (s, 2H), 7.13 (q, J = 8 Hz, 4H), 7.51-7.62
5 (m, 2H), 7.64-7.73 (m, 2H); MS (FAB) m/e (rel intensity)
404 (100), 235 (20), 207 (91), 192 (24); HRMS. Calc'd for
M+H: 404.1835. Found: 404.1882.

Example 53



5

5-[4'-[(5-butyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-
biphenyl]-2-yl]-1H-tetrazole

10

Following general procedure A, 630 mg (5 mmol) of 3-butyl-1H-1,2,4-triazole was reacted with 5 mmol of the alkylating reagent prepared in step 1 of Example 3 to give 2.3 g (76%) of a mixture of the two isomers. Deprotection of the faster isomer provided 5-[4'-[(5-butyl-1H-1,2,4-


15

triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]1H-tetrazole: NMR (CDCl₃) δ 0.87 (t, J = 8 Hz, 3H), 1.35-1.41 (m, 2H), 1.59-1.72 (m, 2H), 2.72 (t, J = 8 Hz, 2H), 5.30 (s, 2H), 7.09 (d, J = 8 Hz, 2H), 7.17 (d, J = 8 Hz, 2H), 7.44 (dd, J = 8 and 2 Hz, 1H), 7.52-7.66 (m, 2H), 7.72 (s, 1H), 8.01 (dd, J

20

= 8 and 2 Hz, 1H); MS (FAB) m/e (rel intensity) 360 (88), 235 (20), 207 (100), 192 (31), 178 (22); HRMS. Calc'd for M+H: 360.1936. Found: 360.1938.

Example 54



5


5-[4'-[(3-butyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-
biphenyl]-2-yl]-1H-tetrazole

10

The slower moving isomer from Example 53 was deprotected to provide 5-[4'-[(3-butyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole: NMR (CDCl₃) δ 0.74 (t, J = 8 Hz, 3H), 1.19-1.33 (m, 2H), 1.51-1.63 (m, 2H), 2.55 (t, J = 8 Hz, 2H), 5.23 (s, 2H), 7.10 (q, J = 8 Hz, 4H), 7.44 (dd, J = 8 and 2 Hz, 1H), 7.50-7.64 (m, 2H), 7.92 (dd, J = 8 and 2 Hz, 1H), 8.02 (s, 1H); MS (FAB) m/e (rel intensity) 360 (84), 135 (18), 207 (100), 192 (26), 178 (16), 126 (22); HRMS. Calc'd for M+H: 360.1936. Found: 360.1971.

15

Example 55



5

5-[4'-[[3-butyl-5-(tricyclo[3.3.1.3.7]dec-1-yl)-1H-1,2,4-
triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

10

Following general procedure A, 1.3 g (5.0 mmol) of 5-butyl-3-adamantyl-1H-1,2,4-triazole was reacted with 5.0 mmol of the alkylating reagent prepared in step 1 of Example 3 to give 2.94 g (72%) of a mixture of the two isomers. Deprotection of the faster moving isomer provided


15

5-[4'-[[3-butyl-5-(tricyclo[3.3.1.3.7]dec-1-yl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole:

20

NMR (CDCl₃) δ 0.90 (t, J = 8 Hz, 3H), 1.27-1.41 (m, 2H), 1.42-1.55 (m, 3H), 1.58-1.73 (m, 5H), 1.76-1.91 (m, 9H), 2.28-2.43 (m, 2H), 5.42 (s, 2H), 6.72-6.83 (m, 2H), 7.07 (d, J = 8 Hz, 2H), 7.45-7.66 (m, 3H), 7.88 (t, J = 8 Hz, 1H); MS (FAB) m/e (rel intensity) 494 (100), 260 (37), 207 (74), 178 (22); HRMS. Calc'd for M+H: 494.3032. Found: 494.3014.

Example 56



5

5-[4'-[[5-butyl-3-(tricyclo[3.3.1.1^{3.7}]dec-1-yl)-1H-1,2,4-
triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole


10

The slower moving isomer from Example 55 was deprotected to provide 5-[4'-[[5-butyl-3-(tricyclo[3.3.1.1^{3.7}]dec-1-yl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole: NMR (CDCl₃) δ 0.87 (t, J = 8 Hz, 3H), 1.31-1.47 (m, 2H), 1.52-1.71 (m, 9H), 1.78 (s, 7H), 1.88 (s, 3H), 2.20-2.35 (m, 2H), 5.22 (s, 2H), 6.87 (d, J = 8 Hz, 2H), 7.13 (d, J = 8 Hz, 2H), 7.42 (dd, J = 8 and 2 Hz, 1H), 7.49-7.63 (m, 2H), 7.93 (dd, J = 8 and 2 Hz, 1H); MS (FAB) m/e (rel intensity) 494 (73), 451 (10), 260 (24), 207 (66), 178 (21); HRMS. Calc'd for

20

M+H: 494.3032. Found: 494.3026.

Example 57



5

5-[4'-[[3-butyl-5-(3-cyclohexylpropyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

10

Following general procedure A, 1.25 g (5 mmol) of 5-butyl-3-(3-cyclohexylpropyl)-1H-1,2,4-triazole was reacted with 5 mmol of the alkylating reagent prepared in step 1 of Example 3 to give 3.1 g (85%) of a mixture of the two isomers. Deprotection of the faster moving isomer provided 5-[4'-[[3-butyl-5-(3-cyclohexylpropyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole:


15

NMR (CDCl₃) δ 0.73-0.92 (m, 2H), 0.87 (t, J = 8 Hz, 3H), 1.06-1.21 (m, 6H), 1.22-1.36 (m, 2H), 1.53-1.71 (m, 9H), 2.51 (t, J = 8 Hz, 2H), 2.61 (t, J = 8 Hz, 2H), 5.22 (s, 2H), 7.01 (d, J = 8 Hz, 2H), 7.12 (d, J = 8 Hz, 2H), 7.42 (dd, J = 8 and 2 Hz, 1H), 7.51-7.64 (m, 2H), 7.94 (dd, J = 8 and 2 Hz, 1H); MS (FAB) m/e (rel intensity) 484 (83), 250 (38), 207 (100), 178 (25); HRMS. Calc'd for M+H:

20

484.3189. Found: 484.3223.

Example 58



5


5-[4'-[[5-butyl-3-(3-cyclohexylpropyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

10 The slower moving isomer from Example 57 was deprotected to provide 5-[4'-[[5-butyl-3-(3-cyclohexylpropyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole: NMR (CDCl₃) δ 0.74-0.90 (m, 2H), 0.83 (t, J = 8 Hz, 3H), 1.05-1.22 (m, 6H), 1.23-1.35 (m, 2H), 1.50-1.70 (m, 9H), 2.40 (t, J = 8 Hz, 2H), 2.57 (t, J = 8 Hz, 2H), 5.20 (s, 2H), 6.96 (d, J = 8 Hz, 2H), 7.10 (d, J = 8 Hz, 2H), 7.43 (dd, J = 8 and 2 Hz, 2H), 7.50-7.64 (m, 2H), 7.92 (dd, J = 8 and 2 Hz, 2H); MS (FAB) m/e (rel intensity) 484 (65), 250 (26), 207 (100), 178 (18); HRMS. Calc'd for M+H: 484.3189. Found: 484.3181.

15

20

Example 59



5


5-[4'-[[5-butyl-3-(fluoromethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

10 Following general procedure A, 790 mg (5.0 mmol) of 5-butyl-3-(fluoromethyl)-1H-1,2,4-triazole was reacted with 5 mmol of the alkylating reagent prepared in step 1 of Example 3 to give 1.3 g (42%) of a mixture of the two isomers. Deprotection of the faster moving isomer provided

15 5-[4'-[[5-butyl-3-(fluoromethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole: NMR (CDCl₃) δ 0.86 (t, J = 8 Hz, 3H), 1.26-1.40 (m, 2H), 1.56-1.69 (m, 2H), 2.67 (t, J = 8 Hz, 2H), 5.16 (s, 1H), 5.27 (s, 2H), 5.32 (s, 1H), 7.06 (d, J = 8 Hz, 2H), 7.16 (d, J = 8 Hz, 2H), 7.43 (dd, J = 8 and 2 Hz, 1H), 7.52-7.65 (m, 2H), 7.99 (dd, J = 8 and 2 Hz, 1H); MS (FAB) m/e (rel intensity) 392 (100), 235 (22), 207 (66), 192 (18), 178 (14); HRMS. Calc'd for M+H: 392.1999. Found: 392.1932.

20

Example 60




5

5-[4'-[[[3-butyl-5-(fluoromethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

The slower moving isomer from Example 59 was
10 deprotected to provide 5-[4'-[[[3-butyl-5-(fluoromethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole: NMR (CDCl₃) δ 0.89 (t, J = 8 Hz, 3H), 1.24-1.40 (m, 2H), 1.59 (m, 2H), 2.65 (t, J = 8 Hz, 2H), 5.38 (s, 3H), 5.55 (s, 1H), 7.21 (s, 4H), 7.43 (d, J = 8 Hz, 1H),
15 7.51-7.65 (m, 2H), 8.05 (d, J = 8 Hz, 1H); MS (FAB) m/e (rel intensity) 392 (90), 235 (56), 207 (100), 192 (51), 178 (24); HRMS. Calc'd for M+H: 392.1999. Found: 392.1944.

Example 61



5


5-[4'-[[[3-butyl-5-(tricyclo[3.3.1.1]^{3.7}]dec-1-ylmethyl)-1H-1,2,4-triazole-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

10

Following general procedure A, 1.4 g (5.0 mmol) of 5-butyl-3-adamantyl-1H-1,2,4-triazole was reacted with 5.0 mmol of the alkylating reagent prepared in step 1 of Example 3 to give a mixture of the two isomers.

15 Deprotection of the faster moving isomer provided 5-[4'-[[[3-butyl-5-(tricyclo[3.3.1.1]^{3.7}]dec-1-ylmethyl)-1H-1,2,4-triazole-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole:
 NMR (CDCl₃) δ 0.84 (t, J = 8 Hz, 3H), 1.16-1.30 (m, 2H), 1.41 (s, 6H), 1.44-1.56 (m, 5H), 1.65 (d, J = 12 Hz, 3H),
 20 1.92 (s, 3H), 2.18 (s, 2H), 2.30 (t, J = 8 Hz, 2H), 5.18 (s, 2H), 6.79 (d, J = 8 Hz, 2H), 7.01 (d, J = 8 Hz, 2H), 7.46 (d, J = 8 Hz, 1H), 7.52 (m, 2H), 7.85 (dd, J = 8 and 2 Hz, 1H); MS (FAB) m/e (rel intensity) 508 (45), 274 (25), 207 (100), 192 (31); HRMS. Calc'd for M+H: 508.3189.
 25 Found: 508.3162.

Example 62



5


5-[4'-[[5-butyl-3-(tricyclo[3.3.1.1^{3,7}]dec-1-yl)methyl]-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

10

The slower moving isomer from Example 61 was deprotected to provide 5-[4'-[[5-butyl-3-(tricyclo[3.3.1.1^{3,7}]dec-1-yl)methyl]-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole: NMR (CDCl₃)

15 δ 0.85 (t, J = 8 Hz, 3H), 1.21-1.37 (m, 8H), 1.44-1.58 (m, 5H), 1.62 (d, J = 10 Hz, 3H), 1.86 (s, 3H), 2.07 (s, 2H), 2.49 (t, J = 8 Hz, 2H), 5.21 (s, 2H), 6.89 (d, J = 8 Hz, 2H), 7.07 (d, J = 8 Hz, 2H), 7.53 (dd, J = 8 and 2 Hz, 1H), 7.52-7.65 (m, 2H), 7.91 (dd, J = 8 and 2 Hz, 1H), MS (FAB)
20 m/e (rel intensity) 508 (42), 274 (22), 207 (100), 192 (34); HRMS. Calc'd for M+H: 508.3189. Found: 508.3142.

Example 63



5

5-butyl-1-[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-ylmethyl]-1H-1,2,4-triazole-3-propanoic acid

10


A 2.18 g sample of the mixture of the two isomers from Example 45 was oxidized to the corresponding aldehydes with oxalyl chloride/dimethyl sulfoxide and subsequently oxidized to the acids with potassium permanganate to give 1.8 g of a mixture of the two propanoic isomers.

15

Deprotection of the faster moving isomer provided 5-butyl-1-[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-ylmethyl]-1H-1,2,4-triazole-3-propanoic acid: NMR (CDCl₃) δ 0.99 (t, J = 8 Hz, 3H), 1.28-1.43 (m, 2H), 1.61-1.74 (m, 2H), 2.67-2.80 (m, 4H), 2.94 (t, J = 6 Hz, 2H), 5.18 (s, 2H), 7.04-7.14 (m, 4H), 7.42 (dd, J = 8 and 2 Hz, 1H), 7.50 (dt, J = 8 and 2 Hz, 1H), 7.59 (dt, J = 8 and 2 Hz, 1H), 7.90 (dd, J = 8 and 2 Hz, 1H); MS (FAB) m/e (rel intensity) 432 (89), 389 (15), 237 (36), 198 (100); HRMS. Calc'd for M+H: 432.2148. Found: 432.2216.

20

Example 64



5


3-butyl-1-[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-ylmethyl]-1H-1,2,4-triazole-5-propanoic acid

10

The slower moving isomer from Example 63 was deprotected to give 3-butyl-1-[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-ylmethyl]-1H-1,2,4-triazole-5-propanoic acid:

15 NMR (CDCl₃) δ 0.92 (t, J = 8 Hz, 3H), 1.30-1.40 (m, 2H), 1.65-1.78 (m, 2H), 2.70 (t, J = 8 Hz, 2H), 2.84-2.91 (m, 2H), 2.93-3.00 (m, 2H), 5.40 (s, 2H), 7.20 (d, J = 8 Hz, 2H), 7.34 (d, J = 8 Hz, 2H), 7.37-7.42 (m, 1H), 7.51-7.62 (m, 2H), 8.30-8.36 (m, 1H); MS (FAB) m/e (rel intensity)
20 432 (100) 198 (45); HRMS. Calc'd for M+H: 432.2148.
Found: 432.2214.

Example 65




5

5-[4'-[(3-butyl-5-cyclohexyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

Following General Procedure A, 2.07 g (10 mmol) of 3-butyl-5-cyclohexyl-1H-1,2,4-triazole was reacted with 6.5 g (11.8 mmol) of the alkylating reagent prepared in step 1 of Example 3 to give 6.3 g (9.2 mmol) of a mixture of the two isomers. Deprotection of the faster moving isomer provided 33 mg of 5-[4'-[(3-butyl-5-cyclohexyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-1,2,4-triazole as a colorless solid: mp 157.5-159.5 °C; NMR (CDCl₃) δ 0.87 (t, J = 7 Hz, 3H), 1.11-1.36 (m, 6H), 1.45-1.70 (m, 6H), 1.70-1.80 (m, 2H), 2.50 (t, J = 7 Hz, 2H), 2.64-2.78 (m, 1H), 5.23 (s, 2H), 6.98 (d, J = 8 Hz, 2H), 7.08 (d, J = 8 Hz, 2H), 7.41 (dd, J = 5 and 1 Hz, 1H), 7.49 (dt, J = 5 and 1 Hz, 1H), 7.57 (dt, J = 5 and 1 Hz, 1H), 7.80 (dd, J = 5 and 1 Hz, 1H): MS (FAB) m/e (rel intensity), 442 (100), 414 (10), 399 (20), 235 (8), 207 (65), 192 (18).

Example 66




5

5-[4'-[(5-butyl-3-cyclohexyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

The slower moving isomer from Example 65 was
 10 deprotected to provide 135 mg of 5-[4'-[(5-butyl-3-
 cyclohexyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-
 yl]-1H-tetrazole as a colorless solid: mp 125-127 °C; NMR
 (CDCl₃) δ 0.85 (t, J = 7 Hz, 3H), 1.12-1.40 (m, J = 8 Hz,
 8H), 1.52-1.75 (m, J = 8 Hz, 4H), 1.81 (d, J = 8 Hz, 2H),
 15 2.50 (t, J = 8 Hz, 3H), 5.20 (s, 2H), 6.94 (d, J = 8 Hz,
 2H), 7.09 (d, J = 8 Hz, 2H), 7.42 (dd, J = 8 and 2 Hz, 1H),
 7.50-7.64 (m, J = 8 Hz, 2H), 7.88 (dd, J = 8 and 2 Hz, 1H);
 MS (FAB) m/e (rel intensity) 442 (100), 414 (10), 399 (20),
 235 (10), 207 (75), 192 (17).

Example 67




5

1,1-dimethylethyl 3-butyl-[1-[2'-(1H-tetrazol-5-yl)[1,1'-
biphenyl]-4-ylmethyl]-1H-1,2,4-triazol-5-yl]propylcarbamate

Following General Procedure A, 6.0 g (20 mmol)
 10 of 3-butyl-5-(N-Boc-3-aminopropyl)-1H-1,2,4-triazole was
 reacted with 13.1 g (24 mmol) of the alkylating reagent
 prepared in step 1 of Example 3. Deprotection of the
 faster moving isomer gave 3.52 g (33%) of 1,1-dimethylethyl
 3-butyl-[1-[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-
 15 ylmethyl]-1H-1,2,4-triazol-5-yl]propylcarbamate as a
 colorless solid: NMR (DMSO-d₆) δ 0.88 (t, J = 8 Hz, 3H),
 1.22-1.34 (m, J = 8 Hz, 2H), 1.36 (s, 9H), 1.54-1.66 (m, J
 = 7 Hz, 2H), 1.66-1.78 (m, J = 7 Hz, 2H), 2.5 (t, J = 8 Hz,
 2H), 2.67 (t, J = 8 Hz, 2H), 2.96 (q, J = 8 Hz, 2H), 5.26
 20 (s, 2H), 7.08 (s, 4H), 7.55 (q, J = 8 Hz, 2H), 7.65 (d, J =
 8 Hz, 2H); MS (FAB) m/e (rel intensity) 517 (60), 489 (10),
 461 (10), 439 (10), 417 (23), 389 (18), 357 (8), 323 (8),
 305 (15), 283 (80), 227 (70), 207 (100), 183 (65); HRMS.
 Calc'd for M+H: 517.3039. Found 517.3001.

Example 68



5

1,1-dimethylethyl 5-butyl-[1-[2'-(1H-tetrazol-5-yl)][1,1'-biphenyl]-4-ylmethyl]-1H-1,2,4-triazol-3-ylpropylcarbamate

10

The slower moving isomer from Example 67 was deprotected to give 3.43 g (33%) of 1,1-dimethylethyl 5-butyl-[1-[2'-(1H-tetrazol-5-yl)][1,1'-biphenyl]-4-ylmethyl]-1H-1,2,4-triazol-3-ylpropylcarbamate as a colorless solid:


15

NMR (DMSO- d_6) δ 0.84 (t, J = 7 Hz, 3H), 1.29 (q, J = 8 Hz, 2H), 1.38 (s, 9H), 1.48-1.59 (m, J = 8 Hz, 2H), 1.72 (t, J = 8 Hz, 2H), 2.53 (t, J = 7 Hz, 2H), 2.67 (t, J = 8 Hz, 2H), 2.97 (q, J = 8 Hz, 2H), 5.28 (s, 2H), 7.08 (s, 4H), 7.50 (d, J = 8 Hz, 1H) 7.55 (d, J = 8 Hz, 1H), 7.61-7.68 (m, J = 7 Hz, 2H); MS (FAB) m/e (rel intensity) 517 (40),

20

439 (10) 417 (40), 390 (20), 357 (10), 207 (100), 192 (40); HRMS. Calc'd for $M+H$: 517.3039. Found: 517.3014.

Example 69




5

5-[4'-[[3-(1,1,2,2,2-pentafluoroethyl)-5-butyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazol

Following General Procedure A, 1.21 g (5.0 mmol) of 3-perfluoroethyl-5-butyl-1H-1,2,4-triazole was reacted with 3.2 g (5.7 mmol) of the alkylating reagent prepared in step 1 of Example 3. Deprotection provided 409 mg of 5-[4'-[[3-(1,1,2,2,2-pentafluoroethyl)-5-butyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole as a colorless oil: NMR (CDCl₃) δ 0.90 (t, J = 7 Hz, 3H), 1.36 (q, J = 7 Hz, 2H), 1.67 (t, J = 7 Hz, 2H), 2.78 (t, J = 7 Hz, 2H), 5.38 (s, 2H), 7.18 (s, 4H), 7.42 (d, J = 8 Hz, 1H), 7.51-7.67 (m, J = 8 Hz, 2H), 7.98 (d, J = 8 Hz, 1H); MS (FAB) m/e (rel intensity) 478 (40), 452 (15), 235 (90), 207 (100), 178 (40), 152 (35); HRMS. Calc'd. for M+H: 478.1778. Found: 478.1807.

Example 70



5


5-[4'-[[[3-(1,1,2,2,3,3,3-heptafluoropropyl)-5-butyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

10 Following General Procedure A, 1.46 g (5.0 mmol) of 5-butyl-3-perfluoropropyl-1H-1,2,4-triazole was reacted with 3.20 g (5.7 mmol) of the alkylating reagent prepared in step 1 of Example 3. Deprotection provided 210 mg of 5-[4'-[[[3-(1,1,2,2,3,3,3-heptafluoropropyl)-5-butyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole as a colorless oil: NMR (CDCl₃) δ 0.90 (t, J = 7 Hz, 3H), 1.36 (q, J = 8 Hz, 2H), 1.63-1.75 (m, J = 8 Hz, 2H), 2.78 (t, J = 8 Hz, 2H), 5.40 (s, 2H), 7.20 (q, J = 7 Hz, 4H), 7.42 (dd, J = 8 and 2 Hz, 1H), 7.52-7.65 (m, J = 8 Hz, 2H), 8.04 (dd, J = 8 and 2 Hz, 1H); MS (FAB) m/e (rel intensity) 528 (35), 502 (15), 294 (12), 235 (90), 207 (100), 136 (15); HRMS. Calc'd for M+H: 528.1747. Found: 528.1701.

15

20

Example 71




5

5-[4'-[[3-butyl-5-(3-aminopropyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

Under a static nitrogen atmosphere 2.40 g (4.7 mmol) of 5-[4'-[[3-butyl-5-(N-Boc-3-aminopropyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole from Example 67 was dissolved in 30 mL of 4N HCl in dioxane at ambient temperature and allowed to stir overnight. The solvents were removed in vacuo; the residue was triturated with diethyl ether and filtered providing 2.08 g (98%) of 5-[4'-[[3-butyl-5-(3-aminopropyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole as the hydrochloride salt: NMR (DMSO- d_6) δ 0.88 (t, J = 7 Hz, 3H); 1.25-1.39 (m, 2H), 1.57-1.69 (m, 2H), 1.92-2.40 (m, 2H), 2.62 (t, J = 7 Hz, 2H), 2.89 (q, J = 8 Hz, 2H), 2.98 (t, J = 8 Hz, 2H), 5.38 (s, 2H), 7.10 (d, J = 8 Hz, 2H), 7.19 (d, J = 8 Hz, 2H), 7.54 (d, J = 8 Hz, 1H), 7.59 (dd, J = 8 and 2 Hz, 1H), 7.66 (s, 1H), 7.70 (dd, J = 8 and 2 Hz, 1H), 8.10 (br s, 2H); MS (FAB) m/e (rel intensity) 417 (38), 389 (10), 357 (10), 207 (100), 166 (50), 139 (10) 115 (10); HRMS. Calc'd for $M+H$: 417.2515. Found: 417.2563.

Example 72




5

5-[4'-[[3-(3-aminopropyl)-5-butyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

Under a static nitrogen atmosphere 2.30 g (4.5 mmol) of 5-[4'-[[3-(N-Boc-3-aminopropyl)-5-butyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole from Example 68 was dissolved in 30 mL of 4N HCl in dioxane at ambient temperature and allowed to stir overnight. The solvents were removed in vacuo; the residue was triturated with diethyl ether and filtered providing 1.99 g (98%) of 5-[4'-[[3-(3-aminopropyl)-5-butyl-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole as the hydrochloride salt: NMR (DMSO- d_6) δ 0.85 (t, J = 7 Hz, 3H), 1.22-1.38 (m, 2H), 1.59 (t, J = 8 Hz, 2H), 1.96 (t, J = 8 Hz, 2H), 2.71-2.79 (m, 2H) 2.80-2.92 (m, 4H), 5.40 (s, 2H) 7.10 (d, J = 8 Hz, 2H), 7.18 (d, J = 8 Hz, 2H) 7.52 (d, J = 8 Hz, 1H), 7.59 (d, J = 8 Hz, 1H), 7.68 (d, J = 8 Hz, 2H); MS (FAB) m/e (rel intensity) 417 (60), 389 (10), 357 (10), 263 (5), 235 (10), 207 (100), 166 (40), 136 (15); HRMS. Calc'd for M+H: 417.2515. Found: 417.2510.

Example 73




5

5-[4'-[(3-butyl-5-cyclohexylmethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

Following General Procedure A, 2.21 g (10.0 mmol) of 3-butyl-5-cyclohexylmethyl-1H-1,2,4-triazole was reacted with 6.55 g (11.8 mmol) of the alkylating reagent prepared in step 1 of Example 3 to give 5.90 g (85%) of a mixture of the two isomers. The faster moving isomer was deprotected to provide 64 mg of 5-[4'-[(3-butyl-5-cyclohexylmethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole as a colorless solid: NMR (CDCl₃) δ 0.92 (t, J = 7 Hz, 3H) 0.94-1.05 (m, 2H) 1.08-1.25 (m, 3H), 1.25-1.41 (m, 2H), 1.55-1.80 (m, 8H), 2.65 (d, J = 7 Hz, 2H), 2.68 (t, J = 7 Hz, 2H), 5.27 (s, 2H), 7.11 (q, J = 8 Hz, 4H), 7.42 (dd, J = 8 and 2 Hz, 1H), 7.49-7.63 (m, 2H), 7.94 (dd, J = 8 and 2 Hz, 1H); MS (FAB) m/e (rel intensity) 456 (100), 428 (10), 413 (20), 280 (10), 235 (10), 222 (25), 207 (80), 192 (30); HRMS. Calc'd for M+H: 456.2876. Found: 456.2839.

Example 74




5

5-[4'-[(3-cyclohexylmethyl-5-butyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

The slower moving isomer from Example 73 was
 10 deprotected to provide 51 mg of 5-[4'-[(3-cyclohexylmethyl-5-butyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole as a colorless solid: NMR (CDCl₃) δ 0.84 (t, J = 8 Hz, 3H), 0.92 (q, J = 8 Hz, 2H), 1.05-1.22 (m, 3H), 1.22-1.36 (m, 2H), 1.52-1.69 (m, 8H), 2.50 (d, J = 7 Hz,
 15 2H), 2.63 (t, J = 8 Hz, 2H), 5.18 (s, 2H), 7.05 (q, J = 8 Hz, 4H), 7.35-7.55 (m, 3H), 7.68 (dd, J = 8 and 2 Hz, 1H); MS (FAB) m/e (rel intensity) 456 (90), 413 (10), 281 (10), 243 (10), 222 (35), 207 (100), 192 (35); HRMS. Calc'd for M+H: 456.2876. Found: 456.2849.

Example 75




5

4'-[(3-ethyl-5-butyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid

Following General Procedure A, 1.0 g (6.0 mmol) of 3-ethyl-5-butyl-1H-1,2,4-triazole was reacted with 2.41 g (7.9 mmol) of the alkylating reagent prepared in step 1 of Example 1. Hydrolysis of the faster moving isomer provided 220 mg of 4'-[(3-ethyl-5-butyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid as a colorless solid: NMR (DMSO-d₆) δ 0.85 (t, J = 7 Hz, 3H), 1.19 (t, J = 7 Hz, 3H), 1.27-1.38 (m, 2H), 1.52-1.63 (m, 2H), 2.59 (q, J = 7 Hz, 2H), 2.74 (t, J = 7 Hz, 2H), 5.35 (s, 2H), 7.21 (d, J = 8 Hz, 2H), 7.31 (d, J = 8 Hz, 2H), 7.35 (dd, J = 8 and 2 Hz, 1H), 7.48 (dt, J = 8 and 2 Hz, 1H), 7.57 (dt, J = 8 and 2 Hz, 1H), 7.72 (dd, J = 8 and 2 Hz, 1H).

Example 76




5

4'-[(3-butyl-5-ethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid

- The slower moving isomer from Example 75 was
- 10 hydrolyzed to provide 174 mg of 4'-[(3-butyl-5-ethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid as a colorless solid: NMR (DMSO-d₆) δ 0.89 (t, J = 7 Hz, 3H), 1.17 (t, J = 7 Hz, 3H), 1.27-1.40 (m, 2H), 1.56-1.68 (m, 2H), 2.58 (t, J = 8 Hz, 2H), 2.77 (q, J = 8 Hz, 2H),
- 15 5.35 s, 2H), 7.21 (d, J = 8 Hz, 2H), 7.32 (d, J = 8 Hz, 2H), 7.36 (dd, J = 8 and 2 Hz, 1H), 7.46 (dt, J = 8 and 2 Hz, 1H), 7.57 (dt, J = 8 and 2 Hz, 1H), 7.73 (dd, J = 8 and 2 Hz, 1H).

Example 77




5

4'-[(3-butyl-5-methyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid

Following General Procedure A, 1.0 g (7.2 mmol) of 3-butyl-5-methyl-1H-1,2,4-triazole was reacted with 2.89 g (9.5 mmol) of the alkylating reagent prepared in step 1 of Example 1. Hydrolysis of the slower moving isomer provided 220 mg of 4'-[(3-butyl-5-methyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid as a colorless solid: NMR (DMSO-d₆) δ 0.88 (t, J = 7 Hz, 3H), 1.25-1.38 (m, 2H), 1.54-1.67 (m, 2H), 2.39 (s, 3H), 2.54 (t, J = 8 Hz, 2H), 5.32 (s, 2H), 7.21 (d, J = 8 Hz, 2H), 7.32 (d, J = 8 Hz, 2H), 7.36 (dd, J = 8 and 2 Hz, 1H), 7.46 (dt, J = 8 and 2 Hz, 1H), 7.56 (dt, J = 8 and 2 Hz, 1H), 7.72 (dd, J = 8 and 2 Hz, 1H).

Example 78



5 4'-[(3-methyl-5-butyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-
biphenyl]-2-carboxylic acid


The faster moving isomer from Example 77 was hydrolyzed to provide 19 mg of 4'-[(3-methyl-5-butyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid as a colorless solid: NMR (DMSO-d₆) δ 0.86 (t, J = 7 Hz, 3H), 1.25-1.38 (m, 2H), 1.52-1.64 (m, 2H), 2.21 (s, 3H), 2.72 (t, J = 7 Hz, 2H), 5.32 (s, 2H), 7.21 (d, J = 8 Hz, 2H), 7.32 (d, J = 8 Hz, 2H), 7.35 (dd, J = 8 and 2 Hz, 1H), 7.45 (dt, J = 8 and 2 Hz, 1H), 7.57 (dt, J = 8 and 2 Hz, 1H), 7.72 (dd, J = 8 and 2 Hz, 1H).

A class of highly preferred specific conjugates of the invention is provided by conjugates formed from a biphenylmethyl 1H-substituted-1,2,4-triazole AII antagonist compound linked to a cleavable glutamyl residue. Each conjugate of this class contains a diamino linker moiety which connects a terminal carboxylic acid moiety on the biphenylmethyl portion of the AII antagonist compound with a terminal carboxylic acid moiety on the gamma carbon of the cleavable glutamyl residue. Example #79 is a detailed description of a conjugate of this class. Other specific conjugates of Examples #80-#144, as shown in Table IV, may

be prepared generally in accordance with the procedures of Example #79.

5

Example 79



10


N-acetylglutamic acid, 5-[[[4'-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]]carbonylhydrazide

To a solution of 10.45 g (34.5 mmol) of N-Boc-L-glutamic acid- α -tertbutyl ester (BACHEM) in 100 mL of methylene chloride under nitrogen was added 3.5 g (17.0 mmol) of solid dicyclohexylcarbodiimide (DCC). The reaction was allowed to stir for 2 h and filtered under nitrogen. The anhydride solution was then added to a solution of 6.03 g (14.9 mmol) of the compound of Example 4 in 75 mL of methylene chloride under nitrogen. The reaction was stirred overnight and concentrated in vacuo. Purification by silica gel chromatography (Waters Prep-500A) using ethyl acetate gave 7.90 g (77%) of pure material by thin-layer chromatography (TLC). This material was redissolved in 100 mL of methylene chloride under nitrogen and cooled to 0°C prior to the addition of 135 mL

of TFA. The stirred reaction was allowed to warm to ambient temperature overnight and concentrated in vacuo.









The crude product was dissolved in 80 mL of
5 acetonitrile/water (1:1) and the pH adjusted to 9 with 1 M
K₂CO₃. The solution was cooled to 0°C and 1.1 mL (11 mmol)
of acetic anhydride and 11 mL (11 mmol) of 1 M K₂CO₃ was
added every 30 min for 5 h; during the course of this
reaction the pH was maintained at 9 and the reaction
10 temperature kept below 5°C. After the last addition, the
reaction was allowed to warm to ambient temperature
overnight. The pH was adjusted to 4 with 3 M HCl and the
reaction concentrated to 300 mL. Purification by reverse
phase chromatography (Waters Delta prep-3000) using
15 isocratic 32% acetonitrile/water (0.05% TFA) gave 6.11 g
(55%-overall yield from the compound of Example 4) of
colorless product: NMR (DMSO-d₆) δ 0.85 (t, J = 8Hz, 3H),
0.88 (t, J = 8Hz, 3H), 1.23-1.39 (m, 4H), 1.53-1.68 (m,
6H), 1.84 (s, 3H), 2.15-2.24 (m, 2H), 2.55 (t, J = 8Hz,
20 2H), 2.70 (t, J = 8Hz, 2H), 4.10-4.20 (m, 1H), 5.31 (s,
2H), 7.11-7.19 (m, 2H), 7.37-7.57 (m, 6H); MS (FAB) m/e
(rel intensity) 577 (42), 389 (10), 195 (82), 182 (100),
167 (63), 152 (28); HRMS. Calcd. for M+H: 577.3138.
Found: 577.3160. Anal. Calcd. for C₃₁H₄₀N₆O₅·1.5 CF₃CO₂H:
25 C, 54.65; H, 5.56; N, 11.25; F, 11.50. Found: C, 54.42;
H, 5.75, N, 11.35, F, 11.85.



256

TABLE IV



Ex. #	R^1	R^2	L	B	E	P
80	C_4H_9 (n)	C_4H_9 (n)	-NH-	H	H	H
81	C_4H_9 (n)	C_4H_9 (n)	-NH-	H	CH_3	H
82	C_4H_9 (n)	C_4H_9 (n)	-NH-	H	CH_3	$COCH_3$
83	C_4H_9 (n)	C_4H_9 (n)	-NH-	H	C_2H_5	$COCH_3$
84	C_4H_9 (n)	C_4H_9 (n)	-NH-	H	C_2H_5	H
85	C_4H_9 (n)	C_4H_9 (n)	-NH-	H	H	$COCH_2Cl$
86	C_4H_9 (n)	C_4H_9 (n)	-NH-	H	H	COC_4H_9 (n)
87	Cl	C_4H_9 (n)	-NH-	H	H	$COCH_3$







Ex. #	R ¹	R ²	L	B	E	P
88	Cl	C ₄ H ₉ (n)	-NH-	H	H	H
89	C ₄ H ₉ (n)	C ₄ H ₉ (n)	-NHCH ₂ CH ₂ -	H	H	COCH ₃
90	C ₄ H ₉ (n)	C ₄ H ₉ (n)	-NHCH ₂ CH ₂ -	H	H	H
91	C ₄ H ₉ (n)	C ₄ H ₉ (n)	-NHCH ₂ CH ₂ -	H	CH ₃	H
92	C ₄ H ₉ (n)	C ₄ H ₉ (n)	-NHCH ₂ CH ₂ -	H	CH ₃	COCH ₃
93	C ₄ H ₉ (n)	C ₄ H ₉ (n)	-NHCH ₂ CH ₂ -	H	C ₂ H ₅	COCH ₃
94	C ₄ H ₉ (n)	C ₄ H ₉ (n)	-NHCH ₂ CH ₂ -	H	C ₂ H ₅	H
95	C ₄ H ₉ (n)	C ₄ H ₉ (n)	-NHCH ₂ CH ₂ -	H	H	COCH ₂ Cl
96	C ₄ H ₉ (n)	C ₄ H ₉ (n)	-NHCH ₂ CH ₂ -	H	H	COC ₄ H ₉ (n)
97	Cl	C ₄ H ₉ (n)	-NHCH ₂ CH ₂ -	H	H	COCH ₃
98	Cl	C ₄ H ₉ (n)	-NHCH ₂ CH ₂ -	H	H	H

Ex. #	R ¹	R ²	L	B	E	P
99	C ₄ H ₉ (n)	C ₄ H ₉ (n)		*	H	COCH ₃
100	C ₄ H ₉ (n)	C ₄ H ₉ (n)		*	H	H
101	C ₄ H ₉ (n)	C ₄ H ₉ (n)		*	CH ₃	H
102	C ₄ H ₉ (n)	C ₄ H ₉ (n)		*	CH ₃	COCH ₃
103	C ₄ H ₉ (n)	C ₄ H ₉ (n)		*	C ₂ H ₅	COCH ₃
104	C ₄ H ₉ (n)	C ₄ H ₉ (n)		*	C ₂ H ₅	H
105	C ₄ H ₉ (n)	C ₄ H ₉ (n)		*	H	COC ₄ H ₉ (n)
106	C ₄ H ₉ (n)	C ₄ H ₉ (n)		H	H	COC ₄ H ₉ (n)

Ex. #	R ¹	R ²	L	B	E	P
107	Cl	C ₄ H ₉ (n)		*	H	COCH ₃
108	Cl	C ₄ H ₉ (n)		*	H	H
109	C ₂ H ₅	C ₄ H ₉ (n)	-NH-	H	H	COCH ₃
110	C ₂ H ₅	C ₄ H ₉ (n)	-NH-	H	H	H
111	C ₃ H ₇ (n)	C ₄ H ₉ (n)	-NH-	H	H	COCH ₃
112	C ₃ H ₇ (n)	C ₄ H ₉ (n)	-NH-	H	H	H
113	C ₅ H ₁₁ (n)	C ₄ H ₉ (n)	-NH-	H	H	COCH ₃
114	C ₅ H ₁₁ (n)	C ₄ H ₉ (n)	-NH-	H	H	H
115	C ₆ H ₁₃ (n)	C ₄ H ₉ (n)	-NH-	H	H	COCH ₃
116	C ₆ H ₁₃ (n)	C ₄ H ₉ (n)	-NH-	H	H	H
117	CF ₂ CH ₂ CH ₂ CH ₃	C ₄ H ₉ (n)	-NH-	H	H	COCH ₃

Ex. #	R ¹	R ²	L	B	E	P
118	CF ₂ CH ₂ CH ₂ CH ₃	C ₄ H ₉ (n)	-NH-	H	H	H
119	CH ₂ CH ₂ CH(CH ₃) ₂	C ₄ H ₉ (n)	-NH-	H	H	COCH ₃
120	CH ₂ CH ₂ CH(CH ₃) ₂	C ₄ H ₉ (n)	-NH-	H	H	H
121	C ₂ H ₅	C ₂ H ₅	-NH-	H	H	COCH ₃
122	C ₂ H ₅	C ₂ H ₅	-NH-	H	H	H
123	C ₃ H ₇ (n)	C ₃ H ₇ (n)	-NH-	H	H	COCH ₃
124	C ₃ H ₇ (n)	C ₃ H ₇ (n)	-NH-	H	H	COCH ₃
125	C ₅ H ₁₁ (n)	C ₅ H ₁₁ (n)	-NH-	H	H	COCH ₃
126	C ₅ H ₁₁ (n)	C ₅ H ₁₁ (n)	-NH-	H	H	H
127	C ₆ H ₁₃ (n)	C ₆ H ₁₃ (n)	-NH-	H	H	COCH ₃
128	C ₆ H ₁₃ (n)	C ₆ H ₁₃ (n)	-NH-	H	H	H


Ex. #	R ¹	R ²	L	B	E	P
129	C ₂ H ₅	C ₂ H ₅	-NHCH ₂ CH ₂ -	H	H	COCH ₃
130	C ₂ H ₅	C ₂ H ₅	-NHCH ₂ CH ₂ -	H	H	H
131	C ₃ H ₇ (n)	C ₃ H ₇ (n)	-NHCH ₂ CH ₂ -	H	H	COCH ₃
132	C ₃ H ₇ (n)	C ₃ H ₇ (n)	-NHCH ₂ CH ₂ -	H	H	H
133	C ₅ H ₁₁ (n)	C ₅ H ₁₁ (n)	-NHCH ₂ CH ₂ -	H	H	COCH ₃
134	C ₅ H ₁₁ (n)	C ₅ H ₁₁ (n)	-NHCH ₂ CH ₂ -	H	H	H
135	C ₆ H ₁₃ (n)	C ₆ H ₁₃ (n)	-NHCH ₂ CH ₂ -	H	H	COCH ₃
136	C ₆ H ₁₃ (n)	C ₆ H ₁₃ (n)	-NHCH ₂ CH ₂ -	H	H	H
137	C ₂ H ₅	C ₂ H ₅		*	H	COCH ₃
138	C ₂ H ₅	C ₂ H ₅		*	H	H

Ex. #	R ¹	R ²	L	B	E	P
139	C ₃ H ₇ (n)	C ₃ H ₇ (n)		*	H	COCH ₃
140	C ₃ H ₇ (n)	C ₃ H ₇ (n)		*	H	H
141	C ₅ H ₁₁ (n) C ₅ H ₁₁ (n)	C ₅ H ₁₁ (n)		*	H	COCH ₃ ⁸¹
142	C ₅ H ₁₁ (n)	C ₅ H ₁₁ (n)		*	H	H
143	C ₆ H ₁₃ (n)	C ₆ H ₁₃ (n)		*	H	COCH ₃
144	C ₆ H ₁₃ (n)	C ₆ H ₁₃ (n)		*	H	H

* B IS INCORPORATED IN A

Another preferred class of specific conjugates of the invention is provided by conjugates formed from a biphenylmethyl 1H-substituted-1,2,4-triazole AII antagonist compound linked to a cleavable glutamyl residue. Each conjugate of this class contains a terminal amino moiety on the triazole portion of the AII antagonist compound which is connected to a terminal carboxylic acid moiety on the gamma carbon of the cleavable glutamyl residue. Example #145 is a detailed description of a conjugate of this class. Other specific conjugates of Examples #146-#426, as shown in Table V, may be prepared generally in accordance with the procedures of Example #145.


Example 145



N²-acetyl-N-[[5-butyl-1-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]-1H-1,2,4-triazol-3-yl]methyl]-L-glutamine

To a solution of 10.45 g (34.5 mmol) of N-Boc-L-glutamic acid- α -tertbutyl ester (BACHEM) in 100 mL of methylene chloride under nitrogen is added 3.5 g (17.0 mmol) of solid dicyclohexylcarbodiimide (DCC). The reaction is allowed to stir for 2 h and filtered under nitrogen. The anhydride solution is then added to a solution of 5.78 g (14.9 mmol) of the compound of Example 3 in 75 mL of methylene chloride under nitrogen. The reaction is stirred overnight and concentrated in vacuo. Purification by silica gel chromatography (Waters Prep-500A) gives pure material by thin-layer chromatography (TLC). This material is redissolved in 100 mL of methylene chloride under nitrogen and is cooled to 0°C prior to the addition of 135 mL of TFA. The stirred reaction is allowed to warm to ambient temperature overnight and is concentrated in vacuo. The crude product is dissolved in 80 mL of acetonitrile/water (1:1) and the pH is adjusted to 9 with 1 M K₂CO₃. The solution is cooled to 0°C and 1.1 mL (11 mmol) of acetic anhydride and 11 mL (11 mmol) of 1 M K₂CO₃ is added every 30 min for 5 h; during the course of

this reaction the pH is maintained at 9 and the reaction temperature is kept below 5°C. After the last addition, the reaction is allowed to warm to ambient temperature overnight. The pH is adjusted to 4 with 3 M HCl and the reaction is concentrated to 300 mL. Purification by reverse phase chromatography (Waters Delta prep-3000) gives N²-acetyl-N-[[5-butyl-1-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl)methyl]-1H-1,2,4-triazol-3-yl)methyl]-L-glutamine.

TABLE V

Ex: #	R ₅	A	B	E	P
146	CO ₂ H	single bond	H	H	COCH ₃
147	CO ₂ H	single bond	H	H	COCH ₂ Cl
148	CO ₂ H	single bond	H	H	COC ₄ H ₉
149	CO ₂ H	single bond	H	CH ₃	COCH ₃
150	CO ₂ H	single bond	H	C ₂ H ₅	COCH ₃
151	CN ₄ H	single bond	H	H	COCH ₃
152	CN ₄ H	single bond	H	H	COCH ₂ Cl
153	CN ₄ H	single bond	H	H	COC ₄ H ₉

Ex: #	R ₅	A	B	E	P
154	CN ₄ H	single bond	H	CH ₃	COCH ₃
155	CN ₄ H	single bond	H	C ₂ H ₅	COCH ₃
156	CO ₂ H	single bond	H	H	H
157	CO ₂ H	single bond	H	CH ₃	H
158	CO ₂ H	single bond	H	C ₂ H ₅	H
159	CN ₄ H	single bond	H	H	H
160	CN ₄ H	single bond	H	CH ₃	H
161	CN ₄ H	single bond	H	C ₂ H ₅	H
162	CO ₂ H	-CH ₂ -	H	H	COCH ₃
163	CO ₂ H	single bond	H	H	COCH ₂ Cl
164	CO ₂ H	single bond	H	H	COC ₄ H ₉
165	CO ₂ H	single bond	H	CH ₃	COCH ₃
166	CO ₂ H	single bond	H	C ₂ H ₅	COCH ₃

268

Ex: #	R ₅	A	B	E	P
167	CN ₄ H	single bond	H	H	COCH ₂ Cl
168	CN ₄ H	single bond	H	H	COC ₄ H ₉
169	CN ₄ H	single bond	H	CH ₃	COCH ₃
170	CN ₄ H	single bond	H	C ₂ H ₅	COCH ₃
171	CO ₂ H	-CH ₂ -	H	H	H
172	CO ₂ H	single bond	H	CH ₃	H
173	CO ₂ H	single bond	H	C ₂ H ₅	H
174	CN ₄ H	-CH ₂ -	H	H	H
175	CN ₄ H	single bond	H	CH ₃	H
176	CN ₄ H	single bond	H	C ₂ H ₅	H
177	CN ₄ H	-CH ₂ -	CH ₃	H	H
178	CN ₄ H	-CH ₂ -	CH ₃	H	COCH ₃
179	CO ₂ H	-CH ₂ CH ₂ -	H	H	COCH ₃

Ex: #	R ₅	A	B	E	P
180	CO ₂ H	single bond	H	H	COCH ₂ Cl
181	CO ₂ H	single bond	H	H	COC ₄ H ₉
182	CO ₂ H	single bond	H	CH ₃	COCH ₃
183	CO ₂ H	single bond	H	C ₂ H ₅	COCH ₃
184	CN ₄ H	-CH ₂ CH ₂ -	H	H	COCH ₃
185	CN ₄ H	single bond	H	H	COCH ₂ Cl
186	CN ₄ H	single bond	H	H	COC ₄ H ₉
187	CN ₄ H	single bond	H	CH ₃	COCH ₃
188	CN ₄ H	single bond	H	C ₂ H ₅	COCH ₃
189	CO ₂ H	-CH ₂ CH ₂ -	H	H	H
190	CO ₂ H	single bond	H	CH ₃	H
191	CO ₂ H	single bond	H	C ₂ H ₅	H
192	CN ₄ H	-CH ₂ CH ₂ -	H	H	H


270

Ex: #	R ₅	A	B	E	P
193	CN ₄ H	single bond	H	CH ₃	H
194	CN ₄ H	single bond	H	C ₂ H ₅	H
195	CO ₂ H	C ₃ H ₆ (n)	H	H	COCH ₃
196	CO ₂ H	single bond	H	H	COCH ₂ Cl
197	CO ₂ H	single bond	H	H	COC ₄ H ₉
198	CO ₂ H	single bond	H	CH ₃	COCH ₃
199	CO ₂ H	single bond	H	C ₂ H ₅	COCH ₃
200	CN ₄ H	C ₃ H ₆ (n)	H	H	COCH ₃
201	CN ₄ H	single bond	H	H	COCH ₂ Cl
202	CN ₄ H	single bond	H	H	COC ₄ H ₉
203	CN ₄ H	single bond	H	CH ₃	COCH ₃
204	CN ₄ H	single bond	H	C ₂ H ₅	COCH ₃
205	CO ₂ H	C ₃ H ₆ (n)	H	H	H





271




Ex: #	R ₅	A	B	E	P
206	CO ₂ H	single bond	H	CH ₃	H
207	CO ₂ H	single bond	H	C ₂ H ₅	H
208	CN ₄ H	C ₃ H ₆ (n)	H	H	H
209	CN ₄ H	single bond	H	CH ₃	H
210	CN ₄ H	single bond	H	C ₂ H ₅	H
211	CO ₂ H	C ₄ H ₈ (n)	H	H	COCH ₃
212	CO ₂ H	single bond	H	H	COCH ₂ Cl
213	CO ₂ H	single bond	H	H	COC ₄ H ₉
214	CO ₂ H	single bond	H	CH ₃	COCH ₃
215	CO ₂ H	single bond	H	C ₂ H ₅	COCH ₃
216	CN ₄ H	C ₄ H ₈ (n)	H	H	COCH ₃
217	CN ₄ H	single bond	H	H	COCH ₂ Cl
218	CN ₄ H	single bond	H	H	COC ₄ H ₉




272

Ex: #	R ₅	A	B	E	P
219	CN ₄ H	single bond	H	CH ₃	COCH ₃
220	CN ₄ H	single bond	H	C ₂ H ₅	COCH ₃
221	CO ₂ H	C ₄ H ₈ (n)	H	H	H
222	CO ₂ H	single bond	H	CH ₃	H
223	CO ₂ H	single bond	H	C ₂ H ₅	H
224	CN ₄ H	C ₄ H ₈ (n)	H	H	H
225	CN ₄ H	single bond	H	CH ₃	H
226	CN ₄ H	single bond	H	C ₂ H ₅	H
227	CO ₂ H		H	H	COCH ₃
228	CO ₂ H	single bond	H	H	COCH ₂ Cl
229	CO ₂ H	single bond	H	H	COC ₄ H ₉
230	CO ₂ H	single bond	H	CH ₃	COCH ₃
231	CO ₂ H	single bond	H	C ₂ H ₅	COCH ₃




273




Ex: #	R ₅	A	B	E	P
232	CN ₄ H		H	H	COCH ₃
233	CN ₄ H	single bond	H	H	COCH ₂ Cl
234	CN ₄ H	single bond	H	H	COC ₄ H ₉
235	CN ₄ H	single bond	H	CH ₃	COCH ₃
236	CN ₄ H	single bond	H	C ₂ H ₅	COCH ₃
237	CO ₂ H		H	H	H
238	CO ₂ H	single bond	H	CH ₃	H
239	CO ₂ H	single bond	H	C ₂ H ₅	H
240	CN ₄ H		H	H	H
241	CN ₄ H	single bond	H	CH ₃	H
242	CN ₄ H	single bond	H	C ₂ H ₅	H
243	CO ₂ H		H	H	COCH ₃
244	CO ₂ H	single bond	H	H	COCH ₂ Cl





Ex: #	R ₅	A	B	E	P
245	CO ₂ H	single bond	H	H	COC ₄ H ₉
246	CO ₂ H	single bond	H	CH ₃	COCH ₃
247	CO ₂ H	single bond	H	C ₂ H ₅	COCH ₃
248	CN ₄ H	-CH ₂ - 	H	H	COCH ₃
249	CN ₄ H	single bond	H	H	COCH ₂ Cl
250	CN ₄ H	single bond	H	H	COC ₄ H ₉
251	CN ₄ H	single bond	H	CH ₃	COCH ₃
252	CN ₄ H	single bond	H	C ₂ H ₅	COCH ₃
253	CO ₂ H	-CH ₂ - 	H	H	H
254	CO ₂ H	single bond	H	CH ₃	H
255	CO ₂ H	single bond	H	C ₂ H ₅	H
256	CN ₄ H	-CH ₂ - 	H	H	H




Ex: #	R ₅	A	B	E	P
257	CN ₄ H	single bond	H	CH ₃	H
258	CN ₄ H	single bond	H	C ₂ H ₅	H
259	CO ₂ H		H	H	COCH ₃
260	CO ₂ H	single bond	H	H	COCH ₂ Cl
261	CO ₂ H	single bond	H	H	COC ₄ H ₉
262	CO ₂ H	single bond	H	CH ₃	COCH ₃
263	CO ₂ H	single bond	H	C ₂ H ₅	COCH ₃
264	CN ₄ H		H	H	COCH ₃
265	CN ₄ H	single bond	H	H	COCH ₂ Cl
266	CN ₄ H	single bond	H	H	COC ₄ H ₉
267	CN ₄ H	single bond	H	CH ₃	COCH ₃
268	CN ₄ H	single bond	H	C ₂ H ₅	COCH ₃
269	CO ₂ H		H	H	H




276




Ex: #	R ₅	A	B	E	P
270	CO ₂ H	single bond	H	CH ₃	H
271	CO ₂ H	single bond	H	C ₂ H ₅	H
272	CN ₄ H		H	H	H
273	CN ₄ H	single bond	H	CH ₃	H
274	CN ₄ H	single bond	H	C ₂ H ₅	H
275	CO ₂ H		H	H	COCH ₃
276	CO ₂ H	single bond	H	H	COCH ₂ Cl
277	CO ₂ H	single bond	H	H	COC ₄ H ₉
278	CO ₂ H	single bond	H	CH ₃	COCH ₃
279	CO ₂ H	single bond	H	C ₂ H ₅	COCH ₃
280	CN ₄ H		H	H	COCH ₃
281	CN ₄ H	single bond	H	H	COCH ₂ Cl
282	CN ₄ H	single bond	H	H	COC ₄ H ₉

Ex: #	R ₅	A	B	E	P
283	CN ₄ H	single bond	H	CH ₃	COCH ₃
284	CN ₄ H	single bond	H	C ₂ H ₅	COCH ₃
285	CO ₂ H	-CH ₂ -  -CH ₂ -	H	H	H
286	CO ₂ H	single bond	H	CH ₃	H
287	CO ₂ H	single bond	H	C ₂ H ₅	H
288	CN ₄ H	-CH ₂ -  -CH ₂ -	H	H	H
289	CN ₄ H	single bond	H	CH ₃	H
290	CN ₄ H	single bond	H	C ₂ H ₅	H
291	CN ₄ H	-CH ₂ CH ₂ -  -	H	H	COCH ₃
292	CO ₂ H	single bond	H	H	COCH ₂ Cl
293	CO ₂ H	single bond	H	H	COC ₄ H ₉
294	CO ₂ H	single bond	H	CH ₃	COCH ₃
295	CO ₂ H	single bond	H	C ₂ H ₅	COCH ₃





Ex: #	R ₅	A	B	E	P
296	CN ₄ H		H	H	H
297	CN ₄ H	single bond	H	H	COCH ₂ Cl
298	CN ₄ H	single bond	H	H	COC ₄ H ₉
299	CN ₄ H	single bond	H	CH ₃	COCH ₃
300	CN ₄ H	single bond	H	C ₂ H ₅	COCH ₃
301	CN ₄ H		H	H	COCH ₃
302	CO ₂ H	single bond	H	CH ₃	H
303	CO ₂ H	single bond	H	C ₂ H ₅	H
304	CN ₄ H		H	H	H
305	CN ₄ H	single bond	H	CH ₃	H
306	CN ₄ H	single bond	H	C ₂ H ₅	H
307	CO ₂ H		H	H	COCH ₃
308	CO ₂ H	single bond	H	H	COCH ₂ Cl




Ex: #	R ₅	A	B	E	P
309	CO ₂ H	single bond	H	H	COC ₄ H ₉
310	CO ₂ H	single bond	H	CH ₃	COCH ₃
311	CO ₂ H	single bond	H	C ₂ H ₅	COCH ₃
312	CN ₄ H		H	H	COCH ₃
313	CN ₄ H	single bond	H	H	COCH ₂ Cl
314	CN ₄ H	single bond	H	H	COC ₄ H ₉
315	CN ₄ H	single bond	H	CH ₃	COCH ₃
316	CN ₄ H	single bond	H	C ₂ H ₅	COCH ₃
317	CO ₂ H		H	H	H
318	CO ₂ H	single bond	H	CH ₃	H
319	CO ₂ H	single bond	H	C ₂ H ₅	H
320	CN ₄ H		H	H	H
321	CN ₄ H	single bond	H	CH ₃	H

Ex: #	R ₅	A	B	E	P
322	CN ₄ H	single bond	H	C ₂ H ₅	H
323	CO ₂ H	-CH ₂ - 	H	H	COCH ₃
324	CO ₂ H	single bond	H	H	COCH ₂ Cl
325	CO ₂ H	single bond	H	H	COC ₄ H ₉
326	CO ₂ H	single bond	H	CH ₃	COCH ₃
327	CO ₂ H	single bond	H	C ₂ H ₅	COCH ₃
328	CN ₄ H	-CH ₂ - 	H	H	COCH ₃
329	CN ₄ H	single bond	H	H	COCH ₂ Cl
330	CN ₄ H	single bond	H	H	COC ₄ H ₉
331	CN ₄ H	single bond	H	CH ₃	COCH ₃
332	CN ₄ H	single bond	H	C ₂ H ₅	COCH ₃
333	CO ₂ H	-CH ₂ - 	H	H	H
334	CO ₂ H	single bond	H	CH ₃	H
335	CO ₂ H	single bond	H	C ₂ H ₅	H





Ex: #	R ₅	A	B	E	P
336	CN ₄ H	-CH ₂ - 	H	H	H
337	CN ₄ H	single bond	H	CH ₃	H
338	CN ₄ H	single bond	H	C ₂ H ₅	H
339	CO ₂ H	 -CH ₂ -	H	H	COCH ₃
340	CO ₂ H	single bond	H	H	COCH ₂ Cl
341	CO ₂ H	single bond	H	H	COC ₄ H ₉
342	CO ₂ H	single bond	H	CH ₃	COCH ₃
343	CO ₂ H	single bond	H	C ₂ H ₅	COCH ₃
344	CN ₄ H	 -CH ₂ -	H	H	COCH ₃
345	CN ₄ H	single bond	H	H	COCH ₂ Cl
346	CN ₄ H	single bond	H	H	COC ₄ H ₉
347	CN ₄ H	single bond	H	CH ₃	COCH ₃
348	CN ₄ H	single bond	H	C ₂ H ₅	COCH ₃




282





Ex: #	R ₅	A	B	E	P
349	CO ₂ H		H	H	H
350	CO ₂ H	single bond	H	CH ₃	H
351	CO ₂ H	single bond	H	C ₂ H ₅	H
352	CN ₄ H		H	H	H
353	CN ₄ H	single bond	H	CH ₃	H
354	CN ₄ H	single bond	H	C ₂ H ₅	H
355	CO ₂ H		H	H	COCH ₃
356	CO ₂ H	single bond	H	H	COCH ₂ Cl
357	CO ₂ H	single bond	H	H	COC ₄ H ₉
358	CO ₂ H	single bond	H	CH ₃	COCH ₃
359	CO ₂ H	single bond	H	C ₂ H ₅	COCH ₃
360	CN ₄ H		H	H	COCH ₃
361	CN ₄ H	single bond	H	H	COCH ₂ Cl
362	CN ₄ H	single bond	H	H	COC ₄ H ₉

Ex: #	R ₅	A	B	E	P
363	CN ₄ H	single bond	H	CH ₃	COCH ₃
364	CN ₄ H	single bond	H	C ₂ H ₅	COCH ₃
365	CO ₂ H		H	H	H
366	CO ₂ H	single bond	H	CH ₃	H
367	CO ₂ H	single bond	H	C ₂ H ₅	H
368	CN ₄ H		H	H	H
369	CN ₄ H	single bond	H	CH ₃	H
370	CN ₄ H	single bond	H	C ₂ H ₅	H
371	CN ₄ H		H	H	COCH ₃
372	CO ₂ H	single bond	H	H	COCH ₂ Cl
373	CO ₂ H	single bond	H	H	COC ₄ H ₉
374	CO ₂ H	single bond	H	CH ₃	COCH ₃
375	CO ₂ H	single bond	H	C ₂ H ₅	COCH ₃


284


Ex: #	R ₅	A	B	E	P
376	CN ₄ H	-CH ₂ CH ₂ - 	H	H	H
377	CN ₄ H	single bond	H	H	COCH ₂ Cl
378	CN ₄ H	single bond	H	H	COC ₄ H ₉
379	CN ₄ H	single bond	H	CH ₃	COCH ₃
380	CN ₄ H	single bond	H	C ₂ H ₅	COCH ₃
381	CN ₄ H	 -CH ₂ CH ₂ -	H	H	COCH ₃
382	CO ₂ H	single bond	H	CH ₃	H
383	CO ₂ H	single bond	H	C ₂ H ₅	H
384	CN ₄ H	 -CH ₂ CH ₂ -	H	H	H
385	CN ₄ H	single bond	H	CH ₃	H
386	CN ₄ H	single bond	H	C ₂ H ₅	H
387	CN ₄ H		*	H	COCH ₃
388	CO ₂ H	single bond	H	H	COCH ₂ Cl

Ex: #	R ₅	A	B	E	P
389	CO ₂ H	single bond	H	H	COC ₄ H ₉
390	CO ₂ H	single bond	H	CH ₃	COCH ₃
391	CO ₂ H	single bond	H	C ₂ H ₅	COCH ₃
392	CN ₄ H		*	H	H
393	CN ₄ H	single bond	H	H	COCH ₂ Cl
394	CN ₄ H	single bond	H	H	COC ₄ H ₉
395	CN ₄ H	single bond	H	CH ₃	COCH ₃
396	CN ₄ H	single bond	H	C ₂ H ₅	COCH ₃
397	CN ₄ H		*	H	COCH ₃
398	CO ₂ H	single bond	H	CH ₃	H
399	CO ₂ H	single bond	H	C ₂ H ₅	H
400	CN ₄ H		*	H	H
401	CN ₄ H	single bond	H	CH ₃	H
402	CN ₄ H	single bond	H	C ₂ H ₅	H

Ex: #	R ₅	A	B	E	P
403	CN ₄ H	-CH ₂ CH ₂ - 	*	H	COCH ₃
404	CO ₂ H	single bond	H	H	COCH ₂ Cl
405	CO ₂ H	single bond	H	H	COC ₄ H ₉
406	CO ₂ H	single bond	H	CH ₃	COCH ₃
407	CO ₂ H	single bond	H	C ₂ H ₅	COCH ₃
408	CN ₄ H	-CH ₂ -CH ₂ - 	*	H	H
409	CN ₄ H	single bond	H	H	COCH ₂ Cl
410	CN ₄ H	single bond	H	H	COC ₄ H ₉
411	CN ₄ H	single bond	H	CH ₃	COCH ₃
412	CN ₄ H	single bond	H	C ₂ H ₅	COCH ₃
413	CN ₄ H	-CH ₂ -N- 	*	H	COCH ₃
414	CO ₂ H	single bond	H	CH ₃	H
415	CO ₂ H	single bond	H	C ₂ H ₅	H
416	CN ₄ H	-CH ₂ -N- 	*	H	H

287


Ex: #	R ₅	A	B	E	P
417	CN ₄ H	single bond	H	CH ₃	H
418	CN ₄ H	single bond	H	C ₂ H ₅	H
419	CN ₄ H	-CH ₂ -CH ₂ -N 	*	H	COCH ₃
420	CN ₄ H	single bond	H	H	COCH ₂ Cl

Ex: #	R ₅	A	B	E	P
421	CN ₄ H	single bond	H	H	COC ₄ H ₉
422	CN ₄ H	single bond	H	CH ₃	COCH ₃
423	CN ₄ H	single bond	H	C ₂ H ₅	COCH ₃
424	CN ₄ H	-CH ₂ CH ₂ -N 	*	H	H
425	CN ₄ H	single bond	H	CH ₃	H
426	CN ₄ H	single bond	H	C ₂ H ₅	H

* B is incorporated in A

Another preferred class of specific conjugates of the invention is provided by conjugates formed from a biphenylmethyl 1H-substituted-1,2,4-triazole AII antagonist compound linked to a cleavable glutamyl residue. Each conjugate of this class contains a diamino linker moiety which connects a terminal carboxylic acid moiety on the triazole portion of the AII antagonist compound with a terminal carboxylic acid moiety on the gamma carbon of the cleavable glutamyl residue. Example #427 is a detailed description of a conjugate of this class. Other specific conjugates of Examples #428-#834, as shown in Table VI, may be prepared generally in accordance with the procedures of Example #427.

Example 427



N-acetyl-L-glutamic acid, 5-[5-butyl-1-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl-1H-1,2,4-triazol-3-yl]acetylhydrazide


Step 1: Preparation of 5-[4'-[(5-butyl-3-hydrazinylcarbonylmethyl-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole

10



Under nitrogen, 8.34 g (20 mmol) of the compound of Example 7 is dissolved in 200 mL of absolute methanol at -10°C (ice/methanol) and is treated dropwise with 3.0 g (25 mmol) of thionyl chloride. After the addition is complete, the reaction is allowed to warm to ambient temperature for 2 h prior to stirring at reflux overnight. All volatiles are removed in vacuo and the residue redissolved in 100 mL of methanol. Under nitrogen, 20 g (625 mmol) of anhydrous hydrazine is added and the reaction is stirred at reflux overnight. Concentration in vacuo gives the crude product; purification by silica gel chromatography provides 5-[4'-[(5-butyl-3-hydrazinocarbonylmethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole.







Step 2: Preparation of N-acetyl-L-glutamic acid, 5-[5-butyl-1-
[[2'-(1H-tetrazol-5-yl)][1,1'-biphenyl]-4-yl]methyl-1H-1,2,4-
triazol-3-yl]acetylhydrazide

5 To a solution of 10.45 g (34.5 mmol) of N-Boc-L-glutamic acid- α -tertbutyl ester (BACHEM) in 100 mL of methylene chloride under nitrogen is added 3.5 g (17.0 mmol) of solid dicyclohexylcarbodiimide (DCC). The reaction is allowed to stir for 2 h and filtered under nitrogen. The anhydride solution is
10 then added to a solution of 6.42 g (14.9 mmol) of the compound of Example 20 in 75 mL of methylene chloride under nitrogen. The reaction is stirred overnight and is concentrated in vacuo. Purification by silica gel chromatography (Waters Prep-500A) gives pure material by thin-layer chromatography (TLC). This
15 material is redissolved in 100 mL of methylene chloride under nitrogen and cooled to 0°C prior to the addition of 135 mL of TFA. The stirred reaction is allowed to warm to ambient temperature overnight and is concentrated in vacuo. The crude product is dissolved in 80 mL of acetonitrile/water (1:1) and the
20 pH is adjusted to 9 with 1 M K₂CO₃. The solution is cooled to 0°C and 1.1 mL (11 mmol) of acetic anhydride and 11 mL (11 mmol) of 1 M K₂CO₃ is added every 30 min for 5 h; during the course of this reaction the pH is maintained at 9 and the reaction temperature is kept below 5°C. After the last addition, the
25 reaction is allowed to warm to ambient temperature overnight. The pH was adjusted to 4 with 3 M HCl and the reaction concentrated to 300 mL. Purification by reverse phase chromatography (Waters Delta prep-3000) gives N-acetyl-L-glutamic acid, 5-[5-butyl-1-[[2'-(1H-tetrazol-5-yl)][1,1'-biphenyl]-4-
30 yl]methyl-1H-1,2,4-triazol-3-yl]acetylhydrazide.









TABLE VI

Ex. #	A	L	B	E	P
428	single bond	-NH-	H	H	COCH ₃
429	single bond	-NH-	H	H	COCH ₂ Cl
430	single bond	-NH-	H	H	COC ₄ H ₉
431	single bond	-NH-	H	CH ₃	COCH ₃
432	single bond	-NH-	H	C ₂ H ₅	COCH ₃
433	single bond	-NH-	H	H	H
434	single bond	-NH-	H	CH ₃	H







Ex. #	A	L	B	E	P
435	single bond	-NH-	H	C ₂ H ₅	H
436	single bond	-NHCH ₂ CH ₂ -	H	H	COCH ₃
437	single bond	-NHCH ₂ CH ₂ -	H	H	COCH ₂ Cl
438	single bond	-NHCH ₂ CH ₂ -	H	H	COC ₄ H ₉
439	single bond	-NHCH ₂ CH ₂ -	H	CH ₃	COCH ₃
440	single bond	-NHCH ₂ CH ₂ -	H	C ₂ H ₅	COCH ₃
441	single bond	-NHCH ₂ CH ₂ -	H	H	H
442	single bond	-NHCH ₂ CH ₂ -	H	CH ₃	H
443	single bond	-NHCH ₂ CH ₂ -	H	C ₂ H ₅	H
444	single bond		*	H	COCH ₃
445	single bond		H	H	COCH ₂ Cl



Ex. #	A	L	B	E	P
446	single bond		H	H	COC ₄ H ₉
447	single bond		H	CH ₃	COCH ₃
448	single bond		H	C ₂ H ₅	COCH ₃
449	single bond		*	H	H
450	single bond		H	CH ₃	H
451	single bond		H	C ₂ H ₅	H
452	CH ₂	-NH-	H	H	COCH ₂ Cl
453	CH ₂	-NH-	H	H	COC ₄ H ₉
454	CH ₂	-NH-	H	CH ₃	COCH ₃




Ex. #	A	L	B	E	P
455	CH ₂	-NH-	H	C ₂ H ₅	COCH ₃
456	CH ₂	-NH-	H	H	H
457	CH ₂	-NH-	H	CH ₃	H
458	CH ₂	-NH-	H	C ₂ H ₅	H
459	CH ₂	-NHCH ₂ CH ₂ -	H	H	COCH ₃
460	CH ₂	-NHCH ₂ CH ₂ -	H	H	COCH ₂ Cl
461	CH ₂	-NHCH ₂ CH ₂ -	H	H	COC 4H ₉
462	CH ₂	-NHCH ₂ CH ₂ -	H	CH ₃	COCH ₃
463	CH ₂	-NHCH ₂ CH ₂ -	H	C ₂ H ₅	COCH ₃
464	CH ₂	-NHCH ₂ CH ₂ -	H	H	H
465	CH ₂	-NHCH ₂ CH ₂ -	H	CH ₃	H
466	CH ₂	-NHCH ₂ CH ₂ -	H	C ₂ H ₅	H









Ex. #	A	L	B	E	P
467	CH ₂		*	H	COCH ₃
468	CH ₂		H	H	COCH ₂ Cl
469	CH ₂		H	H	COC ₄ H ₉
470	CH ₂		H	CH ₃	COCH ₃
471	CH ₂		H	C ₂ H ₅	COCH ₃
472	CH ₂		*	H	H
473	CH ₂		H	CH ₃	H
474	CH ₂		H	C ₂ H ₅	H









Ex. #	A	L	B	E	P
475	CH ₂ CH ₂	-NH-	H	H	COCH ₃
476	CH ₂ CH ₂	-NH-	H	H	COCH ₂ Cl
477	CH ₂ CH ₂	-NH-	H	H	COC ₄ H ₉
478	CH ₂ CH ₂	-NH-	H	CH ₃	COCH ₃
479	CH ₂ CH ₂	-NH-	H	C ₂ H ₅	COCH ₃
480	CH ₂ CH ₂	-NH-	H	H	H
481	CH ₂ CH ₂	-NH-	H	CH ₃	H
482	CH ₂ CH ₂	-NH-	H	C ₂ H ₅	H
483	CH ₂ CH ₂	-NHCH ₂ CH ₂ -	H	H	COCH ₃
484	CH ₂ CH ₂	-NHCH ₂ CH ₂ -	H	H	COCH ₂ Cl
485	CH ₂ CH ₂	-NHCH ₂ CH ₂ -	H	H	COC ₄ H ₉
486	CH ₂ CH ₂	-NHCH ₂ CH ₂ -	H	CH ₃	COCH ₃
487	CH ₂ CH ₂	-NHCH ₂ CH ₂ -	H	C ₂ H ₅	COCH ₃






Ex. #	A	L	B	E	P
488	CH ₂ CH ₂	-NHCH ₂ CH ₂ -	H	H	H
489	CH ₂ CH ₂	-NHCH ₂ CH ₂ -	H	CH ₃	H
490	CH ₂ CH ₂	-NHCH ₂ CH ₂ -	H	C ₂ H ₅	H
491	CH ₂ CH ₂		*	H	COCH ₃
492	CH ₂ CH ₂		H	H	COCH ₂ Cl
493	CH ₂ CH ₂		H	H	COC ₄ H ₉
494	CH ₂ CH ₂		H	CH ₃	COCH ₃
495	CH ₂ CH ₂		H	C ₂ H ₅	COCH ₃
496	CH ₂ CH ₂		*	H	H


Ex. #	A	L	B	E	P
497	CH ₂ CH ₂		H	CH ₃	H
498	CH ₂ CH ₂		H	C ₂ H ₅	H
499	C ₃ H ₆ (n)	-NH-	H	H	COCH ₃
500	C ₃ H ₆ (n)	-NH-	H	H	COCH ₂ Cl
501	C ₃ H ₆ (n)	-NH-	H	H	COC ₄ H ₉
502	C ₃ H ₆ (n)	-NH-	H	CH ₃	COCH ₃
503	C ₃ H ₆ (n)	-NH-	H	C ₂ H ₅	COCH ₃
504	C ₃ H ₆ (n)	-NH-	H	H	H
505	C ₃ H ₆ (n)	-NH-	H	CH ₃	H
506	C ₃ H ₆ (n)	-NH-	H	C ₂ H ₅	H
507	C ₃ H ₆ (n)	-NHCH ₂ CH ₂ -	H	H	COCH ₃










Ex. #	A	L	B	E	P
508	C ₃ H ₆ (n)	-NHCH ₂ CH ₂ -	H	H	COCH ₂ Cl
509	C ₃ H ₆ (n)	-NHCH ₂ CH ₂ -	H	H	COC ₄ H ₉
510	C ₃ H ₆ (n)	-NHCH ₂ CH ₂ -	H	CH ₃	COCH ₃
511	C ₃ H ₆ (n)	-NHCH ₂ CH ₂ -	H	C ₂ H ₅	COCH ₃
512	C ₃ H ₆ (n)	-NHCH ₂ CH ₂ -	H	H	H
513	C ₃ H ₆ (n)	-NHCH ₂ CH ₂ -	H	CH ₃	H
514	C ₃ H ₆ (n)	-NHCH ₂ CH ₂ -	H	C ₂ H ₅	H
515	C ₃ H ₆ (n)		*	H	COCH ₃
516	C ₃ H ₆ (n)		H	H	COCH ₂ Cl
517	C ₃ H ₆ (n)		H	H	COC ₄ H ₉

















Ex. #	A	L	B	E	P
540	C ₄ H ₈ (n)		H	H	COCH ₂ Cl
541	C ₄ H ₈ (n)		H	H	COC ₄ H ₉
542	C ₄ H ₈ (n)		H	CH ₃	COCH ₃
543	C ₄ H ₈ (n)		H	C ₂ H ₅	COCH ₃
544	C ₄ H ₈ (n)		*	H	H
545	C ₄ H ₈ (n)		H	CH ₃	H
546	C ₄ H ₈ (n)		H	C ₂ H ₅	H
547		-NH-	H	H	COCH ₃









Ex. #	A	L	B	E	P
548		-NH-	H	H	COCH ₂ Cl
549		-NH-	H	H	COC ₄ H ₉
550		-NH-	H	CH ₃	COCH ₃
551		-NH-	H	C ₂ H ₅	COCH ₃
552		-NH-	H	H	H
553		-NH-	H	CH ₃	H
554		-NH-	H	C ₂ H ₅	H
555		-NHCH ₂ CH ₂ -	H	H	COCH ₃










Ex. #	A	L	B	E	P
518	C ₃ H ₆ (n)		H	CH ₃	COCH ₃
519	C ₃ H ₆ (n)		H	C ₂ H ₅	COCH ₃
520	C ₃ H ₆ (n)		*	H	H
521	C ₃ H ₆ (n)		H	CH ₃	H
522	C ₃ H ₆ (n)		H	C ₂ H ₅	H
523	C ₄ H ₈ (n)	-NH-	H	H	COCH ₃
524	C ₄ H ₈ (n)	-NH-	H	H	COCH ₂ Cl
525	C ₄ H ₈ (n)	-NH-	H	H	COC ₄ H ₉
526	C ₄ H ₈ (n)	-NH-	H	CH ₃	COCH ₃
527	C ₄ H ₈ (n)	-NH-	H	C ₂ H ₅	COCH ₃

Ex. #	A	L	B	E	P
528	C ₄ H ₈ (n)	-NH-	H	H	H
529	C ₄ H ₈ (n)	-NH-	H	CH ₃	H
530	C ₄ H ₈ (n)	-NH-	H	C ₂ H ₅	H
531	C ₄ H ₈ (n)	-NHCH ₂ CH ₂ -	H	H	COCH ₃
532	C ₄ H ₈ (n)	-NHCH ₂ CH ₂ -	H	H	COCH ₂ Cl
533	C ₄ H ₈ (n)	-NHCH ₂ CH ₂ -	H	H	COC ₄ H ₉
534	C ₄ H ₈ (n)	-NHCH ₂ CH ₂ -	H	CH ₃	COCH ₃
535	C ₄ H ₈ (n)	-NHCH ₂ CH ₂ -	H	C ₂ H ₅	COCH ₃
536	C ₄ H ₈ (n)	-NHCH ₂ CH ₂ -	H	H	H
537	C ₄ H ₈ (n)	-NHCH ₂ CH ₂ -	H	CH ₃	H
538	C ₄ H ₈ (n)	-NHCH ₂ CH ₂ -	H	C ₂ H ₅	H
539	C ₄ H ₈ (n)		*	H	COCH ₃









Ex. #	A	L	B	E	P
556		-NHCH ₂ CH ₂ -	H	H	COCH ₂ Cl
557		-NHCH ₂ CH ₂ -	H	H	COC ₄ H ₉
558		-NHCH ₂ CH ₂ -	H	CH ₃	COCH ₃
559		-NHCH ₂ CH ₂ -	H	C ₂ H ₅	COCH ₃
560		-NHCH ₂ CH ₂ -	H	H	H
561		-NHCH ₂ CH ₂ -	H	CH ₃	H
562		-NHCH ₂ CH ₂ -	H	C ₂ H ₅	H
563			*	H	COCH ₃










Ex. #	A	L	B	E	P
564			H	H	COCH ₂ Cl
565			H	H	COC ₄ H ₉
566			H	CH ₃	COCH ₃
567			H	C ₂ H ₅	COCH ₃
568			*	H	H
569			H	CH ₃	H
570			H	C ₂ H ₅	H
571			H	H	COCH ₃
















Ex. #	A	L	B	E	P
572		-NH-	H	H	COCH ₂ Cl
573		-NH-	H	H	COC ₄ H ₉
574		-NH-	H	CH ₃	COCH ₃
575		-NH-	H	C ₂ H ₅	COCH ₃
576		-NH-	H	H	H
577		-NH-	H	CH ₃	H
578		-NH-	H	C ₂ H ₅	H
579		-NHCH ₂ CH ₂ -	H	H	COCH ₃









Ex. #	A	L	B	E	P
580		$\text{-NHCH}_2\text{CH}_2\text{-}$	H	H	COCH_2Cl
581		$\text{-NHCH}_2\text{CH}_2\text{-}$	H	H	COC_4H_9
582		$\text{-NHCH}_2\text{CH}_2\text{-}$	H	CH_3	COCH_3
583		$\text{-NHCH}_2\text{CH}_2\text{-}$	H	C_2H_5	COCH_3
584		$\text{-NHCH}_2\text{CH}_2\text{-}$	H	H	H
585		$\text{-NHCH}_2\text{CH}_2\text{-}$	H	CH_3	H
586		$\text{-NHCH}_2\text{CH}_2\text{-}$	H	C_2H_5	H
587			*	H	COCH_3










Ex. #	A	L	B	E	P
588			H	H	COCH ₂ Cl
589			H	H	COC ₄ H ₉
590			H	CH ₃	COCH ₃
591			H	C ₂ H ₅	COCH ₃
592			*	H	H
593			H	CH ₃	H
594			H	C ₂ H ₅	H
595			H	H	COCH ₃
















Ex. #	A	L	B	E	P
596	$\text{—CH}_2\text{CH}_2\text{—}$ 	—NH—	H	H	COCH_2Cl
597	$\text{—CH}_2\text{CH}_2\text{—}$ 	—NH—	H	H	COC_4H_9
598	$\text{—CH}_2\text{CH}_2\text{—}$ 	—NH—	H	CH_3	COCH_3
599	$\text{—CH}_2\text{CH}_2\text{—}$ 	—NH—	H	C_2H_5	COCH_3
600	$\text{—CH}_2\text{CH}_2\text{—}$ 	—NH—	H	H	H
601	$\text{—CH}_2\text{CH}_2\text{—}$ 	—NH—	H	CH_3	H
602	$\text{—CH}_2\text{CH}_2\text{—}$ 	—NH—	H	C_2H_5	H
603	$\text{—CH}_2\text{CH}_2\text{—}$ 	$\text{—NHCH}_2\text{CH}_2\text{—}$	H	H	COCH_3









Ex. #	A	L	B	E	P
604		-NHCH ₂ CH ₂ -	H	H	COCH ₂ Cl
605		-NHCH ₂ CH ₂ -	H	H	COC ₄ H ₉
606		-NHCH ₂ CH ₂ -	H	CH ₃	COCH ₃
607		-NHCH ₂ CH ₂ -	H	C ₂ H ₅	COCH ₃
608		-NHCH ₂ CH ₂ -	H	H	H
609		-NHCH ₂ CH ₂ -	H	CH ₃	H
610		-NHCH ₂ CH ₂ -	H	C ₂ H ₅	H
611			*	H	COCH ₃

Ex. #	A	L	B	E	P
612			H	H	COCH ₂ Cl
613			H	H	COC ₄ H ₉
614			H	CH ₃	COCH ₃
615			H	C ₂ H ₅	COCH ₃
616			*	H	H
617			H	CH ₃	H
618			H	C ₂ H ₅	H
619		-NH-	H	H	COCH ₃
















Ex. #	A	L	B	E	P
620		-NH-	H	H	COCH ₂ Cl
621		-NH-	H	H	COC ₄ H ₉
622		-NH-	H	CH ₃	COCH ₃
623		-NH-	H	C ₂ H ₅	COCH ₃
624		-NH-	H	H	H
625		-NH-	H	CH ₃	H
626		-NH-	H	C ₂ H ₅	H
627		-NHCH ₂ CH ₂ -	H	H	COCH ₃









Ex. #	A	L	B	E	P
628		-NHCH ₂ CH ₂ -	H	H	COCH ₂ Cl
629		-NHCH ₂ CH ₂ -	H	H	COC ₄ H ₉
630		-NHCH ₂ CH ₂ -	H	CH ₃	COCH ₃
631		-NHCH ₂ CH ₂ -	H	C ₂ H ₅	COCH ₃
632		-NHCH ₂ CH ₂ -	H	H	H
633		-NHCH ₂ CH ₂ -	H	CH ₃	H
634		-NHCH ₂ CH ₂ -	H	C ₂ H ₅	H
635			*	H	COCH ₃










Ex. #	A	L	B	E	P
636			H	H	COCH ₂ Cl
637			H	H	COC ₄ H ₉
638			H	CH ₃	COCH ₃
639			H	C ₂ H ₅	COCH ₃
640			*	H	H
641			H	CH ₃	H
642			H	C ₂ H ₅	H
643		-NH-	H	H	COCH ₃
















Ex. #	A	L	B	E	P
644	 -CH ₂ -CH ₂ -	-NH-	H	H	COCH ₂ Cl
645	 -CH ₂ -CH ₂ -	-NH-	H	H	COC ₄ H ₉
646	 -CH ₂ -CH ₂ -	-NH-	H	CH ₃	COCH ₃
647	 -CH ₂ -CH ₂ -	-NH-	H	C ₂ H ₅	COCH ₃
648	 -CH ₂ -CH ₂ -	-NH-	H	H	H
649	 -CH ₂ -CH ₂ -	-NH-	H	CH ₃	H
650	 -CH ₂ -CH ₂ -	-NH-	H	C ₂ H ₅	H
651	 -CH ₂ -CH ₂ -	-NHCH ₂ CH ₂ -	H	H	COCH ₃










Ex. #	A	L	B	E	P
652	 -CH ₂ -CH ₂ -	-NHCH ₂ CH ₂ -	H	H	COCH ₂ Cl
653	 -CH ₂ -CH ₂ -	-NHCH ₂ CH ₂ -	H	H	COC ₄ H ₉
654	 -CH ₂ -CH ₂ -	-NHCH ₂ CH ₂ -	H	CH ₃	COCH ₃
655	 -CH ₂ -CH ₂ -	-NHCH ₂ CH ₂ -	H	C ₂ H ₅	COCH ₃
656	 -CH ₂ -CH ₂ -	-NHCH ₂ CH ₂ -	H	H	H
657	 -CH ₂ -CH ₂ -	-NHCH ₂ CH ₂ -	H	CH ₃	H
658	 -CH ₂ -CH ₂ -	-NHCH ₂ CH ₂ -	H	C ₂ H ₅	H
659	 -CH ₂ -CH ₂ -		*	H	COCH ₃













Ex. #	A	L	B	E	P
660			H	H	COCH ₂ Cl
661			H	H	COC ₄ H ₉
662			H	CH ₃	COCH ₃
663			H	C ₂ H ₅	COCH ₃
664			*	H	H
665			H	CH ₃	H
666			H	C ₂ H ₅	H
667		-NH-	H	H	COCH ₃















Ex. #	A	L	B	E	P
668		-NH-	H	H	COCH ₂ Cl
669		-NH-	H	H	COC ₄ H ₉
670		-NH-	H	CH ₃	COCH ₃
671		-NH-	H	C ₂ H ₅	COCH ₃
672		-NH-	H	H	H
673		-NH-	H	CH ₃	H
674		-NH-	H	C ₂ H ₅	H
675		-NHCH ₂ CH ₂ -	H	H	COCH ₃










Ex. #	A	L	B	E	P
676		-NHCH ₂ CH ₂ -	H	H	COCH ₂ Cl
677		-NHCH ₂ CH ₂ -	H	H	COC ₄ H ₉
678		-NHCH ₂ CH ₂ -	H	CH ₃	COCH ₃
679		-NHCH ₂ CH ₂ -	H	C ₂ H ₅	COCH ₃
680		-NHCH ₂ CH ₂ -	H	H	H
681		-NHCH ₂ CH ₂ -	H	CH ₃	H
682		-NHCH ₂ CH ₂ -	H	C ₂ H ₅	H
683			*	H	COCH ₃














Ex. #	A	L	B	E	P
684			H	H	COCH ₂ Cl
685			H	H	COC ₄ H ₉
686			H	CH ₃	COCH ₃
687			H	C ₂ H ₅	COCH ₃
688			*	H	H
689			H	CH ₃	H
690			H	C ₂ H ₅	H
691		-NH-	H	H	COCH ₃












Ex. #	A	L	B	E	P
692		-NH-	H	H	COCH ₂ Cl
693		-NH-	H	H	COC ₄ H ₉
694		-NH-	H	CH ₃	COCH ₃
695		-NH-	H	C ₂ H ₅	COCH ₃
696		-NH-	H	H	H
697		-NH-	H	CH ₃	H
698		-NH-	H	C ₂ H ₅	H
699		-NHCH ₂ CH ₂ -	H	H	COCH ₃
700		-NHCH ₂ CH ₂ -	H	H	COCH ₂ Cl










Ex. #	A	L	B	E	P
701		-NHCH ₂ CH ₂ -	H	H	COC ₄ H ₉
702		-NHCH ₂ CH ₂ -	H	CH ₃	COCH ₃
703		-NHCH ₂ CH ₂ -	H	C ₂ H ₅	COCH ₃
704		-NHCH ₂ CH ₂ -	H	H	H
705		-NHCH ₂ CH ₂ -	H	CH ₃	H
706		-NHCH ₂ CH ₂ -	H	C ₂ H ₅	H
707			*	H	COCH ₃
708			H	H	COCH ₂ Cl
709			H	H	COC ₄ H ₉















Ex. #	A	L	B	E	P
710			H	CH ₃	COCH ₃
711			H	C ₂ H ₅	COCH ₃
712			*	H	H
713			H	CH ₃	H
714			H	C ₂ H ₅	H
715		-NH-	H	H	COCH ₃
716		-NH-	H	H	COCH ₂ Cl
717		-NH-	H	H	COC ₄ H ₉
718		-NH-	H	CH ₃	COCH ₃











Ex. #	A	L	B	E	P
719		-NH-	H	C ₂ H ₅	COCH ₃
720		-NH-	H	H	H
721		-NH-	H	CH ₃	H
722		-NH-	H	C ₂ H ₅	H
723		-NHCH ₂ CH ₂ -	H	H	COCH ₃
724		-NHCH ₂ CH ₂ -	H	H	COCH ₂ Cl
725		-NHCH ₂ CH ₂ -	H	H	COC 4H ₉
726		-NHCH ₂ CH ₂ -	H	CH ₃	COCH ₃
727		-NHCH ₂ CH ₂ -	H	C ₂ H ₅	COCH ₃











Ex. #	A	L	B	E	P
728		$\text{-NHCH}_2\text{CH}_2\text{-}$	H	H	H
729		$\text{-NHCH}_2\text{CH}_2\text{-}$	H	CH ₃	H
730		$\text{-NHCH}_2\text{CH}_2\text{-}$	H	C ₂ H ₅	H
731			*	H	COCH ₃
732			H	H	COCH ₂ Cl
733			H	H	COC ₄ H ₉
734			H	CH ₃	COCH ₃
735			H	C ₂ H ₅	COCH ₃

















Ex. #	A	L	B	E	P
736			*	H	H
737			H	CH ₃	H
738			H	C ₂ H ₅	H
739		-NH-	H	H	COCH ₃
740		-NH-	H	H	COCH ₂ Cl
741		-NH-	H	H	COC ₄ H ₉
742		-NH-	H	CH ₃	COCH ₃
743		-NH-	H	C ₂ H ₅	COCH ₃










Ex. #	A	L	B	E	P
744	$-\text{CH}_2\text{CH}_2-$ 	$-\text{NH}-$	H	H	H
745	$-\text{CH}_2\text{CH}_2-$ 	$-\text{NH}-$	H	CH_3	H
746	$-\text{CH}_2\text{CH}_2-$ 	$-\text{NH}-$	H	C_2H_5	H
747	$-\text{CH}_2\text{CH}_2-$ 	$-\text{NHCH}_2\text{CH}_2-$	H	H	COCH_3
748	$-\text{CH}_2\text{CH}_2-$ 	$-\text{NHCH}_2\text{CH}_2-$	H	H	COCH_2Cl
749	$-\text{CH}_2\text{CH}_2-$ 	$-\text{NHCH}_2\text{CH}_2-$	H	H	COC_4H_9
750	$-\text{CH}_2\text{CH}_2-$ 	$-\text{NHCH}_2\text{CH}_2-$	H	CH_3	COCH_3
751	$-\text{CH}_2\text{CH}_2-$ 	$-\text{NHCH}_2\text{CH}_2-$	H	C_2H_5	C OCH_3
752	$-\text{CH}_2\text{CH}_2-$ 	$-\text{NHCH}_2\text{CH}_2-$	H	H	H










Ex. #	A	L	B	E	P
753		$\text{-NHCH}_2\text{CH}_2\text{-}$	H	CH ₃	H
754		$\text{-NHCH}_2\text{CH}_2\text{-}$	H	C ₂ H ₅	H
755			*	H	COCH ₃
756			H	H	COCH ₂ Cl
757			H	H	COC ₄ H ₉
758			H	CH ₃	COCH ₃
759			H	C ₂ H ₅	COCH ₃
760			*	H	H
















Ex. #	A	L	B	E	P
761			H	CH ₃	H
762			H	C ₂ H ₅	H
763		-NH-	H	H	COCH ₃
764		-NH-	H	H	COCH ₂ Cl
765		-NH-	H	H	COC ₄ H ₉
766		-NH-	H	CH ₃	COCH ₃
767		-NH-	H	C ₂ H ₅	COCH ₃
768		-NH-	H	H	H









Ex. #	A	L	B	E	P
769	 CH ₂ -	-NH-	H	CH ₃	H
770	 CH ₂ -	-NH-	H	C ₂ H ₅	H
771	 CH ₂ -	-NHCH ₂ CH ₂ -	H	H	COCH ₃
772	 CH ₂ -	-NH-	H	H	COCH ₂ Cl
773	 CH ₂ -	-NH-	H	H	COC ₄ H ₉
774	 CH ₂ -	-NH-	H	CH ₃	COCH ₃
775	 CH ₂ -	-NH-	H	C ₂ H ₅	COCH ₃
776	 CH ₂ -	-NHCH ₂ CH ₂ -	H	H	H
777	 CH ₂ -	-NHCH ₂ CH ₂ -	H	CH ₃	H
778	 CH ₂ -	-NHCH ₂ CH ₂ -	H	C ₂ H ₅	H










Ex. #	A	L	B	E	P
779			*	H	COCH ₃
780			H	H	COCH ₂ Cl
781			H	H	COC ₄ H ₉
782			H	CH ₃	COCH ₃
783			C ₂ H ₅		COCH ₃
784			*	H	H
785			H	CH ₃	H
786			H	C ₂ H ₅	H











Ex. #	A	L	B	E	P
787	 CH ₂ CH ₂ -	-NH-	H	H	COCH ₃
788	 CH ₂ CH ₂ -	-NH-	H	H	COCH ₂ Cl
789	 CH ₂ CH ₂ -	-NH-	H	H	COC ₄ H ₉
790	 CH ₂ CH ₂ -	-NH-	H	CH ₃	COCH ₃
791	 CH ₂ CH ₂ -	-NH-	H	C ₂ H ₅	COCH ₃
792	 CH ₂ CH ₂ -	-NH-	H	H	H
793	 CH ₂ CH ₂ -	-NH-	H	CH ₃	H
794	 CH ₂ CH ₂ -	-NH-	H	C ₂ H ₅	H
795	 CH ₂ CH ₂ -	-NHCH ₂ CH ₂ -	H	H	COCH ₃





Ex. #	A	L	B	E	P
796	 CH ₂ CH ₂ -	-NHCH ₂ CH ₂ -	H	H	COCH ₂ Cl
797	 CH ₂ CH ₂ -	-NHCH ₂ CH ₂ -	H	H	COC ₄ H ₉
798	 CH ₂ CH ₂ -	-NHCH ₂ CH ₂ -	H	CH ₃	COCH ₃
799	 CH ₂ CH ₂ -	-NHCH ₂ CH ₂ -	H	C ₂ H ₅	COCH ₃
800	 CH ₂ CH ₂ -	-NHCH ₂ CH ₂ -	H	H	H
801	 CH ₂ CH ₂ -	-NHCH ₂ CH ₂ -	H	CH ₃	H
802	 CH ₂ CH ₂ -	-NHCH ₂ CH ₂ -	H	C ₂ H ₅	H
803	 CH ₂ CH ₂ -		*	H	COCH ₃

Ex. #	A	L	B	E	P
804	 -CH ₂ CH ₂ -		H	H	COCH ₂ Cl
805	 -CH ₂ CH ₂ -		H	H	COC ₄ H ₉
806	 -CH ₂ CH ₂ -		H	CH ₃	COCH ₃
807	 -CH ₂ CH ₂ -		H	C ₂ H ₅	COCH ₃
808	 -CH ₂ CH ₂ -		*	H	H
809	 -CH ₂ CH ₂ -		H	CH ₃	H
810	 -CH ₂ CH ₂ -		H	C ₂ H ₅	H
811	-CH ₂ - 	-NH-	H	H	COCH ₃

Ex. #	A	L	B	E	P
812		-NH-	H	H	COCH ₂ Cl
813		-NH-	H	H	COC ₄ H ₉
814		-NH-	H	CH ₃	COCH ₃
815		-NH-	H	C ₂ H ₅ C	OCH ₃
816		-NH-	H	H	H
817		-NH-	H	CH ₃	H
818		-NH-	H	C ₂ H ₅ H	
819		-NHCH ₂ CH ₂ -	H	H	COCH ₃

Ex. #	A	L	B	E	P
820		-NHCH ₂ CH ₂ -	H	H	COCH ₂ Cl
821		-NHCH ₂ CH ₂ -	H	H	COC ₄ H ₉
822		-NHCH ₂ CH ₂ -	H	CH ₃	COCH ₃
823		-NHCH ₂ CH ₂ -	H	C ₂ H ₅	COCH ₃
824		-NHCH ₂ CH ₂ -	H	H	H
825		-NHCH ₂ CH ₂ -	H	CH ₃	H
826		-NHCH ₂ CH ₂ -	H	C ₂ H ₅	H
827			*	H	COCH ₃

Ex. #	A	L	B	E	P
828			H	H	COCH ₂ Cl
829			H	H	COC ₄ H ₉
830			H	CH ₃	COCH ₃
831			H	C ₂ H ₅	COCH ₃
832			*	H	H

Ex. #	A	L	B	E	P
833			H	CH ₃	H
834			H	C ₂ H ₅	H

BIOLOGICAL EVALUATION

In order to identify suitable angiotension II antagonists for use as the first component of the conjugate of the invention, certain compounds of Examples 1-17 were evaluated variously in three biological assays (Assays "A", "B" and "C"). In a fourth assay, blood-pressure lowering effects of conjugate of the invention of Example 18 were evaluated (Assay "D").

10

Assay A: Angiotensin II Binding Activity

Compounds of Formula I were tested for ability to bind to the smooth muscle angiotensin II receptor using a rat uterine membrane preparation. Angiotensin II (AII) was purchased from Peninsula Labs. ^{125}I -angiotensin II (specific activity of 2200 Ci/mmol) was purchased from Du Pont-New England Nuclear. Other chemicals were obtained from Sigma Chemical Co. This assay was carried out according to the method of Douglas et al [Endocrinology, 106, 120-124 (1980)]. Rat uterine membranes were prepared from fresh tissue. All procedures were carried out at 4°C. Uteri were stripped of fat and homogenized in phosphate-buffered saline at pH 7.4 containing 5 mM EDTA. The homogenate was centrifuged at 1500 x g for 20 min., and the supernatant was recentrifuged at 100,000 x g for 60 min. The pellet was resuspended in buffer consisting of 2 mM EDTA and 50 mM Tris-HCl (pH 7.5) to a final protein concentration of 4 mg/ml. Assay tubes were charged with 0.25 ml of a solution containing 5 mM MgCl_2 , 2 mM EDTA, 0.5% bovine serum albumin, 50 mM Tris-HCl, pH 7.5 and ^{125}I -AII (approximately 10^5 cpm) in the absence or in the presence of unlabelled ligand. The reaction was initiated by the addition of membrane protein and the mixture was incubated at 25°C for 60 min. The incubation was terminated with ice-cold 50 mM Tris-HCl (pH 7.5) and the mixture was filtered to separate membrane-bound labelled peptide from the free ligand. The incubation tube and filter were washed

with ice-cold buffer. Filters were assayed for radioactivity in a Micromedic gamma counter. Nonspecific binding was defined as binding in the presence of 10 μ M of unlabelled AII. Specific binding was calculated as total binding minus nonspecific binding. The receptor binding affinity of an AII antagonist compound was indicated by the concentration (IC_{50}) of the tested AII antagonist which gives 50% displacement of the total specifically bound ^{125}I -AII from the high affinity AII receptor. Binding data were analyzed by a nonlinear least-squares curve fitting program. Results are reported in Table VI.

Assay B: In Vitro Vascular Smooth Muscle-Response for AII

Compounds of Formula I were tested for antagonist activity in rabbit aortic rings. Male New Zealand white rabbits (2-2.5 kg) were sacrificed using an overdose of pentobarbital and exsanguinated via the carotid arteries. The thoracic aorta was removed, cleaned of adherent fat and connective tissue and then cut into 3-mm ring segments. The endothelium was removed from the rings by gently sliding a rolled-up piece of filter paper into the vessel lumen. The rings were then mounted in a water-jacketed tissue bath, maintained at 37°C, between moveable and fixed ends of a stainless steel wire with the moveable end attached to an FT03 Grass transducer coupled to a Model 7D Grass Polygraph for recording isometric force responses. The bath was filled with 20 ml of oxygenated (95% oxygen/5% carbon dioxide) Krebs solution of the following composition (mM): 130 NaCl, 15 $NaHCO_3$, 15 KCl, 1.2 NaH_2PO_4 , 1.2 $MgSO_4$, 2.5 $CaCl_2$, and 11.4 glucose. The preparations were equilibrated for one hour before approximately one gram of passive tension was placed on the rings. Angiotensin II concentration-response curves were then recorded (3×10^{-10} to 1×10^{-5} M). Each concentration of AII was allowed to elicit its maximal contraction, and then AII was washed out repeatedly for 30 minutes before rechallenging with a higher concentration of AII. Aorta rings were exposed to

the test antagonist at 10^{-5} M for 5 minutes before challenging with AII. Adjacent segments of the same aorta ring were used for all concentration-response curves in the presence or absence of the test antagonist. The effectiveness of the test compound was expressed in terms of pA_2 values and were calculated according to H.O. Schild [Br. J. Pharmacol. Chemother., 2,189-206 (1947)]. The pA_2 value is the concentration of the antagonist which increases the EC_{50} value for AII by a factor of 2. Each test antagonist was evaluated in aorta rings from two rabbits. Results are reported in Table VI.

Assay C: In Vivo Intraduodenal Pressor Assay Response for AII Antagonists

Male Sprague-Dawley rats weighing 225-300 grams were anesthetized with Inactin (100 mg/kg, i.p.) and catheters were implanted into the trachea, femoral artery, femoral vein and duodenum. Arterial pressure was recorded from the femoral artery catheter on a Gould chart recorder (Gould, Cleveland, OH). The femoral vein catheter was used for injections of angiotensin II, mecamlamine and atropine. The tracheal catheter allow for airway patency, and the duodenal catheter was used for intraduodenal (i.d.) administration of test compounds. After surgery, the rats were allowed to equilibrate for 30 minutes. Mecamlamine (3 mg/kg, 0.3 ml/kg) and atropine (400 ug/kg, 0.3 ml/kg) were then given i.v. to produce ganglion blockade. These compounds were administered every 90 minutes throughout the test procedure. Angiotensin II was given in bolus doses i.v. (30 ng/kg in saline with 0.5% bovine serum albumin, 0.1 ml/kg) every 10 minutes three times or until the increase in arterial pressure produced was within 3 mmHg for two consecutive AII injections. The last two AII injections were averaged and were taken as the control AII pressor response. Ten minutes after the final control AII injection, the test compound of Formula I (dissolved in sodium bicarbonate) was administered i.d. at a dose of 30

or 100 mg/kg in a volume of 0.2 ml. Angiotensin II injections were then given 5, 10, 20, 30, 45, 60, 75, 90, and 120 minutes after administration of the test compound and response of arterial pressure was monitored. The

5 response to AII was calculated as percent of the control response and then the percent inhibition is calculated as 100 minus the percent control response. Duration of action of a test compound was defined as the time from peak

10 percent inhibition to 50% of peak. One compound at one dose was tested in each rat. Each test compound was tested in two rats and the values for the two rats were averaged. Results are reported in Table IV.

TABLE IV

In Vivo and In Vitro Angiotensin II
Activity of Compounds of the Invention

5	Test Compound Duration Example # (min.)	¹ Assay A	² Assay B	Dose	³ Assay C	
		IC ₅₀ (nM)	pA ₂		Inhibition (%)	
10	1	NT	NT	NT	NT	NT
	2	95	7.37/7.59	10	95	60
				30	98	90-120
	3	5.4	8.70 ± 0.2	10	50	>180
15				30	100	200+
	4	NT	NT	NT	NT	NT
	5	200	7.48/6.91	30	38	20-30
	6	1300	6.55/6.82	100	90	120
	7	84	8.01/8.05	30	90	130
20	8	17,000	NT	NT	NT	NT
	9	700	6.67/6.12	30	80	75
				100	100	130
	10	4.9	8.19/7.59	3	86	100
				30	100	240
25	11	160	6.45/6.77	NT	NT	NT
	12	6.0	8.66/8.59	NT	NT	NT
	13	17	8.70/8.85	NT	NT	NT
	14	7.2	8.84/8.71	NT	NT	NT
	15	16	8.31/8.30	NT	NT	NT
30	16	6.4	8.95/9.24	NT	NT	NT
	17	4.0	8.64/8.40	NT	NT	NT
	18	970	6.14/6.09	NT	NT	NT
	19	12,000	5.18/5.35	NT	NT	NT

	Test	¹ Assay A	² Assay B	Dose	³ Assay C	
	Compound	IC ₅₀	pA ₂		Inhibition	
	Duration					
5	Example #	(nM)		(mg/kg)	(%)	
	(min.)					
	20	78,000	5.89/5.99	100	10	45
	21	87	7.71/7.21	NT	NT	NT
	22	460	6.60/6.46	NT	NT	NT
10	23	430	6.48/7.15	NT	NT	NT
	24	10	7.56/7.73	NT	NT	NT
	25	480	6.80/6.73	NT	NT	NT
	26	3.2	9.83/9.66	10	50	>180
	27	180	NT	NT	NT	NT
15	28	570	5.57/6.00	NT	NT	NT
	29	160	NT	NT	NT	NT
	30	22	7.73/7.88	30	50	>180
	31	14	NT	NT	NT	NT
	32	16	7.68/7.29	NT	NT	NT
20	33	630	6.73/6.36	NT	NT	NT
	34	640	5.34/5.69	NT	NT	NT
	35	41	7.25/7.47	NT	NT	NT
	36	1400	5.92/5.68	NT	NT	NT
	37	340	6.90/6.85	NT	NT	NT
25	38	10	7.82/8.36	NT	NT	NT
	39	10	7.88/7.84	NT	NT	NT
	40	83	7.94/7.61	NT	NT	NT
	41	3700	5.68/5.96	NT	NT	NT
	42	370	6.56/6.26	NT	NT	NT
30	43	19	8.97/8.61	NT	NT	NT
	44	16	8.23/7.70	NT	NT	NT
	45	4.4	8.41/8.24	NT	NT	NT
	46	110	6.80/6.64	NT	NT	NT

5	Test Compound Duration Example # (min.)	¹ Assay A IC ₅₀ (nM)	² Assay B pA ₂	Dose (mg/kg)	³ Assay C Inhibition (%)	
5	47	21	7.85/7.58	NT	NT	NT
	48	680	6.27/6.75	NT	NT	NT
	49	120	7.06/7.07	NT	NT	NT
	50	54	7.71/7.89	NT	NT	NT
	51	8.7	8.39/8.51	NT	NT	NT
10	52	100	8.14/8.12	NT	NT	NT
	53	65	7.56/7.83	NT	NT	NT
	54	3100	6.02	NT	NT	NT
	55	80	6.56/7.13	NT	NT	NT
	56	5.0	9.04/8.35	NT	NT	NT
15	57	2300	6.00	NT	NT	NT
	58	140	6.45/6.57	NT	NT	NT
	59	120	7.23/7.59	NT	NT	NT
	60	2200	6.40/6.03	NT	NT	NT
	61	110	7.29/7.70	NT	NT	NT
20	62	26	8.69/8.61	NT	NT	NT
	63	61	7.77/7.67	NT	NT	NT
	64	54	7.00/6.77	NT	NT	NT
	65	23	7.85/7.75	NT	NT	NT
	66	12	9.34/8.58	NT	NT	NT
25	67	3100	5.88/5.78	NT	NT	NT
	68	8.6	8.19/8.65	NT	NT	NT
	69	15	7.80/8.28	NT	NT	NT
	70	44	7.71/8.05	NT	NT	NT
	71	12,000	*	NT	NT	NT
30	72	83	6.11/6.10	NT	NT	NT
	73	790	7.65/7.46	NT	NT	NT

5	Test Compound Duration Example # (min.)	¹ Assay A IC ₅₀ (nM)	² Assay B pA ₂	Dose (mg/kg)	³ Assay C Inhibition (%)	
	74	6.5	8.56/8.39	NT	NT	NT
	75	570	6.00/5.45	NT	NT	NT
	76	5400	5.52/5.78	NT	NT	NT
10	77	15,000	5.77	NT	NT	NT
	78	480	6.41/6.35	NT	NT	NT

NT = NOT TESTED

15 *Antagonist activity not observed up to 10 μ M of test compound.

¹Assay A: Angiotensin II Binding Activity

²Assay B: In Vitro Vascular Smooth Muscle Response

20 ³Assays C/D: In Vivo Pressor Response (test compounds administered intraduodenally, except for

compounds of Examples #3, #26 and #30 which were given intragastrically).

Assay D: In Vivo Effects of Chronic Infusion of
Conjugate of the Invention

A conjugate of the invention as synthesized in
5 Example 79 was evaluated biologically by in vivo assays
to determine the ability of the conjugate to selectively
inhibit renal action and thereby control blood pressure.
This in vivo experiment was conducted to characterize
10 the effects of the Example 79 conjugate on spontaneously
hypertensive rats (SHR) by acute administration i.v. and
by chronic administration i.v. The Example 18 compound
or saline vehicle was infused continuously for four days
in SHR. Mean arterial pressure was measured (Gould Chart
15 Recorder, model 3800; Statham P23Db pressure transducer)
via an indwelling femoral artery catheter between
10:00a.m. and 2:00 P. M. each day. The Example 79
conjugate (10 mg/hr) or saline was infused via a jugular
vein catheter with a Harvard infusion pump. After
20 administration of the Example 79 conjugate, there was
observed a lowered mean arterial pressure as compared to
the saline vehicle control as reported in Table V and
also in Fig. 1. A test was conducted to determine
whether the Example 79 conjugate would antagonize non-
renal, vascular angiotensin II receptors. In this test
25 AII was administered by bolus injection (100 ng/kg) to
the SHR rats (described above) on the control day and on
days 1, 3 and 4 during conjugate infusion. No evidence
for systemic angiotensin II receptor antagonism was
observed, given the similar pressor responses to
30 injections of angiotensin II on the control day and days
1, 3 and 4 of infusion of the Example 79 conjugate (see
Figure 2). Tachycardia was observed on day 1 of the
conjugate infusion, but heart rate was returned to
control level during the next three days (see Figure 3).

TABLE V

Effect of Ex. #79 Conjugate on Mean
Arterial Pressure: Chronic Administration

5

Time (days): Control 1 2 3 4

Ex. #18 Conjugate (10

10 mg/hr)

MAP (mm Hg) :	183	159	155	154	166
(SD)	±5	±7	±4	±6	±9

350

TABLE VI

Effect of Ex. #18 Conjugate on
AI Pressor Response

5

Time (days): Control 1 3 4

Ex. #18 Conjugate (10 mg/hr)

10

	31	30	29	25
(SD)	± 4	± 6	± 5	± 3

TABLE VII

Effect of Ex. #79 Conjugate on Heart Rate

[illegible]

Also embraced within this invention is a class of pharmaceutical compositions comprising one or more conjugates which comprises a first component selected from angiotensin II antagonist compounds of Formula I linked to
5 a second component provided by an enzyme-cleavable moiety. Such pharmaceutical compositions further comprise one or more non-toxic, pharmaceutically acceptable carriers and/or diluents and/or adjuvants (collectively referred to herein as "carrier" materials) and, if desired, other active
10 ingredients. The conjugates of the present invention may be administered by any suitable route, preferably in the form of a pharmaceutical composition adapted to such a route, and in a dose effective for the treatment intended. Therapeutically effective doses of a conjugate of the
15 present invention required to prevent or arrest the progress of the medical condition are readily ascertained by one of ordinary skill in the art. The conjugates and composition may, for example, be administered intra-vascularly, intraperitoneally, subcutaneously, intra-
20 muscularly or topically.

For oral administration, the pharmaceutical composition may be in the form of, for example, a tablet, capsule, suspension or liquid. The pharmaceutical
25 composition is preferably made in the form of a dosage unit containing a particular amount of the conjugate. Examples of such dosage units are tablets or capsules. These may with advantage contain an amount of conjugate from about 1 to 250 mg, preferably from about 25 to 150 mg. A suitable
30 daily dose for a mammal may vary widely depending on the condition of the patient and other factors. However, a dose of from about 0.1 to 3000 mg/kg body weight, particularly from about 1 to 100 mg/kg body weight, may be appropriate.

35 The conjugate may also be administered by injection as a composition wherein, for example, saline, dextrose or water may be used as a suitable carrier. A suitable daily dose is from about 0.1 to 100 mg/kg body

weight injected per day in multiple doses depending on the disease being treated. A preferred daily dose would be from about 1 to 30 mg/kg body weight. Conjugates indicated for prophylactic therapy will preferably be administered in a daily dose generally in a range from about 0.1 mg to about 100 mg per kilogram of body weight per day. A more preferred dosage will be a range from about 1 mg to about 100 mg per kilogram of body weight. Most preferred is a dosage in a range from about 1 to about 50 mg per kilogram of body weight per day. A suitable dose can be administered, in multiple sub-doses per day. These sub-doses may be administered in unit dosage forms. Typically, a dose or sub-dose may contain from about 1 mg to about 100 mg of active compound per unit dosage form. A more preferred dosage will contain from about 2 mg to about 50 mg of active compound per unit dosage form. Most preferred is a dosage form containing from about 3 mg to about 25 mg of active compound per unit dose.

The dosage regimen for treating a disease condition with the conjugates and/or compositions of this invention is selected in accordance with a variety of factors, including the type, age, weight, sex and medical condition of the patient, the severity of the disease, the route of administration, and the particular conjugate employed, and thus may vary widely.

For therapeutic purposes, the conjugates of this invention are ordinarily combined with one or more adjuvants appropriate to the indicated route of administration. If administered per os, the conjugate may be admixed with lactose, sucrose, starch powder, cellulose esters of alkanolic acids, cellulose alkyl esters, talc, stearic acid, magnesium stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulfuric acids, gelatin, acacia gum, sodium alginate, polyvinylpyrrolidone, and/or polyvinyl alcohol, and then tableted or encapsulated for convenient administration. Such capsules or tablets

may contain a controlled-release formulation as may be provided in a dispersion of conjugate in hydroxypropylmethyl cellulose. Formulations for parenteral administration may be in the form of aqueous or non-aqueous isotonic sterile injection solutions or suspensions. These solutions and suspensions may be prepared from sterile powders or granules having one or more of the carriers or diluents mentioned for use in the formulations for oral administration. The conjugates may be dissolved in water, polyethylene glycol, propylene glycol, ethanol, corn oil, cottonseed oil, peanut oil, sesame oil, benzyl alcohol, sodium chloride, and/or various buffers. Other adjuvants and modes of administration are well and widely known in the pharmaceutical art.

15

Although this invention has been described with respect to specific embodiments, the details of these embodiments are not to be construed as limitations. Various equivalents, changes and modifications may be made without departing from the spirit and scope of this invention, and it is understood that such equivalent embodiments are part of this invention.

20


WHAT IS CLAIMED IS:

1. A conjugate comprising a residue of a biphenylalkyl 1H-substituted-1,2,4-triazole angiotensin II antagonist compound, said conjugate being renal selective.

2. Conjugate of Claim 1 comprising a first residue and a second residue, said first and second residues connected together by a cleavable bond, wherein said first residue is provided by said biphenylalkyl 1H-substituted-1,2,4-triazole angiotensin II antagonist compound, and wherein said second residue is capable of being cleaved from said first residue selectively in the kidney.


3. Conjugate of Claim 2 wherein said first and second residues are provided by precursor compounds wherein the precursor compound of one of said first and second residues has a reactable carboxylic acid moiety and the precursor of the other of said first and second residues has a reactable amino moiety or a moiety convertible to a reactable amino moiety, whereby a cleavable bond may be formed between said carboxylic acid moiety and said amino moiety.

4. Conjugate of Claim 3 wherein said angiotensin II antagonist compound is selected from a class of compounds defined by Formula I:



wherein m is a number selected from one to four, inclusive;

wherein each of R¹ through R¹¹ is independently selected from hydrido, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, formyl, alkoxy, aralkyl, aryl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkylcarbonylalkyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxycarbonylalkyl, aralkoxycarbonylalkyl, aralkylcarbonyloxyalkyl, mercaptocarbonyl, mercaptothiocarbonyl, mercaptoalkyl, alkoxycarbonyloxy, alkylthio, cycloalkylthio, alkylthiocarbonyl, alkylcarbonylthio, alkylthiocarbonyloxy, alkylthiocarbonylthio, alkylthiothiocarbonyl, alkylthiothiocarbonylthio, arylthio, arylthiocarbonyl, arylcarbonylthio, arylthiocarbonyloxy, arylthiocarbonylthio, arylthiothiocarbonyl, arylthiothiocarbonylthio, aralkylthio, aralkylthiocarbonyl, aralkylcarbonylthio, aralkylthiocarbonyloxy, aralkylthiocarbonylthio, alkylthiocarbonyl, aralkylthiocarbonylthio, mercapto, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl wherein each of said heteroaryl- and cyclohetero-containing groups has one or more ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R¹ through R¹¹ may be further independently selected from amino and amido radicals of the formula



wherein X is oxygen atom or sulfur atom;

- 5 wherein each n is a number independently selected from zero to six, inclusive;


- wherein each of R¹² through R²⁴ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino,
 10 monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl, and wherein R¹² and R¹³ taken together, R¹⁴ and R¹⁵ taken together, R¹⁶ and R¹⁷ taken together, R¹⁹ and R²⁰ taken together and R²¹ and R²² taken together may each form a heterocyclic group
 15 having five to seven ring members including the nitrogen atom of said amino or amido radical and which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms and which heterocyclic group may be saturated or partially
 20 unsaturated; wherein R¹² and R¹³ taken together, R¹⁴ and R¹⁵ taken together, R¹⁹ and R²⁰ taken together and R²¹ and R²² taken together may each form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino or amido radical and which aromatic
 25 heterocyclic group may further contain one or more hetero atoms as ring atoms selected from oxygen, nitrogen and sulfur atoms;

- and wherein each of R³ through R¹¹ may be further
 30 independently selected from hydroxy and from acidic moieties of the formula




wherein n is a number selected from zero through three, inclusive, and wherein A is an acidic group selected to
 5 contain at least one acidic hydrogen atom, and the amide, ester and salt derivatives of said acidic moieties; wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, aryl, aralkyl and heteroaryl having one or more
 10 ring atoms selected from oxygen, sulfur and nitrogen atoms;


and wherein any of the foregoing R^1 through R^{24} , Y and A groups having a substitutable position may be substituted with one or more groups selected from hydroxy, alkyl,
 15 alkenyl, alkynyl, aralkyl, hydroxyalkyl, trifluoromethyl, difluoroalkyl, oxo, alkoxy, aryloxy, aralkoxy, aralkylthio, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aroyl, cycloalkenyl, cyano, cyanoamino, nitro, alkylcarbonyloxy, alkoxy carbonyloxy, alkylcarbonyl, alkoxy carbonyl, carboxyl,
 20 mercapto, mercaptocarbonyl, alkylthio, arylthio, alkylthiocarbonyl, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms, and amino
 25 and amido radicals of the formula



wherein X is selected from oxygen atom and sulfur atom;
 30 wherein each of R^{25} , R^{26} , R^{27} , R^{28} and R^{29} is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl, DR^{30} and



wherein D is selected from oxygen atom and sulfur atom and R³⁰ is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl and aryl; wherein each of R²⁵,
 5 R²⁶, R²⁷, R²⁸, R²⁹, R³¹ and R³² is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, carboxyl, alkylsulfinyl, alkylsulfonyl, arylsulfinyl, arylsulfonyl, haloalkylsulfinyl,
 10 haloalkylsulfonyl, aralkyl and aryl, and wherein each of R²⁵, R²⁶, R²⁷, R²⁸, R²⁹, R³¹ and R³² is further independently selected from amino and amido radicals of the formula



wherein X is oxygen atom or sulfur atom;


wherein each of R³³, R³⁴, R³⁵, R³⁶, R³⁷ and R³⁸ is
 20 independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl, and wherein R²⁶ and R²⁷ taken together and R²⁸ and R²⁹ taken together may each
 25 form a heterocyclic group having five to seven ring members including the nitrogen atom of said amino or amido radical, which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms and which heterocyclic group may be
 30 saturated or partially unsaturated; wherein R²⁶ and R²⁷ taken together and R³¹ and R³² taken together may each form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino or amido radical and which aromatic heterocyclic group may further contain
 35 one or more hetero atoms as ring atoms selected from oxygen, nitrogen and sulfur atoms;

with the proviso that at least one of said R¹ through R²⁴,
Y and A substituents contains a terminal primary or
secondary amino moiety or a moiety convertible to a primary
5 or secondary amino moiety.

or a tautomer thereof or a pharmaceutically-acceptable salt
thereof.

10 5. Conjugate of Claim 4 wherein m is one;
wherein each of R¹ through R¹¹ is independently selected
from alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl,
cycloalkylalkyl, alkoxy, aralkyl, aryl, aroyl, aryloxy,
aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl,
15 alkoxy carbonyl, alkenyl, cycloalkenyl, alkynyl,
cycloalkynyl, cyano, nitro, carboxyl, carboxyalkyl,
alkylcarbonyloxy, alkylcarbonyloxyalkyl,
alkoxy carbonylalkyl, aralkoxy carbonylalkyl,
aralkylcarbonyloxyalkyl, mercaptocarbonyl,
20 mercaptothiocarbonyl, mercaptoalkyl, alkoxy carbonyloxy,
alkylthio, cycloalkylthio, alkylthiocarbonyl,
alkylcarbonylthio, alkylthiocarbonyloxy, alkylthio-
carbonylthio, alkylthiothiocarbonyl, alkylthiothio-
carbonylthio, arylthio, arylthiocarbonyl, arylcarbonylthio,
25 arylthiocarbonyloxy, arylthiocarbonylthio,
arylthiothiocarbonyl, arylthiothiocarbonylthio,
aralkylthio, aralkylthiocarbonyl, aralkylcarbonylthio,
aralkylthiocarbonyloxy, aralkylthiocarbonylthio,
aralkylthiocarbonyl, aralkylthiocarbonylthio, mercapto,
30 alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl,
aralkylsulfonyl, arylsulfinyl, arylsulfonyl, phthalimido,
phthalimidoalkyl, heteroaryl, heteroarylalkyl,
cycloheteroalkyl, cycloheteroalkylalkyl and
cycloheteroalkylcarbonylalkyl wherein each of said
35 heteroaryl- and cycloheteroalkyl-containing groups has
one or more hetero ring atoms selected from oxygen, sulfur

and nitrogen atoms, and wherein each of R¹ through R¹¹ may be further independently selected from amino and amido radicals of the formula



wherein X is selected from oxygen atom or sulfur atom;

wherein each n is a number independently selected from zero to six, inclusive;

10

wherein each of R¹² through R²⁴ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

15

and wherein each of R³ through R¹¹ may be further independently selected from hydroxy and from acidic moieties of the formula

20




wherein n is a number selected from zero through three, inclusive; wherein A is an acidic group selected from acids containing one or more atoms selected from oxygen, sulfur, phosphorus and nitrogen atoms, and wherein said acidic group is selected to contain at least one acidic hydrogen atom, and the amide, ester and salt derivatives of said acidic moieties; wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, aryl, aralkyl and

25


30

heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms;

- and wherein any of the foregoing R^1 through R^{24} , Y and A groups having a substitutable position may be substituted with one or more groups selected from alkyl, alkenyl, aralkyl, hydroxyalkyl, trifluoromethyl, difluoroalkyl, alkoxy, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, carboxyl, mercaptocarbonyl, alkylthio, alkylthiocarbonyl, and amino and amido radicals of the formula




- wherein X is selected from oxygen atom and sulfur atom; wherein each of R^{25} , R^{26} , R^{27} , R^{28} and R^{29} is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl, and DR^{30} and



- wherein D is selected from oxygen atom and sulfur atom, and R^{30} is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl and aryl; wherein each of R^{25} , R^{26} , R^{27} , R^{28} , R^{29} , R^{31} and R^{32} is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, alkanoyl, alkoxy carbonyl, carboxyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl, and wherein each of R^{25} , R^{26} , R^{27} , R^{28} , R^{29} , R^{31} and R^{32} is further independently selected from amino and amido radicals of the formula

363



wherein X is selected from oxygen atom or sulfur atom;

5 wherein each of R³³ through R³⁸ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl;

10

with the proviso that at least one of said R¹ through R²⁴, Y and A substituents contains a terminal primary or secondary amino moiety or a moiety convertible to a primary or secondary amino moiety;

15

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

6. Conjugate of Claim 5 wherein m is one;

20

wherein each of R¹ through R¹¹ is independently selected from alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, aralkyl, aryl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, alkenyl, cycloalkenyl, alkynyl,

25

cycloalkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, mercaptocarbonyl, alkoxy carbonyloxy, alkylcarbonyloxyalkyl, alkoxy carbonylalkyl, aralkoxy carbonylalkyl, aralkylcarbonyloxyalkyl, alkylthio, cycloalkylthio, arylthio, aralkylthio,


30

aralkylthiocarbonylthio, mercapto, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl

35

wherein each of said heteroaryl- and cycloheteroalkyl-containing groups has one or more hetero ring atoms

selected from oxygen, sulfur and nitrogen atoms, and wherein each of R^1 through R^{11} may be further independently selected from amino and amido radicals of the formula



wherein X is selected from oxygen atom or sulfur atom;

10 wherein each n is a number independently selected from zero to six, inclusive;


wherein each of R^{12} through R^{24} is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino,
 15 monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

and wherein each of R^3 through R^{11} may be further independently selected from hydroxy and from acidic
 20 moieties of the formula




wherein n is a number selected from zero through three,
 25 inclusive;

wherein A is selected from carboxylic acid and bioisosteres of carboxylic acid selected from



wherein each W is independently selected from oxygen atom, sulfur atom and NR⁴³; wherein each of R³⁹, R⁴⁰, R⁴¹, R⁴² and R⁴³ is independently selected from hydrido, alkyl, haloalkyl, haloalkylsulfonyl, haloalkylcarbonyl, cycloalkyl, cycloalkylalkyl, aryl and aralkyl; wherein each of R³⁹, R⁴⁰, R⁴¹ and R⁴² may be further independently selected from amino radical of the formula



10

wherein each of R⁴⁴ and R⁴⁵ is independently selected from hydrido, alkyl, cycloalkyl, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl, and wherein R⁴⁴ and R⁴⁵ taken together may form a heterocyclic group having five to seven ring members including the nitrogen atom of said amino radical, which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms and which heterocyclic group may be saturated or partially unsaturated; wherein R⁴⁴ and R⁴⁵ taken together may form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino radical and which aromatic heterocyclic group may further contain one or more hetero atoms as ring atoms selected from oxygen, nitrogen and sulfur atoms; wherein each of R⁴⁴ and R⁴⁵ may be further independently selected from hydroxy, alkoxy, alkylthio, aryloxy, arylthio, aralkylthio and aralkoxy; and the amide, ester and salt derivatives of said acidic groups;

30


wherein said bioisostere of carboxylic acid may be further selected from heterocyclic acidic groups consisting of heterocyclic rings of four to about nine ring members, which heterocyclic ring contains at least one hetero atom selected from oxygen, sulfur and nitrogen atoms, which heterocyclic ring may be saturated, fully unsaturated or

35


partially unsaturated, and which heterocyclic ring may be attached at a single position selected from R^3 through R^{11} or may be attached at any two adjacent positions selected from R^3 through R^{11} so as to form a fused-ring system with one of the phenyl rings of the biphenyl moiety of Formula I; and the amide, ester and salt derivatives of said heterocyclic acidic groups;

wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, aryl and aralkyl;

and wherein any of the foregoing R^1 through R^{24} , Y and A groups having a substitutable position may be substituted by one or more groups selected from alkyl, difluoroalkyl, alkenyl, aralkyl, hydroxyalkyl, trifluoromethyl, alkoxy, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, carboxyl, mercaptocarbonyl, alkylthio, alkylthiocarbonyl, and amino and amido radicals of the formula



wherein X is selected from oxygen atom and sulfur atom; wherein each of R^{25} , R^{26} , R^{27} , R^{28} and R^{29} is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl and DR^{30} and



30


wherein D is selected from oxygen atom and sulfur atom, wherein R^{30} is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl and aryl;

wherein each of R²⁵, R²⁶, R²⁷, R²⁸, R²⁹, R³¹ and R³² is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, alkanoyl, alkoxycarbonyl, carboxyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl;

with the proviso that at least one of said R¹ through R²⁴, Y and A substituents contains a terminal primary or secondary amino moiety or a moiety convertible to a primary or secondary amino moiety;

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

7. Conjugate of Claim 6 wherein m is one; wherein each of R¹ and R² is independently selected from alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, aralkyl, aryl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxycarbonylalkyl, aralkoxycarbonylalkyl, aralkylcarbonyloxyalkyl, mercaptoalkyl, alkoxyalkyl, alkylthio, cycloalkylthio, arylthio, aralkylthio, mercapto, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl wherein each of said heteroaryl- and cycloheteroalkyl-containing groups has one or more hetero ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R¹ through R¹¹ may be further independently selected from amino and amido radicals of the formula



wherein X is selected from oxygen atom and sulfur atom;

5

wherein each n is a number independently selected from zero to six, inclusive;


10 wherein each of R¹² through R²⁴ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

15 wherein each of R³ through R¹¹ is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, aralkyl, aryl, aroyl, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl, alkylthio, 20 aralkylthio, mercapto, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl and heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms;


25 and wherein each of R³ through R¹¹ may be an acidic moiety further independently selected from acidic moieties of the formula



30 wherein n is a number selected from zero through three, inclusive; wherein A is selected from carboxylic acid and bioisosteres of carboxylic acid selected from



wherein each W is independently selected from oxygen atom,
 5 sulfur atom and NR^{43} ; wherein each of R^{39} , R^{42} and R^{43} is
 independently selected from hydrido, alkyl, haloalkyl,
 haloalkylsulfonyl, haloalkylcarbonyl, cycloalkyl,
 cycloalkylalkyl, aryl and aralkyl; wherein each of R^{39} and
 10 R^{42} may be further independently selected from amino
 radical of the formula



wherein each of R^{44} and R^{45} is independently selected from
 15 hydrido, alkyl, cycloalkyl, hydroxyalkyl, haloalkyl,
 cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl, and wherein
 R^{44} and R^{45} taken together may form a heterocyclic group
 having five to seven ring members including the nitrogen
 atom of said amino radical, which heterocyclic group may
 20 further contain one or more hetero atoms as ring members
 selected from oxygen, nitrogen and sulfur atoms, and which
 heterocyclic group may be saturated or partially
 unsaturated; wherein R^{44} and R^{45} taken together may form an
 aromatic heterocyclic group having five ring members
 25 including the nitrogen atom of said amino radical and which
 aromatic heterocyclic group may further contain one or more
 hetero atoms as ring atoms selected from oxygen, nitrogen
 and sulfur atoms; and the amide, ester and salt derivatives
 of said acidic groups; wherein said bioisostere of
 30 carboxylic acid may be further selected from heterocyclic
 acidic groups consisting of heterocyclic rings of four to
 about nine ring members, which ring contains at least one
 hetero atom, selected from oxygen, sulfur and nitrogen
 atoms, which heterocyclic ring may be saturated, fully
 35 unsaturated or partially unsaturated, and which

heterocyclic ring may be attached at a single position selected from R^3 through R^{11} or may be attached at any two adjacent positions selected from R^3 through R^{11} so as to form a fused-ring system with one of the phenyl rings of Formula I; and the amide, ester and salt derivatives of said heterocyclic acidic groups;

wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, aryl and aralkyl;


wherein each of R^1 through R^{11} , Y and A independently may be substituted at any substitutable position with one or more groups selected from alkyl, cycloalkyl, cycloalkylalkyl, hydroxy, oxo, trifluoromethyl, difluoroalkyl, alkoxycarbonyl, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, aryl, aralkyl, alkoxy, aryloxy and aralkoxy;

with the proviso that at least one of said R^1 through R^{24} , Y and A substituents contains a terminal primary or secondary amino moiety or a moiety convertible to a primary or secondary amino moiety;

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

8. Conjugate of Claim 7 wherein m is one; wherein each of R^1 and R^2 is independently selected from alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, aralkyl, aryl, benzoyl, phenoxy, phenoxyalkyl, phenalkyloxy, phenylthio, phenalkylthio, aralkoxy, alkoxylalkyl, alkylcarbonyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, carboxylalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxycarbonylalkyl, aralkoxycarbonylalkyl, aralkylcarbonyloxyalkyl, mercaptocarbonyl, mercaptoalkyl, alkoxycarbonyloxy, alkylthio, cycloalkylthio, phthalimido,

phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl wherein each of said heteroaryl- and cycloheteroalkyl-containing groups has one
 5 or more hetero ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R¹ through R¹¹ may be further independently selected from amino and amido radicals of the formula



wherein X is selected from oxygen atom and sulfur atom;

wherein each n is a number independently selected from zero
 15 to six, inclusive;


wherein each of R¹² through R²⁴ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, hydroxyalkyl, alkoxyalkyl, phenalkyl and phenyl;

20 wherein each of R³ through R¹¹ is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, phenalkyl, phenyl, benzoyl, phenoxy, phenalkyloxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, alkenyl, cyano, nitro,
 25 carboxyl, alkylthio, mercapto and heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms;

30 and wherein each of R³ through R¹¹ may be an acidic moiety further independently selected from acidic moieties of the formula




wherein n is a number selected from zero through two,
inclusive; wherein A is selected from carboxylic acid and
5 bioisosteres of carboxylic acid selected from



wherein each W is independently selected from oxygen atom,
10 sulfur atom and NR^{43} ; wherein each of R^{39} , R^{42} and R^{43} is
independently selected from hydrido, alkyl, haloalkyl,
haloalkylsulfonyl, haloalkylcarbonyl, cycloalkyl, phenyl
and benzyl; wherein each of R^{39} and R^{42} may be further
independently selected from amino radical of the formula

15



wherein each of R^{44} and R^{45} is independently selected from
hydrido, alkyl, cycloalkyl, hydroxyalkyl, haloalkyl,
20 alkoxyalkyl, benzyl and phenyl; and the amide, ester and
salt derivatives of said acidic groups;

wherein said bioisostere of carboxylic acid may be further
selected from heterocyclic acidic groups consisting of
25 heterocyclic rings of four to about nine ring members,
which ring contains at least one hetero atom, selected from
oxygen, sulfur and nitrogen atoms, which heterocyclic ring
may be saturated, fully unsaturated or partially
unsaturated, and which heterocyclic ring may be attached at
30 a single position selected from R^3 through R^{11} or may be
attached at any two adjacent positions selected from R^3
through R^{11} so as to form a fused-ring system with one of
the phenyl rings of the biphenyl moiety of Formula I; and
the amide, ester and salt derivatives of said heterocyclic
35 acidic groups;

wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, phenyl, phenalkyl and aralkyl;

5

wherein each of R¹ through R¹¹, Y and A and independently may be substituted at any substitutable position with one or more groups selected from alkyl, cycloalkyl, cycloalkylalkyl, hydroxy, oxo, trifluoromethyl, difluoroalkyl, alkoxy carbonyl, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, aryl, aralkyl, alkoxy, aryloxy and aralkoxy;

10

with the proviso that at least one of said R¹ through R²⁴, Y and A substituents contains a terminal primary or secondary amino moiety or a moiety convertible to a primary or secondary amino moiety;

15

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.


20

9. Conjugate of Claim 8 wherein m is one; wherein each of R¹ and R² is independently selected from alkyl, aminoalkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, aralkyl, aryl, benzoyl, phenoxy, phenoxyalkyl, phenalkyloxy, phenylthio, phenalkylthio, aralkoxy, alkoxyalkyl, acetyl, alkoxy carbonyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, mercaptoalkyl, mercaptocarbonyl, alkoxy carbonyloxy, alkylcarbonyloxyalkyl, alkoxy carbonylalkyl, aralkoxy carbonylalkyl, aralkylcarbonyloxyalkyl, phthalimido, phthalimidoalkyl, imidazoalkyl, tetrazole, tetrazolealkyl, alkylthio, cycloalkylthio, and amino and amido radicals of the formula

25

30

35



wherein X is selected from oxygen atom and sulfur atom;

5


wherein each n is a number independently selected from zero to six, inclusive;

10 wherein each of R¹² through R²⁴ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, hydroxyalkyl, alkoxyalkyl, phenalkyl and phenyl;


15 wherein each of R³ through R¹¹ is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, phenalkyl, phenyl, benzoyl, phenoxy, phenalkyloxy, alkoxyalkyl, acetyl, alkoxy carbonyl, alkenyl, cyano, nitro, carboxyl, alkylthio and mercapto;

20 and wherein each of R³ through R¹¹ may be an acidic moiety further independently selected from acidic moieties consisting of CO₂H, CO₂CH₃, SH, CH₂SH, C₂H₄SH, PO₃H₂, NHSO₂CF₃, NHSO₂C₆F₅, SO₃H, CONHNH₂, CONHNHSO₂CF₃, CONHOCH₃, CONHOC₂H₅, CONHCF₃, OH, CH₂OH, C₂H₄OH, OPO₃H₂, OSO₃H,

25





- wherein each of R⁴⁶, R⁴⁷ and R⁴⁸ is independently selected from H, Cl, CN, NO₂, CF₃, C₂F₅, C₃F₇, CHF₂, CH₂F, CO₂CH₃, CO₂C₂H₅, SO₂CH₃, SO₂CF₃ and SO₂C₆F₅; wherein Z is selected from O, S, NR⁴⁹ and CH₂; wherein R⁴⁹ is selected from hydrido, CH₃ and CH₂C₆H₅; and wherein said acidic moiety may be a heterocyclic acidic group attached at any two adjacent positions of R³ through R¹¹ so as to form a fused ring system so as to include one of the phenyl rings of the biphenyl moiety of Formula I, said biphenyl fused ring system selected from



and the esters, amides and salts of said acidic moieties;



- 5 with the proviso that at least one of said R^1 through R^{24} substituents contains a terminal primary or secondary amino moiety or a moiety convertible to a primary or secondary amino moiety;
- 10 or a tautomer thereof or a pharmaceutically-acceptable salt thereof.



10. Conjugate of Claim 9 wherein m is one;
 wherein each of R^1 and R^2 is independently selected from
 15 amino, aminomethyl, aminoethyl, aminopropyl, CH_2OH , CH_2OCOCH_3 , CH_2Cl , Cl , CH_2OCH_3 , $CH_2OCH(CH_3)_2$, I , CHO ,



- 20 $-CH_2NHCO_2CH_2(CH_3)_2$, $-CH_2NHCO_2C_4H_9$, CH_2NHCO_2 -adamantyl,

- CH₂NHCO₂-(1-naphthyl), -CH₂NHCONHCH₃, -CH₂NHCONHC₂H₅,
 -CH₂NHCONHC₃H₇, -CH₂NHCONHC₄H₉, -CH₂NHCONHCH(CH₃)₂,
 -CH₂NHCONH(1-naphthyl), -CH₂NHCONH(1-adamantyl), CO₂H,
 -CH₂CH₂-CO-N $\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array}$ O, -CH₂CH₂CO-N $\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array}$, -CH₂CH₂CH₂CO₂H,
 5 -CH₂CH₂F, -CH₂OCONHCH₃, -CH₂OCSNHCH₃, -CH₂NHCSOC₃H₇,

- CH₂CH₂CH₂F, -CH₂ONO₂, , -CH₂SH, -CH₂O-,
 H, Cl, NO₂, CF₃, CH₂OH, Br, F, I, methyl, ethyl, n-propyl,
 isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-
 pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl,
 10 cyclohexyl, cyclohexylmethyl, 1-oxoethyl, 1-oxopropyl, 1-
 oxobutyl, 1-oxopentyl, 1,1-dimethoxypropyl, 1,1-
 dimethoxybutyl, 1,1-dimethoxypentyl, hydroxyalkyl, halo,
 difluoromethyl, 1,1-difluoroethyl, 1,1-difluoropropyl, 1,1-
 difluorobutyl and 1,1-difluoropentyl; wherein each of R³
 15 through R¹¹ is hydrido, with the proviso that at least one
 of R⁵, R⁶, R⁸ and R⁹ is an acidic group selected from CO₂H,
 SH, PO₃H₂, SO₃H, CONHNH₂, CONHNHSO₂CF₃, OH,



and




20

wherein each of R⁴⁶ and R⁴⁷ is independently selected from
 Cl, CN, NO₂, CF₃, CO₂CH₃ and SO₂CF₃;

25 with the proviso that at least one of said R¹ through R¹¹
 substituents contains a terminal primary or secondary amino
 moiety or a moiety convertible to a primary or secondary
 amino moiety;

30 or a tautomer thereof or a pharmaceutically-acceptable salt
 thereof.

11. Conjugate of Claim 10 wherein m is one;
 wherein R¹ is selected from amino, aminomethyl, aminoethyl,
 aminopropyl, methyl, ethyl, n-propyl, isopropyl, n-butyl,
 sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl,
 5 neopentyl, phenyl, benzyl, phenethyl, cyclohexyl,
 cyclohexylmethyl, 1-oxoethyl, 1-oxopropyl, 1-oxobutyl, 1-
 oxopentyl, 1,1-dimethoxypropyl, 1,1-dimethoxybutyl, 1,1-
 dimethoxypentyl, hydroxyalkyl, halo, difluoromethyl, 1,1-
 difluoroethyl, 1,1-difluoropropyl, 1,1-difluorobutyl and
 10 1,1-difluoropentyl; wherein R² is selected from ethyl, n-
 propyl, isopropyl, n-butyl, sec-butyl, isobutyl, 4-
 methylbutyl, tert-butyl, n-pentyl and neopentyl; wherein
 each of R³, R⁴, R⁶, R⁷, R⁸, R¹⁰, and R¹¹ is hydrido; wherein
 one of R⁵ and R⁹ is hydrido and the other of R⁵ and R⁹ is
 15 an acidic group selected from CO₂H, SH, PO₃H₂, SO₃H,
 CONHNH₂, CONHNHSO₂CF₃, OH,



- 20 wherein each of R⁴⁶ and R⁴⁷ is independently selected from
 Cl, CN, NO₂, CF₃, CO₂CH₃ and SO₂CF₃;


- with the proviso that at least one of said R¹ through R¹¹
 substituents contains a terminal primary or secondary amino
 25 moiety or a moiety convertible to a primary or secondary
 amino moiety;

or a tautomer thereof or a pharmaceutically-acceptable salt
 thereof.

30

12. Conjugate of Claim 3 wherein said second
 residue forms a kidney-enzyme-cleavable amide bond with the
 residue of said angiotensin II antagonist compound.

13. Conjugate of Claim 3 wherein said second residue is preferably selected from a class of compounds of Formula II:



5

wherein each of R^{50} and R^{51} may be independently selected from hydrido, alkylcarbonyl, alkoxy carbonyl, alkoxyalkyl, hydroxyalkyl and haloalkyl; and wherein G is selected from hydroxyl, halo, mercapto, $-\text{OR}^{52}$, $-\text{SR}^{53}$ and $-\text{NR}^{54}$ with each of R^{52} , R^{53} and R^{54} independently selected from alkyl; and wherein R^{54} may be further selected from hydrido; with the proviso that said Formula II compound is selected such that formation of the cleavable amide bond occurs at carbonyl moiety attached at the gamma-position carbon of said Formula II compound.

10

15

14. Conjugate of Claim 13 wherein each G substituent is hydroxy.

20


15. Conjugate of Claim 14 wherein each G substituent is hydroxy; wherein R^{50} is hydrido; and wherein R^{51} is selected from

25

$-\text{CR}^{55}$ wherein R^{55} is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, iso-butyl, tert-butyl, n-pentyl, neopentyl, n-hexyl and chloromethyl.

30

16. Conjugate of Claim 15 wherein said second residue is




17. Conjugate of Claim 3 wherein said first residue is a biphenylalkyl 1H-substituted-1,2,4-triazole
 5 angiotensin II antagonist compound containing a terminal primary or secondary amino moiety selected from amino and linear or branched aminoalkyl moieties containing linear or branched alkyl groups selected from aminomethyl, aminoethyl, aminopropyl, aminoisopropyl, aminobutyl,
 10 aminosecbutyl, aminoisobutyl, aminotertbutyl, aminopentyl, aminoisopentyl and aminoneopentyl.

18. Conjugate of Claim 3 wherein said first residue is provided by a biphenylalkyl 1H-substituted-
 15 1,2,4-triazole angiotensin II antagonist compound containing a moiety convertible to a primary or secondary amino terminal moiety.

19. Conjugate of Claim 18 wherein said moiety
 20 convertible to an amino terminal moiety is a carboxylic acid group reactable with an amino moiety of a diamino-terminated linker group to provide a terminal amino moiety which may then be further reacted with a carboxylic acid moiety of a compound providing said second residue so as to
 25 form a hydrolyzable amide bond.

20. Conjugate of Claim 19 wherein said diamino-terminated linker group is a divalent radical of Formula III:

30




(III)


wherein each of R²⁰⁰ and R²⁰¹ may be independently selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, alkoxyalkyl, hydroxyalkyl, aralkyl, aryl, haloalkyl, amino, monoalkylamino, dialkylamino, cyanoamino, carboxyalkyl, alkylsulfinyl, alkylsulfonyl, arylsulfinyl and arylsulfonyl; and wherein n is zero or a number selected from three through seven, inclusive.

21. Conjugate of Claim 20 wherein each of R²⁰⁰ and R²⁰¹ is hydrido.

22. Conjugate of Claim 19 wherein said diamino-terminated linker group is a divalent radical of Formula IV:




wherein each of Q and T is one or more groups independently selected from



wherein each of R²⁰² through R²⁰⁵ is independently selected from hydrido, hydroxy, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl, alkoxy, aralkoxy, aryloxy, alkoxyalkyl, haloalkyl, hydroxyalkyl, halo, cyano, amino, monoalkylamino, dialkylamino, carboxy, carboxyalkyl, alkanoyl, alkenyl, cycloalkenyl and alkynyl.

23. Conjugate of Claim 22 wherein said diamino-terminated linker group is a divalent radical of Formula V:




wherein each of R^{202} and R^{203} is independently selected from hydrido, hydroxy, alkyl, phenalkyl, phenyl, alkoxy, benzyloxy, phenoxy, alkoxyalkyl, hydroxyalkyl, halo, amino, monoalkylamino, dialkylamino, carboxy, carboxyalkyl and alkanoyl; and wherein each of p and q is a number independently selected from one through six, inclusive; with the proviso that when each of R^{202} and R^{203} is selected from halo, hydroxy, amino, monoalkylamino and dialkylamino, then the carbon to which R^{202} or R^{203} is attached in Formula V is not adjacent to a nitrogen atom of Formula V.

24. Conjugate of Claim 23 wherein each of R^{202} and R^{203} is independently selected from hydrido, hydroxy, alkyl, alkoxy, amino, monoalkylamino, carboxy, carboxyalkyl and alkanoyl; and wherein each of p and q is a number independently selected from two through four, inclusive.

25. Conjugate of Claim 24 wherein each of R^{202} and R^{203} is independently selected from hydrido, amino, monoalkylamino and carboxyl; and wherein each of p and q is independently selected from the numbers two and three.

26. Conjugate of Claim 25 wherein each of R^{202} and R^{203} is hydrido; and wherein each of p and q is two.

27. Conjugate of Claim 19 wherein said diamino-terminated linker group is a divalent radical of Formula VI:



5

wherein each of R^{214} through R^{217} is independently selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, hydroxyalkyl, alkoxyalkyl, aralkyl, aryl, haloalkyl, amino, monoalkylamino, dialkylamino, cyanoamino, carboxyalkyl, alkylsulfinyl, alkylsulfonyl, arylsulfinyl and arylsulfonyl; and wherein p is a number selected from one through six inclusive.

15

28. Conjugate of Claim 27 wherein each of R^{214} and R^{215} is hydrido; wherein each of R^{216} and R^{217} is independently selected from hydrido, alkyl, phenalkyl, phenyl, alkoxyalkyl, hydroxyalkyl, haloalkyl and carboxyalkyl; and wherein p is two or three.

20

29. Conjugate of Claim 28 wherein each of R^{214} and R^{215} is hydrido; wherein each of R^{216} and R^{217} is independently selected from hydrido and alkyl; and wherein p is two.

25

30. Conjugate of Claim 29 wherein each of R^{214} , R^{215} , R^{216} and R^{217} is hydrido; and wherein p is two.

30

31. Conjugate of Claim 11 wherein said angiotensin II antagonist compound is selected from the group consisting of methyl 4'-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylate; 4'-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;

35

- 4'-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid, hydrazide;
4'-[(5-butyl-3-chloro-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 5 4'-[(3-butyl-5-chloro-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-propyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 10 4'-[(5-butyl-3-isopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-secbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-isobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 15 4'-[(5-butyl-3-tertbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-pentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-isopentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 20 4'-[(5-butyl-3-cyclohexyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-cyclohexylmethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 25 4'-[[5-butyl-3-(2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-cyclohexanoyl-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 30 4'-[[5-butyl-3-(1-oxo-2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-phenyl-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 35 4'-[(5-butyl-3-phenylmethyl-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-butyl-3-(2-phenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-benzoyl-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[[5-butyl-3-(1-oxo-2-phenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-butyl-3-(1,1-dimethoxypropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
5 4'-[[5-butyl-3-(1,1-dimethoxybutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-butyl-3-(1-oxopropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-butyl-3-(1-oxobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
10 4'-[[5-butyl-3-(1-oxopentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-butyl-3-(1,1-difluoroethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
15 4'-[[5-butyl-3-(1,1-difluoropropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-butyl-3-(1,1-difluorobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[3-butyl-5-(1,1-difluorobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
20 4'-[[5-butyl-3-(1,1-difluoropentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(3,5-dipropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
25 4'-[(3,5-isopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(3,5-disecbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(3,5-diisobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
30 4'-[(3,5-ditertbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(3,5-dipentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
35 4'-[(3,5-diisopentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
5-[4'-[(5-butyl-3-amino-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;

- 5-[4'-[(5-butyl-3-aminomethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-butyl-3-aminoethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5 5-[4'-[(5-butyl-3-aminopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-butyl-3-aminobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminophenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
10 5-[4'-[[3-butyl-5-(4-aminophenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminophenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
15 5-[4'-[[3-butyl-5-(4-aminomethylphenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminomethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminoethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
20 5-[4'-[[3-butyl-5-(4-aminocyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminocyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
25 5-[4'-[[3-butyl-5-(4-aminocyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminomethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminomethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
30 5-[4'-[[3-butyl-5-(4-aminoethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-butyl-3-carboxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
35 5-[4'-[(5-butyl-3-carboxymethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-butyl-3-carboxyethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;

- 5-[4'-[(5-butyl-3-carboxypropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
 5-[4'-[(5-butyl-3-carboxybutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
 5-[4'-[[3-butyl-5-(4-carboxyphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
 5-[4'-[[3-butyl-5-(4-carboxyphenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
 5-[4'-[[3-butyl-5-(4-carboxyphenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
 5-[4'-[[3-butyl-5-(4-carboxymethylphenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
 5-[4'-[[3-butyl-5-(4-carboxymethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
 5-[4'-[[3-butyl-5-(4-carboxyethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
 5-[4'-[[3-butyl-5-(4-carboxycyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
 5-[4'-[[3-butyl-5-(4-carboxycyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
 5-[4'-[[3-butyl-5-(4-carboxycyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
 5-[4'-[[3-butyl-5-(4-carboxymethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
 5-[4'-[[3-butyl-5-(4-carboxymethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole; and
 5-[4'-[[3-butyl-5-(4-carboxyethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole.

30

32. Conjugate of Claim 21 which is N-acetylglutamic acid, 5-[[4'-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]]carbonylhydrazide.

35

33. Conjugate of Claim 17 which is N²-acetyl-N-[[5-butyl-1-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl)methyl]-1H-1,2,4-triazol-3-yl)methyl]-L-glutamine.

34. Conjugate of Claim 21 which is N-acetyl-L-glutamic acid, 5-[5-butyl-1-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl-1H-1,2,4-triazol-3-yl]acetylhydrazide.

5


35. A pharmaceutical composition comprising one or more pharmaceutically-acceptable carriers or diluents and a therapeutically-effective amount of a renal-selective conjugate, said renal selective conjugate comprising a
10 residue of a biphenylalkyl 1H-substituted-1,2,4-triazole angiotensin II antagonist compound.

36. The composition of Claim 35 comprising a first residue and a second residue, said first and second
15 residues connected together by a cleavable bond, wherein said first residue is provided by said biphenylalkyl 1H-substituted-1,2,4-triazole angiotensin II antagonist compound, and wherein said second residue is capable of being cleaved from said first residue selectivity in the
20 kidney.

37. The composition of Claim 36 wherein said first and second residues are provided by precursor compounds wherein the precursor compound of one of said
25 first and second residues has a reactable carboxylic acid moiety and the precursor of the other of said first and second residues has a reactable amino moiety or a moiety convertible to a reactable amino moiety, whereby a cleavable bond may be formed between said carboxylic acid
30 moiety and said amino moiety.

38. The composition of Claim 37 wherein said angiotensin II antagonist compound is selected from a class of compounds defined by Formula I:


35



wherein m is a number selected from one to four, inclusive;

- 5 wherein each of R¹ through R¹¹ is independently selected from hydrido, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, formyl, alkoxy, aralkyl, aryl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, 10 alkylcarbonyl, alkylcarbonylalkyl, alkoxy carbonyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxy carbonylalkyl, aralkoxy carbonylalkyl, aralkylcarbonyloxyalkyl, mercaptocarbonyl, 15 mercaptothiocarbonyl, mercaptoalkyl, alkoxy carbonyloxy, alkylthio, cycloalkylthio, alkylthiocarbonyl, alkylcarbonylthio, alkylthiocarbonyloxy, alkylthiocarbonylthio, alkylthiothiocarbonyl, alkylthiothiocarbonylthio, arylthio, arylthiocarbonyl, 20 arylcarbonylthio, arylthiocarbonyloxy, arylthiocarbonylthio, arylthiothiocarbonyl, arylthiothiocarbonylthio, aralkylthio, aralkylthiocarbonyl, aralkylcarbonylthio, aralkylthiocarbonyloxy, aralkylthiocarbonylthio, alkylthiocarbonyl, 25 aralkylthiocarbonylthio, mercapto, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl 30 wherein each of said heteroaryl- and cyclohetero-containing groups has one or more ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R¹ through

R¹¹ may be further independently selected from amino and amido radicals of the formula



5

wherein X is oxygen atom or sulfur atom;

wherein each n is a number independently selected from zero to six, inclusive;

10

wherein each of R¹² through R²⁴ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl, and wherein R¹² and R¹³ taken together, R¹⁴ and R¹⁵ taken together, R¹⁶ and R¹⁷ taken together, R¹⁹ and R²⁰ taken together and R²¹ and R²² taken together may each form a heterocyclic group having five to seven ring members including the nitrogen atom of said amino or amido radical and which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms and which heterocyclic group may be saturated or partially unsaturated; wherein R¹² and R¹³ taken together, R¹⁴ and R¹⁵ taken together, R¹⁹ and R²⁰ taken together and R²¹ and R²² taken together may each form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino or amido radical and which aromatic heterocyclic group may further contain one or more hetero atoms as ring atoms selected from oxygen, nitrogen and sulfur atoms;

30

and wherein each of R^3 through R^{11} may be further independently selected from hydroxy and from acidic moieties of the formula

5




wherein n is a number selected from zero through three, inclusive, and wherein A is an acidic group selected to contain at least one acidic hydrogen atom, and the amide, ester and salt derivatives of said acidic moieties; wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, aryl, aralkyl and heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms;

15


and wherein any of the foregoing R^1 through R^{24} , Y and A groups having a substitutable position may be substituted with one or more groups selected from hydroxy, alkyl, alkenyl, alkynyl, aralkyl, hydroxyalkyl, trifluoromethyl, difluoroalkyl, oxo, alkoxy, aryloxy, aralkoxy, aralkylthio, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aroyl, cycloalkenyl, cyano, cyanoamino, nitro, alkylcarbonyloxy, alkoxy carbonyloxy, alkylcarbonyl, alkoxy carbonyl, carboxyl, mercapto, mercaptocarbonyl, alkylthio, arylthio, alkylthiocarbonyl, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms, and amino and amido radicals of the formula

30




wherein X is selected from oxygen atom and sulfur atom; wherein each of R^{25} , R^{26} , R^{27} , R^{28} and R^{29} is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl, DR^{30} and

35



- wherein D is selected from oxygen atom and sulfur atom and
- 5 R³⁰ is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl and aryl; wherein each of R²⁵, R²⁶, R²⁷, R²⁸, R²⁹, R³¹ and R³² is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, alkylcarbonyl, alkoxy
- 10 alkoxy, carbonyl, carboxyl, alkylsulfinyl, alkylsulfonyl, arylsulfinyl, arylsulfonyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl, and wherein each of R²⁵, R²⁶, R²⁷, R²⁸, R²⁹, R³¹ and R³² is further independently selected from amino and amido radicals of the
- 15 formula



- wherein X is oxygen atom or sulfur atom;
- 20 wherein each of R³³, R³⁴, R³⁵, R³⁶, R³⁷ and R³⁸ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl, and wherein R²⁶ and R²⁷ taken together and R²⁸ and R²⁹ taken together may each form a heterocyclic group having five to seven ring members including the nitrogen atom of said amino or amido radical, which heterocyclic group may further contain one or more
- 25 hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms and which heterocyclic group may be saturated or partially unsaturated; wherein R²⁶ and R²⁷ taken together and R³¹ and R³² taken together may each form an aromatic heterocyclic group having five ring members
- 30

including the nitrogen atom of said amino or amido radical and which aromatic heterocyclic group may further contain one or more hetero atoms as ring atoms selected from oxygen, nitrogen and sulfur atoms;

5


with the proviso that at least one of said R¹ through R²⁴, Y and A substituents contains a terminal primary or secondary amino moiety or a moiety convertible to a primary or secondary amino moiety.

10

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

39. The composition of Claim 38 wherein m is one; wherein each of R¹ through R¹¹ is independently selected from alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, aralkyl, aryl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxycarbonylalkyl, aralkoxycarbonylalkyl, aralkylcarbonyloxyalkyl, mercaptocarbonyl, mercaptothiocarbonyl, mercaptoalkyl, alkoxycarbonyloxy, alkylthio, cycloalkylthio, alkylthiocarbonyl, alkylcarbonylthio, alkylthiocarbonyloxy, alkylthiocarbonylthio, alkylthiothiocarbonyl, alkylthiothiocarbonylthio, arylthio, arylthiocarbonyl, arylcarbonylthio, arylthiocarbonyloxy, arylthiocarbonylthio, arylthiothiocarbonyl, arylthiothiocarbonylthio, aralkylthio, aralkylthiocarbonyl, aralkylcarbonylthio, aralkylthiocarbonyloxy, aralkylthiocarbonylthio, aralkylthiocarbonyl, aralkylthiocarbonylthio, mercapto, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and

- cycloheteroalkylcarbonylalkyl wherein each of said heteroaryl- and cycloheteroalkyl-containing groups has one or more hetero ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R¹ through R¹¹ may be further independently selected from amino and amido radicals of the formula



- wherein X is selected from oxygen atom or sulfur atom;
- wherein each n is a number independently selected from zero to six, inclusive;
- wherein each of R¹² through R²⁴ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;
- and wherein each of R³ through R¹¹ may be further independently selected from hydroxy and from acidic moieties of the formula




- wherein n is a number selected from zero through three, inclusive; wherein A is an acidic group selected from acids containing one or more atoms selected from oxygen, sulfur, phosphorus and nitrogen atoms, and wherein said acidic group is selected to contain at least one acidic hydrogen atom, and the amide, ester and salt derivatives of said acidic moieties; wherein Y is a spacer group independently

selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, aryl, aralkyl and heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms;

5


and wherein any of the foregoing R^1 through R^{24} , Y and A groups having a substitutable position may be substituted with one or more groups selected from alkyl, alkenyl, aralkyl, hydroxyalkyl, trifluoromethyl, difluoroalkyl, alkoxy, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, carboxyl, mercaptocarbonyl, alkylthio, alkylthiocarbonyl, and amino and amido radicals of the formula

15



wherein X is selected from oxygen atom and sulfur atom; wherein each of R^{25} , R^{26} , R^{27} , R^{28} and R^{29} is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl, and DR^{30} and


20



wherein D is selected from oxygen atom and sulfur atom, and R^{30} is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl and aryl; wherein each of R^{25} , R^{26} , R^{27} , R^{28} , R^{29} , R^{31} and R^{32} is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, alkanoyl, alkoxy carbonyl, carboxyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl, and wherein each of R^{25} , R^{26} , R^{27} , R^{28} , R^{29} , R^{31} and R^{32} is further independently selected from amino and amido radicals of the formula

25

30



wherein X is selected from oxygen atom or sulfur atom;

5

wherein each of R³³ through R³⁸ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl;

10

with the proviso that at least one of said R¹ through R²⁴, Y and A substituents contains a terminal primary or secondary amino moiety or a moiety convertible to a primary or secondary amino moiety;

15

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

20

40. The composition of Claim 39 wherein m is one; wherein each of R¹ through R¹¹ is independently selected from alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, aralkyl, aryl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, mercaptocarbonyl, alkoxy carbonyloxy, alkylcarbonyloxyalkyl, alkoxy carbonylalkyl, aralkoxy carbonylalkyl, aralkylcarbonyloxyalkyl, alkylthio, cycloalkylthio, arylthio, aralkylthio, aralkylthiocarbonylthio, mercapto, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl wherein each of said


25

30

35

heteroaryl- and cycloheteroalkyl-containing groups has one or more hetero ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R¹ through R¹¹ may be further independently selected from amino and amido

5 radicals of the formula



10 wherein X is selected from oxygen atom or sulfur atom;

wherein each n is a number independently selected from zero to six, inclusive;

15 wherein each of R¹² through R²⁴ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;


20 and wherein each of R³ through R¹¹ may be further independently selected from hydroxy and from acidic moieties of the formula




25

wherein n is a number selected from zero through three, inclusive;

wherein A is selected from carboxylic acid and bioisosteres
30 of carboxylic acid selected from



wherein each W is independently selected from oxygen atom, sulfur atom and NR⁴³; wherein each of R³⁹, R⁴⁰, R⁴¹, R⁴² and R⁴³ is independently selected from hydrido, alkyl, haloalkyl, haloalkylsulfonyl, haloalkylcarbonyl, cycloalkyl, cycloalkylalkyl, aryl and aralkyl; wherein each of R³⁹, R⁴⁰, R⁴¹ and R⁴² may be further independently selected from amino radical of the formula




wherein each of R⁴⁴ and R⁴⁵ is independently selected from hydrido, alkyl, cycloalkyl, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl, and wherein R⁴⁴ and R⁴⁵ taken together may form a heterocyclic group having five to seven ring members including the nitrogen atom of said amino radical, which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms and which heterocyclic group may be saturated or partially unsaturated; wherein R⁴⁴ and R⁴⁵ taken together may form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino radical and which aromatic heterocyclic group may further contain one or more hetero atoms as ring atoms selected from oxygen, nitrogen and sulfur atoms; wherein each of R⁴⁴ and R⁴⁵ may be further independently selected from hydroxy, alkoxy, alkylthio, aryloxy, arylthio, aralkylthio and aralkoxy; and the amide, ester and salt derivatives of said acidic groups;

wherein said bioisostere of carboxylic acid may be further selected from heterocyclic acidic groups consisting of heterocyclic rings of four to about nine ring members,


which heterocyclic ring contains at least one hetero atom selected from oxygen, sulfur and nitrogen atoms, which heterocyclic ring may be saturated, fully unsaturated or partially unsaturated, and which heterocyclic ring may be attached at a single position selected from R³ through R¹¹ or may be attached at any two adjacent positions selected from R³ through R¹¹ so as to form a fused-ring system with one of the phenyl rings of the biphenyl moiety of Formula I; and the amide, ester and salt derivatives of said heterocyclic acidic groups;

wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, aryl and aralkyl;

and wherein any of the foregoing R¹ through R²⁴, Y and A groups having a substitutable position may be substituted by one or more groups selected from alkyl, difluoroalkyl, alkenyl, aralkyl, hydroxyalkyl, trifluoromethyl, alkoxy, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, carboxyl, mercaptocarbonyl, alkylthio, alkylthiocarbonyl, and amino and amido radicals of the formula



wherein X is selected from oxygen atom and sulfur atom; wherein each of R²⁵, R²⁶, R²⁷, R²⁸ and R²⁹ is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl and DR³⁰ and



wherein D is selected from oxygen atom and sulfur atom,
wherein R³⁰ is selected from hydrido, alkyl, cycloalkyl,
cycloalkylalkyl, aralkyl and aryl;

5 wherein each of R²⁵, R²⁶, R²⁷, R²⁸, R²⁹, R³¹ and R³² is
independently selected from hydrido, alkyl, cycloalkyl,
cyano, hydroxyalkyl, haloalkyl, cycloalkylalkyl,
alkoxyalkyl, alkanoyl, alkoxycarbonyl, carboxyl,
haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl;

10


with the proviso that at least one of said R¹ through R²⁴,
Y and A substituents contains a terminal primary or
secondary amino moiety or a moiety convertible to a primary
or secondary amino moiety;

15

or a tautomer thereof or a pharmaceutically-acceptable salt
thereof.

41. The composition of Claim 40 wherein m is
20 one; wherein each of R¹ and R² is independently selected
from alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl,
cycloalkylalkyl, alkoxy, aralkyl, aryl, aroyl, aryloxy,
aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl,
alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl,
25 cycloalkynyl, cyano, nitro, carboxyl, carboxyalkyl,
alkylcarbonyloxy, alkylcarbonyloxyalkyl,
alkoxycarbonylalkyl, aralkoxycarbonylalkyl,
aralkylcarbonyloxyalkyl, mercaptocarbonyl, mercaptoalkyl,
alkoxycarbonyloxy, alkylthio, cycloalkylthio, arylthio,
30 aralkylthio, mercapto, alkylsulfinyl, alkylsulfonyl,
aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl,
arylsulfonyl, phthalimido, phthalimidoalkyl, heteroaryl,
heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl
and cycloheteroalkylcarbonylalkyl wherein each of said
35 heteroaryl- and cycloheteroalkyl-containing groups has one
or more hetero ring atoms selected from oxygen, sulfur and
nitrogen atoms, and wherein each of R¹ through R¹¹ may be

further independently selected from amino and amido radicals of the formula



5

wherein X is selected from oxygen atom and sulfur atom;

10 wherein each n is a number independently selected from zero to six, inclusive;


15 wherein each of R¹² through R²⁴ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

20 wherein each of R³ through R¹¹ is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, aralkyl, aryl, aroyl, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl, alkylthio, aralkylthio, mercapto, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, 25 arylsulfonyl and heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms;

30 and wherein each of R³ through R¹¹ may be an acidic moiety further independently selected from acidic moieties of the formula




wherein n is a number selected from zero through three, inclusive; wherein A is selected from carboxylic acid and bioisosteres of carboxylic acid selected from



5

wherein each W is independently selected from oxygen atom, sulfur atom and NR^{43} ; wherein each of R^{39} , R^{42} and R^{43} is independently selected from hydrido, alkyl, haloalkyl, haloalkylsulfonyl, haloalkylcarbonyl, cycloalkyl, cycloalkylalkyl, aryl and aralkyl; wherein each of R^{39} and R^{42} may be further independently selected from amino radical of the formula

10



15

wherein each of R^{44} and R^{45} is independently selected from hydrido, alkyl, cycloalkyl, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl, and wherein R^{44} and R^{45} taken together may form a heterocyclic group having five to seven ring members including the nitrogen atom of said amino radical, which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms, and which heterocyclic group may be saturated or partially unsaturated; wherein R^{44} and R^{45} taken together may form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino radical and which aromatic heterocyclic group may further contain one or more hetero atoms as ring atoms selected from oxygen, nitrogen and sulfur atoms; and the amide, ester and salt derivatives of said acidic groups; wherein said bioisostere of carboxylic acid may be further selected from heterocyclic acidic groups consisting of heterocyclic rings of four to about nine ring members, which ring contains at least one

20

25

30

35

hetero atom, selected from oxygen, sulfur and nitrogen atoms, which heterocyclic ring may be saturated, fully unsaturated or partially unsaturated, and which heterocyclic ring may be attached at a single position
5 selected from R^3 through R^{11} or may be attached at any two adjacent positions selected from R^3 through R^{11} so as to form a fused-ring system with one of the phenyl rings of Formula I; and the amide, ester and salt derivatives of said heterocyclic acidic groups;

10

wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, aryl and aralkyl;

15 wherein each of R^1 through R^{11} , Y and A independently may be substituted at any substitutable position with one or more groups selected from alkyl, cycloalkyl, cycloalkylalkyl, hydroxy, oxo, trifluoromethyl, difluoroalkyl, alkoxycarbonyl, cyano, nitro, alkylsulfonyl,
20 haloalkylsulfonyl, aryl, aralkyl, alkoxy, aryloxy and aralkoxy;


with the proviso that at least one of said R^1 through R^{24} , Y and A substituents contains a terminal primary or
25 secondary amino moiety or a moiety convertible to a primary or secondary amino moiety;

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

30

42. The composition of Claim 41 wherein m is one; wherein each of R^1 and R^2 is independently selected from alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, aralkyl, aryl, benzoyl, phenoxy,
35 phenoxyalkyl, phenalkyloxy, phenylthio, phenalkylthio, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl,

- alkoxycarbonylalkyl, aralkoxycarbonylalkyl,
 aralkylcarbonyloxyalkyl, mercaptocarbonyl, mercaptoalkyl,
 alkoxycarbonyloxy, alkylthio, cycloalkylthio, phthalimido,
 phthalimidoalkyl, heteroaryl, heteroarylalkyl,
 5 cycloheteroalkyl, cycloheteroalkylalkyl and
 cycloheteroalkylcarbonylalkyl wherein each of said
 heteroaryl- and cycloheteroalkyl-containing groups has one
 or more hetero ring atoms selected from oxygen, sulfur and
 nitrogen atoms, and wherein each of R¹ through R¹¹ may be
 10 further independently selected from amino and amido
 radicals of the formula




- 15 wherein X is selected from oxygen atom and sulfur atom;
- wherein each n is a number independently selected from zero
 to six, inclusive;
- 20 wherein each of R¹² through R²⁴ is independently selected
 from hydrido, alkyl, cycloalkyl, cyano, amino,
 hydroxyalkyl, alkoxyalkyl, phenalkyl and phenyl;
- wherein each of R³ through R¹¹ is independently selected
 25 from hydrido, hydroxy, alkyl, hydroxyalkyl, halo,
 haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, phenalkyl,
 phenyl, benzoyl, phenoxy, phenalkyloxy, alkoxyalkyl,
 alkylcarbonyl, alkoxycarbonyl, alkenyl, cyano, nitro,
 carboxyl, alkylthio, mercapto and heteroaryl having one or
 30 more ring atoms selected from oxygen, sulfur and nitrogen
 atoms;

and wherein each of R³ through R¹¹ may be an acidic moiety further independently selected from acidic moieties of the formula




5

wherein n is a number selected from zero through two, inclusive; wherein A is selected from carboxylic acid and bioisosteres of carboxylic acid selected from



10

wherein each W is independently selected from oxygen atom, sulfur atom and NR⁴³; wherein each of R³⁹, R⁴² and R⁴³ is independently selected from hydrido, alkyl, haloalkyl, haloalkylsulfonyl, haloalkylcarbonyl, cycloalkyl, phenyl and benzyl; wherein each of R³⁹ and R⁴² may be further independently selected from amino radical of the formula



20

wherein each of R⁴⁴ and R⁴⁵ is independently selected from hydrido, alkyl, cycloalkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, benzyl and phenyl; and the amide, ester and salt derivatives of said acidic groups;

25

wherein said bioisostere of carboxylic acid may be further selected from heterocyclic acidic groups consisting of heterocyclic rings of four to about nine ring members, which ring contains at least one hetero atom, selected from oxygen, sulfur and nitrogen atoms, which heterocyclic ring may be saturated, fully unsaturated or partially unsaturated, and which heterocyclic ring may be attached at a single position selected from R³ through R¹¹ or may be attached at any two adjacent positions selected from R³ through R¹¹ so as to form a fused-ring system with one of

the phenyl rings of the biphenyl moiety of Formula I; and the amide, ester and salt derivatives of said heterocyclic acidic groups;


- 5 wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, phenyl, phenalkyl and aralkyl;

- 10 wherein each of R^1 through R^{11} , Y and A and independently may be substituted at any substitutable position with one or more groups selected from alkyl, cycloalkyl, cycloalkylalkyl, hydroxy, oxo, trifluoromethyl, difluoroalkyl, alkoxycarbonyl, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, aryl, aralkyl, alkoxy, aryloxy and
15 aralkoxy;

- with the proviso that at least one of said R^1 through R^{24} , Y and A substituents contains a terminal primary or secondary amino moiety or a moiety convertible to a primary
20 or secondary amino moiety;

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

- 25 43. The composition of Claim 42 wherein m is one; wherein each of R^1 and R^2 is independently selected from alkyl, aminoalkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, aralkyl, aryl, benzoyl, phenoxy, phenoxyalkyl, phenalkyloxy, phenylthio,
30 phenalkylthio, aralkoxy, alkoxyalkyl, acetyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, mercaptoalkyl, mercaptocarbonyl, alkoxycarbonyloxy, alkylcarbonyloxyalkyl, alkoxycarbonylalkyl,
35 aralkoxycarbonylalkyl, aralkylcarbonyloxyalkyl, phthalimido, phthalimidoalkyl, imidazoalkyl, tetrazole, tetrazolealkyl, alkylthio, cycloalkylthio, and amino and amido radicals of the formula




5 wherein X is selected from oxygen atom and sulfur atom;

wherein each n is a number independently selected from zero to six, inclusive;


10 wherein each of R¹² through R²⁴ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, hydroxyalkyl, alkoxyalkyl, phenalkyl and phenyl;

15 wherein each of R³ through R¹¹ is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, phenalkyl, phenyl, benzoyl, phenoxy, phenalkyloxy, alkoxyalkyl, acetyl, alkoxy carbonyl, alkenyl, cyano, nitro, carboxyl, alkylthio and mercapto;

20 and wherein each of R³ through R¹¹ may be an acidic moiety further independently selected from acidic moieties consisting of CO₂H, CO₂CH₃, SH, CH₂SH, C₂H₄SH, PO₃H₂, NHSO₂CF₃, NHSO₂C₆F₅, SO₃H, CONHNH₂, CONHNHSO₂CF₃, CONHOCH₃,
 25 CONHOC₂H₅, CONHCF₃, OH, CH₂OH, C₂H₄OH, OPO₃H₂, OSO₃H,




wherein each of R⁴⁶, R⁴⁷ and R⁴⁸ is independently selected
 5 from H, Cl, CN, NO₂, CF₃, C₂F₅, C₃F₇, CHF₂, CH₂F, CO₂CH₃,
 CO₂C₂H₅, SO₂CH₃, SO₂CF₃ and SO₂C₆F₅; wherein Z is selected
 from O, S, NR⁴⁹ and CH₂; wherein R⁴⁹ is selected from
 hydrido, CH₃ and CH₂C₆H₅; and wherein said acidic moiety
 10 may be a heterocyclic acidic group attached at any two
 adjacent positions of R³ through R¹¹ so as to form a fused
 ring system so as to include one of the phenyl rings of the
 biphenyl moiety of Formula I, said biphenyl fused ring
 system selected from



and the esters, amides and salts of said acidic moieties;

- 5 with the proviso that at least one of said R^1 through R^{24} substituents contains a terminal primary or secondary amino moiety or a moiety convertible to a primary or secondary amino moiety;
- 10 or a tautomer thereof or a pharmaceutically-acceptable salt thereof.



44. The composition of Claim 43 wherein m is one; wherein each of R^1 and R^2 is independently selected
- 15 from amino, aminomethyl, aminoethyl, aminopropyl, CH_2OH , CH_2OCOCH_3 , CH_2Cl , Cl , CH_2OCH_3 , $CH_2OCH(CH_3)_2$, I , CHO ,



-CH₂NHCO₂-(1-naphthyl), -CH₂NHCONHCH₃, -CH₂NHCONHC₂H₅,
 -CH₂NHCONHC₃H₇, -CH₂NHCONHC₄H₉, -CH₂NHCONHCH(CH₃)₂,
 -CH₂NHCONH(1-naphthyl), -CH₂NHCONH(1-adamantyl), CO₂H,



-CH₂CH₂CO-N $\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array}$ O, -CH₂CH₂CO-N $\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array}$, -CH₂CH₂CH₂CO₂H,

5 -CH₂CH₂F, -CH₂OCONHCH₃, -CH₂OCSNHCH₃, -CH₂NHCSOC₃H₇,


-CH₂CH₂CH₂F, -CH₂ONO₂, , -CH₂SH, -CH₂O-,

H, Cl, NO₂, CF₃, CH₂OH, Br, F, I, methyl, ethyl, n-propyl,
 isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-
 pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl,
 10 cyclohexyl, cyclohexylmethyl, 1-oxoethyl, 1-oxopropyl, 1-
 oxobutyl, 1-oxopentyl, 1,1-dimethoxypropyl, 1,1-
 dimethoxybutyl, 1,1-dimethoxypentyl, hydroxyalkyl, halo,
 difluoromethyl, 1,1-difluoroethyl, 1,1-difluoropropyl, 1,1-
 difluorobutyl and 1,1-difluoropentyl; wherein each of R³

15 through ¹¹ is hydrido, with the proviso that at least one
 of R⁵, R⁶, R⁸ and R⁹ is an acidic group selected from CO₂H,
 SH, PO₃H₂, SO₃H, CONHNH₂, CONHNHSO₂CF₃, OH,



and




20

wherein each of R⁴⁶ and R⁴⁷ is independently selected from
 Cl, CN, NO₂, CF₃, CO₂CH₃ and SO₂CF₃;

25 with the proviso that at least one of said R¹ through R¹¹
 substituents contains a terminal primary or secondary amino
 moiety or a moiety convertible to a primary or secondary
 amino moiety;

30 or a tautomer thereof or a pharmaceutically-acceptable salt
 thereof.

45. The composition of Claim 44 wherein m is one; wherein R¹ is selected from amino, aminomethyl, aminoethyl, aminopropyl, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclohexyl, cyclohexylmethyl, 1-oxoethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl, 1,1-dimethoxypropyl, 1,1-dimethoxybutyl, 1,1-dimethoxypentyl, hydroxyalkyl, halo, difluoromethyl, 1,1-difluoroethyl, 1,1-difluoropropyl, 1,1-difluorobutyl and 1,1-difluoropentyl; wherein R² is selected from ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, 4-methylbutyl, tert-butyl, n-pentyl and neopentyl; wherein each of R³, R⁴, R⁶, R⁷, R⁸, R¹⁰, and R¹¹ is hydrido; wherein one of R⁵ and R⁹ is hydrido and the other of R⁵ and R⁹ is an acidic group selected from CO₂H, SH, PO₃H₂, SO₃H, CONHNH₂, CONHNHSO₂CF₃, OH,



- 20 wherein each of R⁴⁶ and R⁴⁷ is independently selected from Cl, CN, NO₂, CF₃, CO₂CH₃ and SO₂CF₃;

- with the proviso that at least one of said R¹ through R¹¹ substituents contains a terminal primary or secondary amino moiety or a moiety convertible to a primary or secondary amino moiety;


or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

30

46. The composition of Claim 37 wherein said second residue forms a kidney-enzyme-cleavable amide bond with the residue of said angiotensin II antagonist compound.

35

47. The composition of Claim 37 wherein said second residue is preferably selected from a class of compounds of Formula II:




5 wherein each of R⁵⁰ and R⁵¹ may be independently selected from hydrido, alkylcarbonyl, alkoxycarbonyl, alkoxyalkyl, hydroxyalkyl and haloalkyl; and wherein G is selected from
 10 hydroxyl, halo, mercapto, -OR⁵², -SR⁵³ and
 NR^{54} with each of R⁵², R⁵³ and R⁵⁴ independently selected from alkyl; and wherein R⁵⁴ may be further selected from
 15 hydrido; with the proviso that said Formula II compound is selected such that formation of the cleavable amide bond occurs at carbonyl moiety attached at the gamma-position carbon of said Formula II compound.

48. The composition of Claim 47 wherein each G substituent is hydroxy.

49. The composition of Claim 48 wherein each G substituent is hydroxy; wherein R⁵⁰ is hydrido; and wherein R⁵¹ is selected from

25 O
 \parallel
 CR^{55} wherein R⁵⁵ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, iso-butyl, tert-butyl, n-pentyl, neopentyl, n-hexyl and chloromethyl.

50. The composition of Claim 49 wherein said
 30 second residue is



51. Conjugate of Claim 37 wherein said first residue is provided by a biphenylalkyl 1H-substituted-
 5 1,2,4-triazole angiotensin II antagonist compound containing a terminal primary or secondary amino moiety selected from amino and linear or branched aminoalkyl moieties containing linear or branched alkyl groups selected from aminomethyl, aminoethyl, aminopropyl,
 10 aminoisopropyl, aminobutyl, aminosecbutyl, aminoisobutyl, aminotertbutyl, aminopentyl, aminoisopentyl and aminoneopentyl.

52. The composition of Claim 37 wherein said
 15 first residue is provided by a biphenylalkyl 1H-substituted-1,2,4-triazole angiotensin II antagonist compound containing a moiety convertible to a primary or secondary amino terminal moiety.

20 53. The composition of Claim 52 wherein said moiety convertible to an amino terminal moiety is a carboxylic acid group reactable with an amino moiety of a diamino-terminated linker group to provide a terminal amino moiety which may then be further reacted with a carboxylic
 25 acid moiety of a compound providing said second residue so as to form a hydrolyzable amide bond.


54. The composition of Claim 53 wherein said
 30 diamino-terminated linker group is a divalent radical of Formula III:




wherein each of R²⁰⁰ and R²⁰¹ may be independently selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, alkoxyalkyl, hydroxyalkyl, aralkyl, aryl, haloalkyl, amino, monoalkylamino, dialkylamino, cyanoamino, carboxyalkyl, alkylsulfinyl, alkylsulfonyl, arylsulfinyl and arylsulfonyl; and wherein n is zero or a number selected from three through seven, inclusive.

55. The composition of Claim 54 wherein each of R²⁰⁰ and R²⁰¹ is hydrido.

56. The composition of Claim 53 wherein said diamino-terminated linker group is a divalent radical of Formula IV:




wherein each of Q and T is one or more groups independently selected from



wherein each of R²⁰² through R²⁰⁵ is independently selected from hydrido, hydroxy, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl, alkoxy, aralkoxy, aryloxy, alkoxyalkyl, haloalkyl, hydroxyalkyl, halo, cyano, amino,

monoalkylamino, dialkylamino, carboxy, carboxyalkyl, alkanoyl, alkenyl, cycloalkenyl and alkynyl.

57. The composition of Claim 56 wherein said
 5 diamino-terminated linker group is a divalent radical of
 Formula V:




- 10 wherein each of R²⁰² and R²⁰³ is independently selected
 from hydrido, hydroxy, alkyl, phenalkyl, phenyl, alkoxy,
 benzyloxy, phenoxy, alkoxyalkyl, hydroxyalkyl, halo, amino,
 monoalkylamino, dialkylamino, carboxy, carboxyalkyl and
 alkanoyl; and wherein each of p and q is a number
 15 independently selected from one through six, inclusive;
 with the proviso that when each of R²⁰² and R²⁰³ is selected
 from halo, hydroxy, amino, monoalkylamino and dialkylamino,
 then the carbon to which R²⁰² or R²⁰³ is attached in
 Formula V is not adjacent to a nitrogen atom of Formula V.
 20

58. The composition of Claim 57 wherein each of
 R²⁰² and R²⁰³ is independently selected from hydrido,
 hydroxy, alkyl, alkoxy, amino, monoalkylamino, carboxy,
 25 carboxyalkyl and alkanoyl; and wherein each of p and q is a
 number independently selected from two through four,
 inclusive.

59. The composition of Claim 58 wherein each of R²⁰² and R²⁰³ is independently selected from hydrido, amino, monoalkylamino and carboxyl; and wherein each of p and q is independently selected from the numbers two and three.

60. The composition of Claim 59 wherein each of R²⁰² and R²⁰³ is hydrido; and wherein each of p and q is two.

61. The composition of Claim 53 wherein said diamino-terminated linker group is a divalent radical of Formula VI:



wherein each of R²¹⁴ through R²¹⁷ is independently selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, hydroxyalkyl, alkoxyalkyl, aralkyl, aryl, haloalkyl, amino, monoalkylamino, dialkylamino, cyanoamino, carboxyalkyl, alkylsulfinyl, alkylsulfonyl, arylsulfinyl and arylsulfonyl; and wherein p is a number selected from one through six inclusive.

62. The composition of Claim 61 wherein each of R²¹⁴ and R²¹⁵ is hydrido; wherein each of R²¹⁶ and R²¹⁷ is independently selected from hydrido, alkyl, phenalkyl, phenyl, alkoxyalkyl, hydroxyalkyl, haloalkyl and carboxyalkyl; and wherein p is two or three.

63. The composition of Claim 62 wherein each of R²¹⁴ and R²¹⁵ is hydrido; wherein each of R²¹⁶ and R²¹⁷ is independently selected from hydrido and alkyl; and wherein p is two.

64. The composition of Claim 63 wherein each of R²¹⁴, R²¹⁵, R²¹⁶ and R²¹⁷ is hydrido; and wherein p is two.

65. The composition of Claim 45 wherein said
5 angiotensin II antagonist compound is selected from the group consisting of
methyl 4'-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-
biphenyl]-2-carboxylate;
4'-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-
10 biphenyl]-2-carboxylic acid;
4'-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-
biphenyl]-2-carboxylic acid, hydrazide;
4'-[(5-butyl-3-chloro-1H-1,2,4-triazol-1-yl)methyl][1,1'-
biphenyl]-2-carboxylic acid;
15 4'-[(3-butyl-5-chloro-1H-1,2,4-triazol-1-yl)methyl][1,1'-
biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-propyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-
biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-isopropyl-1H-1,2,4-triazol-1-
20 yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-secbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-
biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-isobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-
biphenyl]-2-carboxylic acid;
25 4'-[(5-butyl-3-tertbutyl-1H-1,2,4-triazol-1-
yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-pentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-
biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-isopentyl-1H-1,2,4-triazol-1-
30 yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-cyclohexyl-1H-1,2,4-triazol-1-
yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-cyclohexylmethyl-1H-1,2,4-triazol-1-
yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
35 4'-[[5-butyl-3-(2-cyclohexylethyl))-1H-1,2,4-triazol-1-
yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-cyclohexanoyl-1,2,4-triazol-1-
yl)methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[[5-butyl-3-(1-oxo-2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-phenyl-1,2,4,-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
5 4'-[(5-butyl-3-phenylmethyl-1,2,4,-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-butyl-3-(2-phenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-benzoyl-1,2,4,-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
10 4'-[[5-butyl-3-(1-oxo-2-phenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-butyl-3-(1,1-dimethoxypropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
15 4'-[[5-butyl-3-(1,1-dimethoxybutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-butyl-3-(1-oxopropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-butyl-3-(1-oxobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
20 4'-[[5-butyl-3-(1-oxopentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-butyl-3-(1,1-difluoroethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
25 4'-[[5-butyl-3-(1,1-difluoropropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-butyl-3-(1,1-difluorobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[3-butyl-5-(1,1-difluorobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
30 4'-[[5-butyl-3-(1,1-difluoropentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(3,5-dipropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
35 4'-[(3,5-isopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(3,5-disecbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[(3,5-diisobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(3,5-ditertbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 5 4'-[(3,5-dipentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(3,5-diisopentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 10 5-[4'-[(5-butyl-3-amino-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-butyl-3-aminomethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-butyl-3-aminoethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 15 5-[4'-[(5-butyl-3-aminopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-butyl-3-aminobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminophenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 20 5-[4'-[[3-butyl-5-(4-aminophenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminophenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 25 5-[4'-[[3-butyl-5-(4-aminomethylphenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminomethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminoethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 30 5-[4'-[[3-butyl-5-(4-aminocyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminocyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
- 35 5-[4'-[[3-butyl-5-(4-aminocyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminomethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;

- 5-[4'-[[3-butyl-5-(4-aminomethylcyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminoethylcyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5 5-[4'-[(5-butyl-3-carboxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-butyl-3-carboxymethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-butyl-3-carboxyethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
10 5-[4'-[(5-butyl-3-carboxypropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-butyl-3-carboxybutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
15 5-[4'-[[3-butyl-5-(4-carboxyphenyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-carboxyphenylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-carboxyphenylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
20 5-[4'-[[3-butyl-5-(4-carboxymethylphenylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-carboxymethylphenyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
25 5-[4'-[[3-butyl-5-(4-carboxyethylphenyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-carboxycyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-carboxycyclohexylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
30 5-[4'-[[3-butyl-5-(4-carboxycyclohexylethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-carboxymethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
35 tetrazole;
5-[4'-[[3-butyl-5-(4-carboxymethylcyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole; and

5-[4'-[[3-butyl-5-(4-carboxyethylcyclohexyl)-1H-1,2,4-triazol-1-yl]methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole.

66. The composition of Claim 55 wherein said
5 angiotensin II antagonist compound is N-acetylglutamic acid, 5-[[4'-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]]carbonylhydrazide.

67. The composition of Claim 51 wherein said
10 angiotensin II antagonist compound is N²-acetyl-N-[[5-butyl-1-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]-1H-1,2,4-triazol-3-yl]methyl]-L-glutamine.

68. The composition of Claim 55 wherein said
15 angiotensin II antagonist compound is N-acetyl-L-glutamic acid, 5-[5-butyl-1-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]-1H-1,2,4-triazol-3-yl]acetylhydrazide.

69. A method for treating a circulatory
20 disorder, said method comprising administering to a patient afflicted with or susceptible to said disorder a therapeutically-effective amount of a renal-selective conjugate, said renal-selective conjugate comprising a residue of a biphenylalkyl 1H-substituted-1,2,4-triazole
25 angiotensin II antagonist compound.


70. The method of Claim 69 comprising a first residue and a second residue, said first and second residues connected together by a cleavable bond, wherein
30 said first residue is provided by said biphenylalkyl 1H-substituted-1,2,4-triazole angiotensin II antagonist compound, and wherein said second residue is capable of being cleaved from said first residue selectively in the kidney.

35

71. The method of Claim 70 wherein said first and second residues are provided by precursor compounds wherein the precursor compound of one of said first and

second residues has a reactable carboxylic acid moiety and the precursor of the other of said first and second residues has a reactable amino moiety or a moiety convertible to a reactable amino moiety, whereby a
 5 cleavable bond may be formed between said carboxylic acid moiety and said amino moiety.

72. The method of Claim 71 wherein said angiotensin II antagonist compound is selected from a class
 10 of compounds defined by Formula I:




15 wherein m is a number selected from one to four, inclusive;

wherein each of R¹ through R¹¹ is independently selected from hydrido, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, formyl, alkoxy, aralkyl, aryl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkylcarbonylalkyl, alkoxy carbonyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxy carbonylalkyl, aralkoxy carbonylalkyl, aralkylcarbonyloxyalkyl, mercaptocarbonyl, mercaptothiocarbonyl, mercaptoalkyl, alkoxy carbonyloxy, alkylthio, cycloalkylthio, alkylthiocarbonyl, alkylcarbonylthio, alkylthiocarbonyloxy, alkylthiocarbonylthio, alkylthiothiocarbonyl, alkylthiothiocarbonylthio, arylthio, arylthiocarbonyl, arylcarbonylthio, arylthiocarbonyloxy, arylthiocarbonylthio, arylthiothiocarbonyl, arylthiothiocarbonylthio, aralkylthio, aralkylthiocarbonyl,

20
25
30

aralkylcarbonylthio, aralkylthiocarbonyloxy,
 aralkylthiocarbonylthio, alkylthiocarbonyl,
 aralkylthiocarbonylthio, mercapto, alkylsulfinyl,
 alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl,
 5 arylsulfinyl, arylsulfonyl, phthalimido, phthalimidoalkyl,
 heteroaryl, heteroarylalkyl, cycloheteroalkyl,
 cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl
 wherein each of said heteroaryl- and cyclohetero-containing
 groups has one or more ring atoms selected from oxygen,
 10 sulfur and nitrogen atoms, and wherein each of R¹ through
 R¹¹ may be further independently selected from amino and
 amido radicals of the formula



15

wherein X is oxygen atom or sulfur atom;

wherein each n is a number independently selected from zero
 to six, inclusive;

20

wherein each of R¹² through R²⁴ is independently selected
 from hydrido, alkyl, cycloalkyl, cyano, amino,
 monoalkylamino, dialkylamino, hydroxyalkyl,
 cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl, and wherein
 25 R¹² and R¹³ taken together, R¹⁴ and R¹⁵ taken together, R¹⁶
 and R¹⁷ taken together, R¹⁹ and R²⁰ taken together and R²¹
 and R²² taken together may each form a heterocyclic group
 having five to seven ring members including the nitrogen
 atom of said amino or amido radical and which heterocyclic
 30 group may further contain one or more hetero atoms as ring
 members selected from oxygen, nitrogen and sulfur atoms and
 which heterocyclic group may be saturated or partially

unsaturated; wherein R¹² and R¹³ taken together, R¹⁴ and R¹⁵ taken together, R¹⁹ and R²⁰ taken together and R²¹ and R²² taken together may each form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino or amido radical and which aromatic heterocyclic group may further contain one or more hetero atoms as ring atoms selected from oxygen, nitrogen and sulfur atoms;

10 and wherein each of R³ through R¹¹ may be further independently selected from hydroxy and from acidic moieties of the formula


$$-Y_n A$$

15


wherein n is a number selected from zero through three, inclusive, and wherein A is an acidic group selected to contain at least one acidic hydrogen atom, and the amide, ester and salt derivatives of said acidic moieties;

20 wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, aryl, aralkyl and heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms;


25 and wherein any of the foregoing R¹ through R²⁴, Y and A groups having a substitutable position may be substituted with one or more groups selected from hydroxy, alkyl, alkenyl, alkynyl, aralkyl, hydroxyalkyl, trifluoromethyl, difluoroalkyl, oxo, alkoxy, aryloxy, aralkoxy, aralkylthio, 30 alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aroyl, cycloalkenyl, cyano, cyanoamino, nitro, alkylcarbonyloxy, alkoxy carbonyloxy, alkylcarbonyl, alkoxy carbonyl, carboxyl, mercapto, mercaptocarbonyl, alkylthio, arylthio, alkylthiocarbonyl, alkylsulfanyl, alkylsulfonyl, 35 aralkylsulfanyl, aralkylsulfonyl, arylsulfanyl, arylsulfonyl, heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms, and amino and amido radicals of the formula



- wherein X is selected from oxygen atom and sulfur atom;
 5 wherein each of R²⁵, R²⁶, R²⁷, R²⁸ and R²⁹ is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl, DR³⁰ and



- 10 wherein D is selected from oxygen atom and sulfur atom and R³⁰ is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl and aryl; wherein each of R²⁵, R²⁶, R²⁷, R²⁸, R²⁹, R³¹ and R³² is independently selected
 15 from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, carboxyl, alkylsulfinyl, alkylsulfonyl, arylsulfinyl, arylsulfonyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl, and wherein each of
 20 R²⁵, R²⁶, R²⁷, R²⁸, R²⁹, R³¹ and R³² is further independently selected from amino and amido radicals of the formula



25

wherein X is oxygen atom or sulfur atom;


- wherein each of R³³, R³⁴, R³⁵, R³⁶, R³⁷ and R³⁸ is independently selected from hydrido, alkyl, cycloalkyl,
 30 cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl, and wherein R²⁶ and

- R²⁷ taken together and R²⁸ and R²⁹ taken together may each form a heterocyclic group having five to seven ring members including the nitrogen atom of said amino or amido radical, which heterocyclic group may further contain one or more
5 hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms and which heterocyclic group may be saturated or partially unsaturated; wherein R²⁶ and R²⁷ taken together and R³¹ and R³² taken together may each form an aromatic heterocyclic group having five ring members
10 including the nitrogen atom of said amino or amido radical and which aromatic heterocyclic group may further contain one or more hetero atoms as ring atoms selected from oxygen, nitrogen and sulfur atoms;
- 15 with the proviso that at least one of said R¹ through R²⁴, Y and A substituents contains a terminal primary or secondary amino moiety or a moiety convertible to a primary or secondary amino moiety.
- 20 or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

73. The method of Claim 72 wherein m is one; wherein each of R¹ through R¹¹ is independently selected
25 from alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, aralkyl, aryl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl, carboxyalkyl,
30 alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxy carbonylalkyl, aralkoxy carbonylalkyl, aralkylcarbonyloxyalkyl, mercaptocarbonyl, mercaptothiocarbonyl, mercaptoalkyl, alkoxy carbonyloxy, alkylthio, cycloalkylthio, alkylthiocarbonyl,
35 alkylcarbonylthio, alkylthiocarbonyloxy, alkylthio-carbonylthio, alkylthiothiocarbonyl, alkylthiothio-carbonylthio, arylthio, arylthiocarbonyl, arylcarbonylthio, arylthiocarbonyloxy, arylthiocarbonylthio,

arylthiothiocarbonyl, arylthiothiocarbonylthio,
 aralkylthio, aralkylthiocarbonyl, aralkylcarbonylthio,
 aralkylthiocarbonyloxy, aralkylthiocarbonylthio,
 aralkylthiocarbonyl, aralkylthiocarbonylthio, mercapto,
 5 alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl,
 aralkylsulfonyl, arylsulfinyl, arylsulfonyl, phthalimido,
 phthalimidoalkyl, heteroaryl, heteroarylalkyl,
 cycloheteroalkyl, cycloheteroalkylalkyl and
 cycloheteroalkylcarbonylalkyl wherein each of said
 10 heteroaryl- and cycloheteroalkyl-containing groups has
 one or more hetero ring atoms selected from oxygen, sulfur
 and nitrogen atoms, and wherein each of R¹ through R¹¹ may
 be further independently selected from amino and amido
 radicals of the formula

15



wherein X is selected from oxygen atom or sulfur atom;

20 wherein each n is a number independently selected from zero to six, inclusive;


wherein each of R¹² through R²⁴ is independently selected
 from hydrido, alkyl, cycloalkyl, cyano, amino,
 25 monoalkylamino, dialkylamino, hydroxyalkyl,
 cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

and wherein each of R³ through R¹¹ may be further
 independently selected from hydroxy and from acidic
 30 moieties of the formula




wherein n is a number selected from zero through three, inclusive; wherein A is an acidic group selected from acids containing one or more atoms selected from oxygen, sulfur, phosphorus and nitrogen atoms, and wherein said acidic group is selected to contain at least one acidic hydrogen atom, and the amide, ester and salt derivatives of said acidic moieties; wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, aryl, aralkyl and heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms;

and wherein any of the foregoing R^1 through R^{24} , Y and A groups having a substitutable position may be substituted with one or more groups selected from alkyl, alkenyl, aralkyl, hydroxyalkyl, trifluoromethyl, difluoroalkyl, alkoxy, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxycarbonyl, carboxyl, mercaptocarbonyl, alkylthio, alkylthiocarbonyl, and amino and amido radicals of the formula




wherein X is selected from oxygen atom and sulfur atom; wherein each of R^{25} , R^{26} , R^{27} , R^{28} and R^{29} is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl, and DR^{30} and



wherein D is selected from oxygen atom and sulfur atom, and R^{30} is selected from hydrido, alkyl, cycloalkyl,

- cycloalkylalkyl, aralkyl and aryl; wherein each of R²⁵, R²⁶, R²⁷, R²⁸, R²⁹, R³¹ and R³² is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, alkanoyl, alkoxycarbonyl, carboxyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl, and wherein each of R²⁵, R²⁶, R²⁷, R²⁸, R²⁹, R³¹ and R³² is further independently selected from amino and amido radicals of the formula

10




wherein X is selected from oxygen atom or sulfur atom;

- wherein each of R³³ through R³⁸ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl;
- with the proviso that at least one of said R¹ through R²⁴, Y and A substituents contains a terminal primary or secondary amino moiety or a moiety convertible to a primary or secondary amino moiety;
- or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

74. The method of Claim 73 wherein m is one;
- wherein each of R¹ through R¹¹ is independently selected from alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, aralkyl, aryl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, mercaptocarbonyl, alkoxycarbonyloxy,

- alkylcarbonyloxyalkyl, alkoxycarbonylalkyl,
 aralkoxycarbonylalkyl, aralkylcarbonyloxyalkyl, alkylthio,
 cycloalkylthio, arylthio, aralkylthio,
 aralkylthiocarbonylthio, mercapto, alkylsulfinyl,
 5 alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl,
 arylsulfinyl, arylsulfonyl, phthalimido, phthalimidoalkyl,
 heteroaryl, heteroarylalkyl, cycloheteroalkyl,
 cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl
 wherein each of said heteroaryl- and cycloheteroalkyl-
 10 containing groups has one or more hetero ring atoms
 selected from oxygen, sulfur and nitrogen atoms, and
 wherein each of R¹ through R¹¹ may be further independently
 selected from amino and amido radicals of the formula



- wherein X is selected from oxygen atom or sulfur atom;
- 20 wherein each n is a number independently selected from zero to six, inclusive;


- wherein each of R¹² through R²⁴ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino,
 25 monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

- and wherein each of R³ through R¹¹ may be further independently selected from hydroxy and from acidic
 30 moieties of the formula




wherein n is a number selected from zero through three, inclusive;

- 5 wherein A is selected from carboxylic acid and bioisosteres of carboxylic acid selected from



- 10 wherein each W is independently selected from oxygen atom, sulfur atom and NR^{43} ; wherein each of R^{39} , R^{40} , R^{41} , R^{42} and R^{43} is independently selected from hydrido, alkyl, haloalkyl, haloalkylsulfonyl, haloalkylcarbonyl, cycloalkyl, cycloalkylalkyl, aryl and aralkyl; wherein each
- 15 of R^{39} , R^{40} , R^{41} and R^{42} may be further independently selected from amino radical of the formula




- 20 wherein each of R^{44} and R^{45} is independently selected from hydrido, alkyl, cycloalkyl, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl, and wherein R^{44} and R^{45} taken together may form a heterocyclic group having five to seven ring members including the nitrogen
- 25 atom of said amino radical, which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms and which heterocyclic group may be saturated or partially unsaturated; wherein R^{44} and R^{45} taken together may form an
- 30 aromatic heterocyclic group having five ring members including the nitrogen atom of said amino radical and which aromatic heterocyclic group may further contain one or more hetero atoms as ring atoms selected from oxygen, nitrogen and sulfur atoms; wherein each of R^{44} and R^{45} may be
- 35 further independently selected from hydroxy, alkoxy,

alkylthio, aryloxy, arylthio, aralkylthio and aralkoxy; and the amide, ester and salt derivatives of said acidic groups;

- 5 wherein said bioisostere of carboxylic acid may be further selected from heterocyclic acidic groups consisting of heterocyclic rings of four to about nine ring members, which heterocyclic ring contains at least one hetero atom selected from oxygen, sulfur and nitrogen atoms, which
 10 heterocyclic ring may be saturated, fully unsaturated or partially unsaturated, and which heterocyclic ring may be attached at a single position selected from R³ through R¹¹ or may be attached at any two adjacent positions selected from R³ through R¹¹ so as to form a fused-ring system with
 15 one of the phenyl rings of the biphenyl moiety of Formula I; and the amide, ester and salt derivatives of said heterocyclic acidic groups;


- wherein Y is a spacer group independently selected from one
 20 or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, aryl and aralkyl;

- and wherein any of the foregoing R¹ through R²⁴, Y and A groups having a substitutable position may be substituted
 25 by one or more groups selected from alkyl, difluoroalkyl, alkenyl, aralkyl, hydroxyalkyl, trifluoromethyl, alkoxy, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, carboxyl, mercaptocarbonyl, alkylthio, alkylthiocarbonyl, and amino and amido radicals of the
 30 formula



- wherein X is selected from oxygen atom and sulfur atom;
 35 wherein each of R²⁵, R²⁶, R²⁷, R²⁸ and R²⁹ is selected from

hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl and DR³⁰ and



5

wherein D is selected from oxygen atom and sulfur atom, wherein R³⁰ is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl and aryl;

10

wherein each of R²⁵, R²⁶, R²⁷, R²⁸, R²⁹, R³¹ and R³² is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, alkanoyl, alkoxycarbonyl, carboxyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl;

15


with the proviso that at least one of said R¹ through R²⁴, Y and A substituents contains a terminal primary or secondary amino moiety or a moiety convertible to a primary or secondary amino moiety;

20

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

75. The method of Claim 74 wherein m is one;
- 25 wherein each of R¹ and R² is independently selected from alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, aralkyl, aryl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxycarbonylalkyl, aralkoxycarbonylalkyl, aralkylcarbonyloxyalkyl, mercaptocarbonyl, mercaptoalkyl, alkoxycarbonyloxy, alkylthio, cycloalkylthio, arylthio, aralkylthio, mercapto, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, phthalimido, phthalimidoalkyl, heteroaryl,
- 35

heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl wherein each of said heteroaryl- and cycloheteroalkyl-containing groups has one or more hetero ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R^1 through R^{11} may be further independently selected from amino and amido radicals of the formula



wherein X is selected from oxygen atom and sulfur atom;

wherein each n is a number independently selected from zero to six, inclusive;

wherein each of R^{12} through R^{24} is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;


wherein each of R^3 through R^{11} is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, aralkyl, aryl, aroyl, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl, alkylthio, aralkylthio, mercapto, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl and heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms;

and wherein each of R^3 through R^{11} may be an acidic moiety further independently selected from acidic moieties of the formula



5


wherein n is a number selected from zero through three, inclusive; wherein A is selected from carboxylic acid and bioisosteres of carboxylic acid selected from



10

wherein each W is independently selected from oxygen atom, sulfur atom and NR^{43} ; wherein each of R^{39} , R^{42} and R^{43} is independently selected from hydrido, alkyl, haloalkyl, haloalkylsulfonyl, haloalkylcarbonyl, cycloalkyl, cycloalkylalkyl, aryl and aralkyl; wherein each of R^{39} and R^{42} may be further independently selected from amino radical of the formula

15



20

wherein each of R^{44} and R^{45} is independently selected from hydrido, alkyl, cycloalkyl, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl, and wherein R^{44} and R^{45} taken together may form a heterocyclic group having five to seven ring members including the nitrogen atom of said amino radical, which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms, and which heterocyclic group may be saturated or partially unsaturated; wherein R^{44} and R^{45} taken together may form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino radical and which aromatic heterocyclic group may further contain one or more hetero atoms as ring atoms selected from oxygen, nitrogen

25

30

35

and sulfur atoms; and the amide, ester and salt derivatives of said acidic groups; wherein said bioisostere of carboxylic acid may be further selected from heterocyclic acidic groups consisting of heterocyclic rings of four to
5 about nine ring members, which ring contains at least one hetero atom, selected from oxygen, sulfur and nitrogen atoms, which heterocyclic ring may be saturated, fully unsaturated or partially unsaturated, and which heterocyclic ring may be attached at a single position
10 selected from R^3 through R^{11} or may be attached at any two adjacent positions selected from R^3 through R^{11} so as to form a fused-ring system with one of the phenyl rings of Formula I; and the amide, ester and salt derivatives of said heterocyclic acidic groups;

15 wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, aryl and aralkyl;

20 wherein each of R^1 through R^{11} , Y and A independently may be substituted at any substitutable position with one or more groups selected from alkyl, cycloalkyl, cycloalkylalkyl, hydroxy, oxo, trifluoromethyl, difluoroalkyl, alkoxycarbonyl, cyano, nitro, alkylsulfonyl,
25 haloalkylsulfonyl, aryl, aralkyl, alkoxy, aryloxy and aralkoxy;


with the proviso that at least one of said R^1 through R^{24} , Y and A substituents contains a terminal primary or
30 secondary amino moiety or a moiety convertible to a primary or secondary amino moiety;

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

35

76. The method of Claim 75 wherein m is one; wherein each of R^1 and R^2 is independently selected from alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl,

- cycloalkylalkyl, alkoxy, aralkyl, aryl, benzoyl, phenoxy, phenoxyalkyl, phenalkyloxy, phenylthio, phenalkylthio, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl,
- 5 carboxyalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxycarbonylalkyl, aralkoxycarbonylalkyl, aralkylcarbonyloxyalkyl, mercaptocarbonyl, mercaptoalkyl, alkoxycarbonyloxy, alkylthio, cycloalkylthio, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl,
- 10 cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl wherein each of said heteroaryl- and cycloheteroalkyl-containing groups has one or more hetero ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R¹ through R¹¹ may be
- 15 further independently selected from amino and amido radicals of the formula



- 20 wherein X is selected from oxygen atom and sulfur atom;

wherein each n is a number independently selected from zero to six, inclusive;

- 25 wherein each of R¹² through R²⁴ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, hydroxyalkyl, alkoxyalkyl, phenalkyl and phenyl;

- wherein each of R³ through R¹¹ is independently selected
- 30 from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, phenalkyl, phenyl, benzoyl, phenoxy, phenalkyloxy, alkoxyalkyl,

alkylcarbonyl, alkoxycarbonyl, alkenyl, cyano, nitro, carboxyl, alkylthio, mercapto and heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms;


5

and wherein each of R^3 through R^{11} may be an acidic moiety further independently selected from acidic moieties of the formula



10


wherein n is a number selected from zero through two, inclusive; wherein A is selected from carboxylic acid and bioisosteres of carboxylic acid selected from



15

wherein each W is independently selected from oxygen atom, sulfur atom and NR^{43} ; wherein each of R^{39} , R^{42} and R^{43} is independently selected from hydrido, alkyl, haloalkyl, haloalkylsulfonyl, haloalkylcarbonyl, cycloalkyl, phenyl and benzyl; wherein each of R^{39} and R^{42} may be further independently selected from amino radical of the formula

20



25

wherein each of R^{44} and R^{45} is independently selected from hydrido, alkyl, cycloalkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, benzyl and phenyl; and the amide, ester and salt derivatives of said acidic groups;

30


wherein said bioisostere of carboxylic acid may be further selected from heterocyclic acidic groups consisting of heterocyclic rings of four to about nine ring members, which ring contains at least one hetero atom, selected from oxygen, sulfur and nitrogen atoms, which heterocyclic ring

35

- may be saturated, fully unsaturated or partially unsaturated, and which heterocyclic ring may be attached at a single position selected from R^3 through R^{11} or may be attached at any two adjacent positions selected from R^3 through R^{11} so as to form a fused-ring system with one of the phenyl rings of the biphenyl moiety of Formula I; and the amide, ester and salt derivatives of said heterocyclic acidic groups;
- 5
- 10 wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, phenyl, phenalkyl and aralkyl;
- wherein each of R^1 through R^{11} , Y and A and independently
- 15 may be substituted at any substitutable position with one or more groups selected from alkyl, cycloalkyl, cycloalkylalkyl, hydroxy, oxo, trifluoromethyl, difluoroalkyl, alkoxycarbonyl, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, aryl, aralkyl, alkoxy, aryloxy and
- 20 aralkoxy;
- with the proviso that at least one of said R^1 through R^{24} , Y and A substituents contains a terminal primary or secondary amino moiety or a moiety convertible to a primary
- 25 or secondary amino moiety;
- or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

- 30 77. The method of Claim 76 wherein m is one; wherein each of R^1 and R^2 is independently selected from alkyl, aminoalkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, aralkyl, aryl, benzoyl, phenoxy, phenoxyalkyl, phenalkyloxy, phenylthio,
- 35 phenalkylthio, aralkoxy, alkoxyalkyl, acetyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, mercaptoalkyl, mercaptocarbonyl, alkoxycarbonyloxy,

- alkylcarbonyloxyalkyl, alkoxycarbonylalkyl,
 aralkoxycarbonylalkyl, aralkylcarbonyloxyalkyl,
 phthalimido, phthalimidoalkyl, imidazoalkyl, tetrazole,
 tetrazolealkyl, alkylthio, cycloalkylthio, and amino and
 5 amido radicals of the formula




- 10 wherein X is selected from oxygen atom and sulfur atom;
 wherein each n is a number independently selected from zero
 to six, inclusive;


- 15 wherein each of R¹² through R²⁴ is independently selected
 from hydrido, alkyl, cycloalkyl, cyano, amino,
 hydroxyalkyl, alkoxyalkyl, phenalkyl and phenyl;

- wherein each of R³ through R¹¹ is independently selected
 20 from hydrido, hydroxy, alkyl, hydroxyalkyl, halo,
 haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, phenalkyl,
 phenyl, benzoyl, phenoxy, phenalkyloxy, alkoxyalkyl,
 acetyl, alkoxycarbonyl, alkenyl, cyano, nitro, carboxyl,
 alkylthio and mercapto;

- 25 and wherein each of R³ through R¹¹ may be an acidic moiety
 further independently selected from acidic moieties
 consisting of CO₂H, CO₂CH₃, SH, CH₂SH, C₂H₄SH, PO₃H₂,
 NHSO₂CF₃, NHSO₂C₆F₅, SO₃H, CONHNH₂, CONHNHSO₂CF₃, CONHOCH₃,
 30 CONHOC₂H₅, CONHCF₃, OH, CH₂OH, C₂H₄OH, OPO₃H₂, OSO₃H,






wherein each of R⁴⁶, R⁴⁷ and R⁴⁸ is independently selected
 5 from H, Cl, CN, NO₂, CF₃, C₂F₅, C₃F₇, CHF₂, CH₂F, CO₂CH₃,
 CO₂C₂H₅, SO₂CH₃, SO₂CF₃ and SO₂C₆F₅; wherein Z is selected
 from O, S, NR⁴⁹ and CH₂; wherein R⁴⁹ is selected from
 hydrido, CH₃ and CH₂C₆H₅; and wherein said acidic moiety
 10 may be a heterocyclic acidic group attached at any two
 adjacent positions of R³ through R¹¹ so as to form a fused
 ring system so as to include one of the phenyl rings of the
 biphenyl moiety of Formula I, said biphenyl fused ring
 system selected from



and the esters, amides and salts of said acidic moieties;

- 5 with the proviso that at least one of said R^1 through R^{24} substituents contains a terminal primary or secondary amino moiety or a moiety convertible to a primary or secondary amino moiety;
- 10 or a tautomer thereof or a pharmaceutically-acceptable salt thereof.



78. The method of Claim 77 wherein m is one; wherein each of R^1 and R^2 is independently selected from
- 15 amino, aminomethyl, aminoethyl, aminopropyl, CH_2OH , CH_2OCOCH_3 , CH_2Cl , Cl , CH_2OCH_3 , $CH_2OCH(CH_3)_2$, I , CHO ,





- 20 $-CH_2NHCO_2CH_2(CH_3)_2$, $-CH_2NHCO_2C_4H_9$, CH_2NHCO_2 -adamantyl,

-CH₂NHCO₂-(1-naphthyl), -CH₂NHCONHCH₃, -CH₂NHCONHC₂H₅,
 -CH₂NHCONHC₃H₇, -CH₂NHCONHC₄H₉, -CH₂NHCONHCH(CH₃)₂,
 -CH₂NHCONH(1-naphthyl), -CH₂NHCONH(1-adamantyl), CO₂H,


-CH₂CH₂CO-N $\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array}$ O, -CH₂CH₂CO-N $\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array}$, -CH₂CH₂CH₂CO₂H,
 5 -CH₂CH₂F, -CH₂OCONHCH₃, -CH₂OCSNHCH₃, -CH₂NHCSOC₃H₇,

-CH₂CH₂CH₂F, -CH₂ONO₂, , -CH₂SH, -CH₂O-,

H, Cl, NO₂, CF₃, CH₂OH, Br, F, I, methyl, ethyl, n-propyl,
 isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-
 pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl,
 10 cyclohexyl, cyclohexylmethyl, 1-oxoethyl, 1-oxopropyl, 1-
 oxobutyl, 1-oxopentyl, 1,1-dimethoxypropyl, 1,1-
 dimethoxybutyl, 1,1-dimethoxypentyl, hydroxyalkyl, halo,
 difluoromethyl, 1,1-difluoroethyl, 1,1-difluoropropyl, 1,1-
 difluorobutyl and 1,1-difluoropentyl; wherein each of R³
 15 through ¹¹ is hydrido, with the proviso that at least one
 of R⁵, R⁶, R⁸ and R⁹ is an acidic group selected from CO₂H,
 SH, PO₃H₂, SO₃H, CONHNH₂, CONHNHSO₂CF₃, OH,



and



20

wherein each of R⁴⁶ and R⁴⁷ is independently selected from
 Cl, CN, NO₂, CF₃, CO₂CH₃ and SO₂CF₃;



25

with the proviso that at least one of said R¹ through R¹¹
 substituents contains a terminal primary or secondary amino
 moiety or a moiety convertible to a primary or secondary
 amino moiety;


30

or a tautomer thereof or a pharmaceutically-acceptable salt
 thereof.

79. The method of Claim 78 wherein m is one;
 wherein R¹ is selected from amino, aminomethyl, aminoethyl,
 aminopropyl, methyl, ethyl, n-propyl, isopropyl, n-butyl,
 sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl,
 5 neopentyl, phenyl, benzyl, phenethyl, cyclohexyl,
 cyclohexylmethyl, 1-oxoethyl, 1-oxopropyl, 1-oxobutyl, 1-
 oxopentyl, 1,1-dimethoxypropyl, 1,1-dimethoxybutyl, 1,1-
 dimethoxypentyl, hydroxyalkyl, halo, difluoromethyl, 1,1-
 difluoroethyl, 1,1-difluoropropyl, 1,1-difluorobutyl and
 10 1,1-difluoropentyl; wherein R² is selected from ethyl, n-
 propyl, isopropyl, n-butyl, sec-butyl, isobutyl, 4-
 methylbutyl, tert-butyl, n-pentyl and neopentyl; wherein
 each of R³, R⁴, R⁶, R⁷, R⁸, R¹⁰, and R¹¹ is hydrido; wherein
 one of R⁵ and R⁹ is hydrido and the other of R⁵ and R⁹ is
 15 an acidic group selected from CO₂H, SH, PO₃H₂, SO₃H,
 CONHNH₂, CONHNHSO₂CF₃, OH,



and



20 wherein each of R⁴⁶ and R⁴⁷ is independently selected from
 Cl, CN, NO₂, CF₃, CO₂CH₃ and SO₂CF₃;


with the proviso that at least one of said R¹ through R¹¹
 substituents contains a terminal primary or secondary amino
 25 moiety or a moiety convertible to a primary or secondary
 amino moiety;

or a tautomer thereof or a pharmaceutically-acceptable salt
 thereof.

30

80. The method of Claim 71 wherein said second
 residue forms a kidney-enzyme-cleavable amide bond with the
 residue of said angiotensin II antagonist compound.

81. The method of Claim 71 wherein said second residue is preferably selected from a class of compounds of Formula II:



5

wherein each of R^{50} and R^{51} may be independently selected from hydrido, alkylcarbonyl, alkoxycarbonyl, alkoxyalkyl, hydroxyalkyl and haloalkyl; and wherein G is selected from hydroxyl, halo, mercapto, $-\text{OR}^{52}$, $-\text{SR}^{53}$ and $-\text{NR}^{54}$ with each of R^{52} , R^{53} and R^{54} is independently selected from hydrido and alkyl; with the proviso that said Formula II compound is selected such that formation of the cleavable amide bond occurs at carbonyl moiety attached at the gamma-position carbon of said Formula II compound.

10

15

82. The method of Claim 81 wherein each G substituent is hydroxy.

20


83. The method of Claim 82 wherein each G substituent is hydroxy; wherein R^{50} is hydrido; and wherein R^{51} is selected from

25

$\begin{array}{c} \text{O} \\ \parallel \\ -\text{CR}^{55} \end{array}$ wherein R^{55} is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, iso-butyl, tert-butyl, n-pentyl, neopentyl, n-hexyl and chloromethyl.

84. The method of Claim 83 wherein said second residue is

30




85. The method of Claim 71 wherein said first residue is a biphenylalkyl 1H-substituted-1,2,4-triazole
 5 angiotensin II antagonist compound containing a terminal primary or secondary amino moiety selected from amino and linear or branched aminoalkyl moieties containing linear or branched alkyl groups selected from aminomethyl,
 aminoethyl, aminopropyl, aminoisopropyl, aminobutyl,
 10 aminosecbutyl, aminoisobutyl, aminotertbutyl, aminopentyl, aminoisopentyl and aminoneopentyl.

86. The method of Claim 71 wherein said first residue is provided by a biphenylalkyl 1H-substituted-
 15 1,2,4-triazole angiotensin II antagonist compound containing a moiety convertible to a primary or secondary amino terminal moiety.

87. The method of Claim 86 wherein said moiety
 20 convertible to an amino terminal moiety is a carboxylic acid group reactable with an amino moiety of a diamino-terminated linker group to provide a terminal amino moiety which may then be further reacted with a carboxylic acid moiety of a compound providing said second residue so as to
 25 form a hydrolyzable amide bond.

88. The method of Claim 87 wherein said diamino-terminated linker group is a divalent radical of
 Formula III:


30




wherein each of R²⁰⁰ and R²⁰¹ may be independently selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, alkoxyalkyl, hydroxyalkyl, aralkyl, aryl, haloalkyl, amino, monoalkylamino, dialkylamino, cyanoamino, carboxyalkyl, alkylsulfinyl, alkylsulfonyl, arylsulfinyl and arylsulfonyl; and wherein n is zero or a number selected from three through seven, inclusive.

89. The method of Claim 88 wherein each of R²⁰⁰ and R²⁰¹ is hydrido.

90. The method of Claim 87 wherein said diamino-terminated linker group is a divalent radical of Formula IV:




wherein each of Q and T is one or more groups independently selected from



wherein each of R²⁰² through R²⁰⁵ is independently selected from hydrido, hydroxy, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl, alkoxy, aralkoxy, aryloxy, alkoxyalkyl, haloalkyl, hydroxyalkyl, halo, cyano, amino, monoalkylamino, dialkylamino, carboxy, carboxyalkyl, alkanoyl, alkenyl, cycloalkenyl and alkynyl.

91. The method of Claim 90 wherein said diamino-terminated linker group is a divalent radical of Formula V:



wherein each of R^{202} and R^{203} is independently selected
 5 from hydrido, hydroxy, alkyl, phenalkyl, phenyl, alkoxy, benzyloxy, phenoxy, alkoxyalkyl, hydroxyalkyl, halo, amino, monoalkylamino, dialkylamino, carboxy, carboxyalkyl and alkanoyl; and wherein each of p and q is a number
 independently selected from one through six, inclusive;
 10 with the proviso that when each of R^{202} and R^{203} is selected from halo, hydroxy, amino, monoalkylamino and dialkylamino, then the carbon to which R^{202} or R^{203} is attached in Formula V is not adjacent to a nitrogen atom of Formula V.


15

92. The method of Claim 91 wherein each of R^{202} and R^{203} is independently selected from hydrido, hydroxy, alkyl, alkoxy, amino, monoalkylamino, carboxy, carboxyalkyl and alkanoyl; and wherein each of p and q is a number
 20 independently selected from two through four, inclusive.

93. The method of Claim 92 wherein each of R^{202} and R^{203} is independently selected from hydrido, amino, monoalkylamino and carboxyl; and wherein each of p and q is
 25 independently selected from the numbers two and three.

94. The method of Claim 93 wherein each of R^{202} and R^{203} is hydrido; and wherein each of p and q is two.

95. The method of Claim 87 wherein said diamino-terminated linker group is a divalent radical of Formula VI:



5

wherein each of R²¹⁴ through R²¹⁷ is independently selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, hydroxyalkyl, alkoxyalkyl, aralkyl, aryl, haloalkyl, amino, monoalkylamino, dialkylamino, cyanoamino, carboxyalkyl, alkylsulfinyl, alkylsulfonyl, arylsulfinyl and arylsulfonyl; and wherein p is a number selected from one through six inclusive.

15

96. The method of Claim 95 wherein each of R²¹⁴ and R²¹⁵ is hydrido; wherein each of R²¹⁶ and R²¹⁷ is independently selected from hydrido, alkyl, phenalkyl, phenyl, alkoxyalkyl, hydroxyalkyl, haloalkyl and carboxyalkyl; and wherein p is two or three.

20

97. The method of Claim 96 wherein each of R²¹⁴ and R²¹⁵ is hydrido; wherein each of R²¹⁶ and R²¹⁷ is independently selected from hydrido and alkyl; and wherein p is two.

25

98. The method of Claim 97 wherein each of R²¹⁴, R²¹⁵, R²¹⁶ and R²¹⁷ is hydrido; and wherein p is two.

99. The method of Claim 79 wherein said angiotensin II antagonist compound is selected from the group consisting of

30

methyl 4'-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylate;

4'-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-

35

biphenyl]-2-carboxylic acid;

- 4'-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid, hydrazide;
4'-[(5-butyl-3-chloro-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 5 4'-[(3-butyl-5-chloro-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-propyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 10 4'-[(5-butyl-3-isopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-secbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-isobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 15 4'-[(5-butyl-3-tertbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-pentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-isopentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 20 4'-[(5-butyl-3-cyclohexyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-cyclohexylmethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 25 4'-[[5-butyl-3-(2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-cyclohexanoyl-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 30 4'-[[5-butyl-3-(1-oxo-2-cyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-phenyl-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
- 35 4'-[(5-butyl-3-phenylmethyl-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-butyl-3-(2-phenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(5-butyl-3-benzoyl-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;

- 4'-[[5-butyl-3-(1-oxo-2-phenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-butyl-3-(1,1-dimethoxypropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
5 4'-[[5-butyl-3-(1,1-dimethoxybutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-butyl-3-(1-oxopropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-butyl-3-(1-oxobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
10 4'-[[5-butyl-3-(1-oxopentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-butyl-3-(1,1-difluoroethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
15 4'-[[5-butyl-3-(1,1-difluoropropyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[5-butyl-3-(1,1-difluorobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[[3-butyl-5-(1,1-difluorobutyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
20 4'-[[5-butyl-3-(1,1-difluoropentyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(3,5-dipropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
25 4'-[(3,5-isopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(3,5-disecbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(3,5-diisobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
30 4'-[(3,5-ditertbutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
4'-[(3,5-dipentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
35 4'-[(3,5-diisopentyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-carboxylic acid;
5-[4'-[(5-butyl-3-amino-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;

- 5-[4'-[(5-butyl-3-aminomethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-butyl-3-aminoethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5 5-[4'-[(5-butyl-3-aminopropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-butyl-3-aminobutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminophenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
10 5-[4'-[[3-butyl-5-(4-aminophenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminophenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
15 5-[4'-[[3-butyl-5-(4-aminomethylphenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminomethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminoethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
20 5-[4'-[[3-butyl-5-(4-aminocyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminocyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
25 5-[4'-[[3-butyl-5-(4-aminocyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminomethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[[3-butyl-5-(4-aminomethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
30 5-[4'-[[3-butyl-5-(4-aminoethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-butyl-3-carboxy-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
35 5-[4'-[(5-butyl-3-carboxymethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-butyl-3-carboxyethyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;

5-[4'-[(5-butyl-3-carboxypropyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(5-butyl-3-carboxybutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5 5-[4'-[(3-butyl-5-(4-carboxyphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(3-butyl-5-(4-carboxyphenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(3-butyl-5-(4-carboxyphenylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
10 5-[4'-[(3-butyl-5-(4-carboxymethylphenylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(3-butyl-5-(4-carboxymethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
15 5-[4'-[(3-butyl-5-(4-carboxyethylphenyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(3-butyl-5-(4-carboxycyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(3-butyl-5-(4-carboxycyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
20 5-[4'-[(3-butyl-5-(4-carboxycyclohexylethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
5-[4'-[(3-butyl-5-(4-carboxymethylcyclohexylmethyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole;
25 tetrazole;
5-[4'-[(3-butyl-5-(4-carboxymethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole; and
5-[4'-[(3-butyl-5-(4-carboxyethylcyclohexyl)-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]-1H-tetrazole.

30

100. The method of Claim 89 wherein said renal-selective conjugate is N-acetylglutamic acid, 5-[[4'-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl][1,1'-biphenyl]-2-yl]]carbonylhydrazide.

35

101. The method of Claim 85 wherein said renal-selective conjugate is N²-acetyl-N-[[5-butyl-1-[[2'-(1H-

tetrazol-5-yl)[1,1'-biphenyl]-4-yl)methyl]-1H-1,2,4-triazol-3-yl)methyl]-L-glutamine.

102. The method of Claim 89 wherein said renal-
5 selective conjugate is N-acetyl-L-glutamic acid, 5-[5-butyl-1-[[2'-(1H-tetrazol-5-yl)[1,1'-biphenyl]-4-yl)methyl-1H-1,2,4-triazol-3-yl]acetylhydrazide.

103. The method of Claim 69 wherein said
10 circulatory disorder is a hypertensive-related disorder.

104. The method of Claim 103 wherein said hypertensive-related disorder is chronic hypertension.

15 105. The method of Claim 69 wherein said circulatory disorder is a sodium-retaining disorder.

106. The method of Claim 105 wherein said sodium-retaining disorder is congestive heart failure.
20

107. The method of Claim 105 wherein said sodium-retaining disorder is cirrhosis.

108. The method of Claim 105 wherein said
25 sodium-retaining disorder is nephrosis.




FIG. 1




FIG. 2




FIG. 3